

## BOUNDARY CONTROL OF THE MAXWELL DYNAMICAL SYSTEM: LACK OF CONTROLLABILITY BY TOPOLOGICAL REASONS

MIKHAIL BELISHEV<sup>1</sup> AND ALEKSANDR GLASMAN<sup>2</sup>

**Abstract.** The paper deals with a boundary control problem for the Maxwell dynamical system in a bounded domain  $\Omega \subset \mathbf{R}^3$ . Let  $\Omega^T \subset \Omega$  be the subdomain filled by waves at the moment  $T$ ,  $T_*$  the moment at which the waves fill the whole of  $\Omega$ . The following effect occurs: for small enough  $T$  the system is approximately controllable in  $\Omega^T$  whereas for larger  $T < T_*$  a lack of controllability is possible. The subspace of unreachable states is of finite dimension determined by topological characteristics of  $\Omega^T$ .

**AMS Subject Classification.** 93B05, 35B37, 35Q60, 78A25, 93C20.

Received June 11, 1999. Revised December 30, 1999.

### INTRODUCTION

Let  $\Omega \subset \mathbf{R}^3$  be a bounded domain with a smooth boundary  $\Gamma$ . We consider the Maxwell system

$$\begin{aligned} \varepsilon e_t &= \operatorname{rot} h; & \mu h_t &= -\operatorname{rot} e & \text{in } \Omega \times (0, T); \\ \operatorname{div} \varepsilon e &= 0, & \operatorname{div} \mu h &= 0 & \text{in } \Omega; \\ e|_{t=0} &= 0, & h|_{t=0} &= 0; \\ \nu \times e|_{\Gamma \times [0, T]} &= f, \end{aligned}$$

where  $\varepsilon, \mu$  are smooth positive scalar functions (permeabilities) given in  $\overline{\Omega}$ ,  $\nu$  is a normal on  $\Gamma$ ,  $f$  is a boundary control; let  $\{e^f(x, t), h^f(x, t)\}$  be a solution (wave).

Permeabilities determine the velocity  $c = (\varepsilon\mu)^{1/2}$  and the optical metric

$$d\tau^2 = \frac{|dx|^2}{c^2},$$

which turns  $\Omega$  into a Riemannian manifold; we denote  $\operatorname{dist}_c$  the corresponding distance. Let

$$\Omega^T := \{x \in \Omega \mid \operatorname{dist}_c(x, \Gamma) < T\}, \quad T > 0$$

---

*Keywords and phrases:* Maxwell's dynamical system, boundary control, unreachable states, topology of a domain.

<sup>1</sup> Saint-Petersburg Department of Steklov Mathematical Institute, Fontanka 27, Saint-Petersburg 191011, Russia;  
e-mail: [belishev@bel.pdmi.ras.ru](mailto:belishev@bel.pdmi.ras.ru)

Supported by RFBR, grant 98-01-00314.

<sup>2</sup> Saint-Petersburg State University, Saint-Petersburg, Russia.

Supported by RFBR, grant 99-01-00107.

be a near-boundary layer of optical thickness  $T$ ; the surface

$$\Gamma^T := \{x \in \Gamma \mid \text{dist}_c(x, \Gamma) = T\}$$

is an inner component of  $\partial\Omega^T$ . The magnitude

$$T_* := \inf\{T > 0 \mid \Omega^T = \Omega\}$$

coincides with time needed for waves moving into  $\Omega$  from  $\Gamma$  to fill the whole of the domain.

Introduce the (electric) reachable set

$$\mathcal{E}^T := \{e^f(\cdot, T) \mid f \in L_2((0, T); H^1(\Gamma)), f \cdot \nu = 0 \text{ on } \Gamma\}$$

( $H^1(\dots)$  is the Sobolev class); let

$$J^T := \left\{ y \in L_2(\Omega) \mid \text{div } \varepsilon y = 0 \text{ in } \Omega, \text{ supp } y \subset \overline{\Omega}^T \right\}$$

be the space of  $\varepsilon$ -solenoidal fields localized in  $\overline{\Omega}^T$ . By finiteness of  $c$ , the embedding

$$\mathcal{E}^T \subset J^T, \quad T > 0$$

occures. The main question under consideration is a density of this embedding. Our results are the following.

Let us say that  $\Omega^T$  satisfies the EP-condition (existence of potential) if any cycle (simple smooth closed curve) lying in  $\Omega^T$  may be continuously deformed into a cycle lying on  $\Gamma$ .

**Theorem 1.** *If  $\Omega^T$  satisfies the EP-condition the equality*

$$\text{clos } \mathcal{E}^T = J^T \tag{*}$$

*holds.*

In particular, for small enough  $T$  relation (\*) is valid, *i.e.* the electric component of the Maxwell system is approximately controllable.

If the EP-condition is violated the unreachable subspace

$$\mathcal{N}^T = J^T \ominus \mathcal{E}^T,$$

turns out to be nontrivial, *i.e.* the system is not controllable. The subspace  $\mathcal{N}^T$  is of a finite dimension determined by topological characteristics of  $\Omega^T$ . For example, if  $\Omega$  is homeomorphic to a ball and  $\Omega \setminus \overline{\Omega}^T$  is homeomorphic to a ball with  $n$  handles then  $\dim \mathcal{N}^T = n$ .

A lack of controllability described above is of purely topological nature: it is not connected with a presence of real obstacles in  $\Omega$ . In particular, if the system is not controllable at the moment  $t = T_0$ , however, the equality (\*) may be restored later for some  $T > T_0$ .

## 1. DOMAINS AND SPACES

Let  $\Omega \subset \mathbf{R}^3$  be a bounded domain with a boundary  $\Gamma \in C^\infty$ ,  $\varepsilon, \mu \in C^\infty(\overline{\Omega})$  strictly positive functions (permeabilities); denote  $c := (\varepsilon\mu)^{-1/2}$ .

Equip  $\Omega$  with the optical metric

$$d\tau^2 = \frac{|dx|^2}{c^2};$$

let  $\text{dist}_c$  be the corresponding distance; introduce the eikonal  $\tau(x) := \text{dist}_c(x, \Gamma)$ ,  $x \in \overline{\Omega}$ . The eikonal determines an increasing family of subdomains

$$\Omega^T := \{x \in \Omega \mid \tau(x) < T\}, \quad T > 0$$

and the level surfaces

$$\Gamma^T := \{x \in \Omega \mid \tau(x) = T\}, \quad T \geq 0$$

( $\Gamma^0 = \Gamma$ ); denote

$$T_* := \inf \{T > 0 \mid \Omega^T = \Omega\} = \max_{\Omega} \tau(\cdot).$$

Let us introduce spaces and classes of  $\mathbf{R}^3$ -valued functions (fields) used in the paper:

the Sobolev classes  $H^s(\dots)$ ;

the space of  $\varepsilon$ -solenoidal fields  $J := \{y \in L_{2,\varepsilon}(\Omega) \mid \text{div } \varepsilon y = 0 \text{ in } \Omega\}$  (with measure  $\varepsilon dx$ );

the subspace  $J^T := \{y \in J \mid \text{supp } y \subset \overline{\Omega}^T\}$  of fields localized in  $\overline{\Omega}^T$ ;

the class  $J_+ := J \cap H^1(\Omega)$  (with  $H^1$ -topology) and its dual  $J_- := (J_+)'$  with respect to  $J$ ;

the space of tangent fields  $\mathcal{T} := \{g \in L_2(\Gamma) \mid \nu \cdot g = 0 \text{ on } \Gamma\}$  ( $\nu$  is a normal);

the class  $\mathcal{T}_+ := \mathcal{T} \cap H^1(\Gamma)$  (with  $H^1$ -topology) and its dual  $\mathcal{T}_- := (\mathcal{T}_+)'$  with respect to  $L_2(\Gamma)$ ;

the space of controls  $\mathcal{F}^T := L_2([0, T]; \mathcal{T})$ ;

the class  $\mathcal{F}_+^T := L_2((0, T); \mathcal{T}_+)$  and its dual  $\mathcal{F}_-^T := (\mathcal{F}_+^T)' = L_2((0, T); \mathcal{T}_-)$  with respect to  $\mathcal{F}^T$ .

## 2. THE MAXWELL SYSTEM WITH BOUNDARY CONTROL. ELECTRIC SUBSYSTEM

Denote  $Q^T := \Omega \times (0, T)$ ,  $\Sigma^T := \Gamma \times [0, T]$  and consider the system

$$\varepsilon e_t = \text{rot } h, \quad \mu h_t = -\text{rot } e \quad \text{in } Q^T; \tag{2.1}$$

$$e|_{t=0} = 0, \quad h|_{t=0} = 0; \tag{2.2}$$

$$\nu \times e|_{\Sigma^T} = f, \tag{2.3}$$

with (electric) boundary control  $f$ ; let  $\{e^f(x, t), h^f(x, t)\}$  be its solution. Note that (2.1, 2.2) imply

$$\text{div } \varepsilon e = 0, \quad \text{div } \mu h = 0 \quad \text{in } \Omega.$$

For  $f \in \mathcal{F}_+^T$  problem (2.1–2.3) is uniquely solvable in an appropriate class (see [7, 10]). The well known fact is that solutions (waves) propagate with velocity  $c$ :

$$\text{supp } \{e^f, h^f\} \subset \{(x, t) \in \overline{Q}^T \mid t \geq \tau(x)\}. \tag{2.4}$$

The electric component satisfies

$$e_{tt} + \frac{1}{\varepsilon} \text{rot} \frac{1}{\mu} \text{rote} = 0 \quad \text{in } Q^T; \tag{2.5}$$

$$e|_{t=0} = e_t|_{t=0} = 0 \quad \text{in } \Omega; \tag{2.6}$$

$$\nu \times e|_{\Sigma^T} = f. \tag{2.7}$$

For  $f \in \mathcal{F}_+^T$  the inclusion  $e^f \in C([0, T]; J)$  holds, and the map  $f \rightarrow e^f$  is continuous in corresponding norms; this property ensures a continuity of the map  $W^T : f \rightarrow e^f(\cdot, T)$  from  $\mathcal{F}_+^T$  into  $J$ .

**Theorem 2.** *For times  $T < T_*$  the map  $W^T$  is injective.*

*Proof.* Choose  $g \in \text{Ker}W^T$ ; let  $e^g, h^g$  be the solution of (2.1–2.3). Consider the extensions:

$$e(\cdot, t) := \begin{cases} 0 & -\infty < t < 0; \\ e^g(\cdot, t) & 0 \leq t < T; \\ -e^g(\cdot, 2T - t) & T \leq t < 2T; \\ 0 & 2T \leq t < \infty \end{cases}$$

and

$$h(\cdot, t) := \begin{cases} 0 & -\infty < t < 0; \\ h^g(\cdot, t) & 0 \leq t < T; \\ h^g(\cdot, 2T - t) & T \leq t < 2T; \\ 0 & 2T \leq t < \infty. \end{cases}$$

By virtue of  $e^g(\cdot, T) = 0$ , extending by oddness one doesn't violate a continuity of  $e^g$  and the pair  $\{e, h\}$  turns out to be a solution of the system

$$\varepsilon e_t = \text{rot } h, \quad \mu h_t = -\text{rot } e \quad \text{in } \Omega \times (-\infty, \infty). \quad (2.8)$$

Relation (2.4) implies  $\text{supp } \{e(\cdot, t), h(\cdot, t)\} \subset \overline{\Omega}^T$  for any  $t$  that leads to

$$e = 0, \quad h = 0 \quad \text{in } (\Omega \setminus \overline{\Omega}^T) \times (-\infty, \infty). \quad (2.9)$$

Applying the Fourier transform on time to (2.8, 2.9) we get

$$ik\varepsilon \tilde{e}(\cdot, k) = \text{rot } \tilde{h}(\cdot, k), \quad -ik\mu \tilde{h}(\cdot, k) = \text{rot } \tilde{e}(\cdot, k) \quad \text{in } \Omega; \quad (2.10)$$

$$\tilde{e}(\cdot, k) = 0, \quad \tilde{h}(\cdot, k) = 0 \quad \text{in } \Omega \setminus \overline{\Omega}^T \quad (2.11)$$

for all  $k \in (-\infty, \infty)$ . By virtue of  $\text{div } \varepsilon \tilde{e}(\cdot, k) = 0$ ,  $\text{div } \mu \tilde{h}(\cdot, k) = 0$ , system (2.10) turns out to be elliptic, its solution vanishing on a nonvoid open subset (see (2.11)). By known uniqueness theorem (see [11], Th. 8.17) the solution vanishes in  $\Omega$  identically that implies  $\tilde{e} = 0$ , then  $e = 0$ ,  $e^g = 0$ , and, finally,  $g = 0$ . Thus,  $\text{Ker } W^T = \{0\}$ ; the theorem is proved.

A simple generalization of the proof enables to obtain the following interesting result. Let us say that a subset  $\omega \subset \Omega^T$  belongs to the class  $\mathcal{D}^T$  if  $\text{dist}_c(\omega, \partial\Omega^T) > 0$ , i.e.  $\omega$  is separated from  $\Gamma \cup \Gamma^T$ , and the (open) set  $\Omega^T \setminus \overline{\omega}$  is connected. Put also  $\emptyset \in \mathcal{D}^T$  by definition.

**Lemma 1.** *Let  $T < T_*$ ,  $\{e^f, h^f\}$  satisfy (2.1–2.3) for  $f \in \mathcal{F}_+^T$ . If  $\text{supp } e^f(\cdot, T) \in \mathcal{D}^T$  then  $f = 0$  and  $e^f = 0$ ,  $h^f = 0$ .*

The analogous result for the scalar wave equation was established in [1]. Notice that Theorem 2 is a simple corollary of Lemma 1.

### 3. BOUNDARY CONTROL PROBLEM

Let us return back to the system (2.1–2.3). As Theorem 2 shows, for times  $T < T_*$  electric component  $e^f(\cdot, T)$  determines uniquely control  $f$  which, in turn, determines magnetic component  $h^f(\cdot, T)$ . Therefore, managing  $f$  one can't control both of the components simultaneously. Thus, in the case  $T < T_*$ , the following statement of the boundary control problem (BCP) turns out to be natural: given  $y \in J^T$  to find control  $f \in \mathcal{F}_+^T$  such that the equality

$$e^f(\cdot, T) = y$$

holds. By virtue of Theorem 2 the BCP has no more than one solution.

The operator  $W^T : \mathcal{F}^T \rightarrow J$ ,  $\text{Dom } W^T = \mathcal{F}_+^T$ ,  $W^T f := e^f(\cdot, T)$  is well defined due to Section 2; it is injective for  $T < T_*$ . By virtue of (2.4),  $W^T$  acts into the subspace  $J^T$ . The set

$$\mathcal{E}^T := \text{Ran } W^T = \{e^f(\cdot, T) \mid f \in \mathcal{F}_+^T\}$$

is said to be reachable (at the moment  $t = T$ ). The goal of the paper is to treat the embedding  $\mathcal{E}^T \subset J^T$ .

In the case of  $T < T_*$  Lemma 1 shows that any nonzero  $y \in J^T : \text{supp } y \in \mathcal{D}^T$  doesn't belong to  $\mathcal{E}^T$ . Thus, the set  $J^T \setminus \mathcal{E}^T$  is rich enough and the equality  $\mathcal{E}^T = J^T$  (exact controllability) certainly doesn't hold. This raises the question of whether the equality  $\text{clos } \mathcal{E}^T = J^T$  (approximate controllability) holds, which is main subject of the paper.

#### 4. DUAL SYSTEM

The system

$$\varepsilon \varphi_t = \text{rot } \psi, \quad \mu \psi_t = -\text{rot } \varphi \quad \text{in } Q^T; \quad (4.1)$$

$$\varphi|_{t=T} = y, \quad \psi|_{t=T} = 0; \quad (4.2)$$

$$\nu \times \varphi|_{\Sigma^T} = 0; \quad (4.3)$$

is called dual to system (2.1–2.3); let  $\varphi = \varphi^y(x, t)$ ,  $\psi = \psi^y(x, t)$  be its solution. The following is something of the properties of  $\{\varphi^y, \psi^y\}$  (see [9, 10]):

- (i) for  $y \in J$  one has  $\varphi^y \in C([0, T]; J)$ ;  $\psi^y \in C([0, T]; L_2(\Omega))$ ;  $\text{div } \mu \psi^y = 0$ ;  $\nu \cdot \psi^y = 0$  on  $\Sigma^T$ ;
- (ii) the map  $y \rightarrow \nu \cdot \psi^y|_{\Sigma^T}$  acts continuously from  $J$  into  $\mathcal{F}_-^T$ ;
- (iii) by finiteness of velocity of wave propagation, solution  $\{\varphi^y, \psi^y\}$  in the subdomain  $\{(x, t) \in Q^T \mid t > \tau(x)\}$  is determined by  $y|_{\Omega^T}$  (doesn't depend on  $y|_{\Omega \setminus \Omega^T}$ );
- (iv) the duality relation

$$(e^f(\cdot, T), y)_J = -(f, \psi^y|_{\Sigma^T})_{\mathcal{F}^T} \quad (4.4)$$

holds for any  $f \in \mathcal{F}_+^T$ ,  $y \in J$ .

#### 5. UNREACHABLE STATES

The subspace

$$\mathcal{N}^T := J^T \ominus \text{clos } \mathcal{E}^T$$

is said to be unreachable. To describe  $\mathcal{N}^T$  let us introduce the set  $\mathcal{N}_*^T$  of  $y \in J^T$  such that:

- 1)  $y$  is  $C^\infty$ -smooth in  $\Omega^T \cup \Gamma$ ;
- 2)  $\nu \times y = 0$  on  $\Gamma$ ;
- 3)  $\text{rot } y = 0$  in  $\Omega^T$ .

**Theorem 3.** For any  $T > 0$  the equality

$$\mathcal{N}^T = \mathcal{N}_*^T \quad (5.0)$$

holds.

*Proof.* (i) Choose  $y \in \mathcal{N}_*^T$ ; As is easy to check, the pair  $\{y(x), 0\}$  satisfies (4.1–4.3) for  $t > \tau(x)$  (see (iii), Sect. 4). Therefore, by uniqueness of solution of the dual system one has

$$\varphi^y(x, t) = y(x), \quad \psi^y(x, t) = 0 \quad \text{in } \{(x, t) \in Q^T \mid t > \tau(x)\};$$

in particular,  $\psi^y = 0$  holds on  $\Sigma^T$ . Duality (4.4) leads to  $(e^f(\cdot, T), y)_J = 0$  for any  $f \in \mathcal{F}_+^T$ ; hence  $y \perp \mathcal{E}^T$ , i.e.  $y \in \mathcal{N}^T$ , and we get  $\mathcal{N}_*^T \subset \mathcal{N}^T$ . To prove the theorem one needs to check the opposite inclusion  $\mathcal{N}_*^T \supset \mathcal{N}^T$ .

(ii) Choose  $y \in \mathcal{N}^T$ ; let  $\{\varphi^y, \psi^y\}$  be the corresponding solution of (4.1–4.3). Boundary condition (4.3), duality (4.4) and property (i), Section 4 lead to

$$\nu \times \varphi^y = 0, \quad \psi^y = 0 \quad \text{on } \Sigma^T, \quad (5.1)$$

the latter equality being understood in accordance with (ii), Section 4.

Extending the solution as follows

$$\varphi(\cdot, t) := \begin{cases} \varphi^y(\cdot, t), & 0 \leq t < T, \\ \varphi^y(\cdot, 2T - t), & T \leq t < 2T; \end{cases}$$

$$\psi(\cdot, t) := \begin{cases} \psi^y(\cdot, t), & 0 \leq t < T, \\ -\psi^y(\cdot, 2T - t), & T \leq t < 2T; \end{cases}$$

and taking into account (5.1) one can check that  $\varphi, \psi$  satisfy

$$\varepsilon \varphi_t = \text{rot } \psi, \quad \mu \psi_t = -\text{rot } \varphi, \quad \text{in } Q^{2T}; \quad (5.2)$$

$$\nu \times \varphi = 0, \quad \psi = 0 \quad \text{on } \Sigma^{2T}. \quad (5.3)$$

(iii) To deal with classical solutions we apply smoothing with respect to time. Choose a scalar function  $\chi \in C_0^\infty(-\infty, \infty)$ :

$$\chi(-t) = \chi(t), \quad \chi(t) \geq 0, \quad \text{supp } \chi \subset [-1, 1], \quad \int_{-1}^1 \chi(t) dt = 1,$$

and denote  $\chi_\delta(t) := \frac{1}{\delta} \chi(\frac{t}{\delta})$  ( $\delta > 0$ ), so that  $\chi_\delta$  converges to the Dirac function as  $\delta$  tends to zero. The vector valued functions

$$\varphi^\delta(\cdot, t) := \chi_\delta(t) * \varphi(\cdot, t), \quad \psi^\delta(\cdot, t) := \chi_\delta(t) * \psi(\cdot, t)$$

are defined in  $Q_\delta^{2T} := \Omega \times (\delta, 2T - \delta)$  and satisfy

$$\varepsilon \varphi_t^\delta = \text{rot } \psi^\delta, \quad \mu \psi_t^\delta = -\text{rot } \varphi^\delta, \quad \text{in } Q_\delta^{2T}; \quad (5.4)$$

$$\nu \times \varphi^\delta = 0 \quad \text{on } \Sigma_\delta^{2T} \quad (5.5)$$

$$\psi^\delta = 0 \quad \text{on } \Sigma_\delta^{2T} \quad (5.6)$$

where  $\Sigma_\delta^{2T} := \Gamma \times [\delta, 2T - \delta]$ . A peculiar feature of the Maxwell system is that time smoothing leads to smoothing with respect to space variables. This may be justified, for instance, by means of the Fourier method expanding  $\varphi(\cdot, t)$ ,  $\psi(\cdot, t)$  over the eigenbasis of the Maxwell operator associated with system (2.1–2.3) (see [11]). Smoothed solutions turns out to be classical:  $\varphi^\delta, \psi^\delta \in C^\infty(\overline{Q}_\delta^{2T})$ .

(iv) A simple fact of the vector analysis is that relation (5.6) implies

$$\nu \cdot \text{rot } \psi^\delta = 0 \quad \text{on } \Sigma_\delta^{2T}. \quad (5.7)$$

Multiplying (5.4) by  $\nu$  on  $\Gamma$  we get

$$\nu \cdot \varphi_t^\delta = \frac{1}{\varepsilon} \nu \cdot \text{rot } \psi^\delta = 0 \quad \text{on } \Sigma_\delta^{2T} \quad (5.8)$$

in view of (5.7). Relations (5.5, 5.8) lead to

$$\varphi_t^\delta = 0 \quad \text{on } \Sigma_\delta^{2T}; \quad (5.9)$$

(5.9) and (5.4) give

$$\operatorname{rot} \psi^\delta = 0 \quad \text{on } \Sigma_\delta^{2T}. \tag{5.10}$$

We omit the proof of the following auxiliary result.

**Proposition 5.1.** *If  $\eta \in C^1(\overline{\Omega}^\xi)$  satisfies  $\operatorname{div} \mu \eta = 0$  in  $\Omega^\xi$  and  $\eta = \operatorname{rot} \eta = 0$  on  $\Gamma$  then  $\frac{\partial \eta}{\partial \nu} = 0$  on  $\Gamma$ .*

The equality

$$\frac{\partial \psi^\delta}{\partial \nu} = 0 \quad \text{on } \Sigma_\delta^{2T} \tag{5.11}$$

follows from (5.6, 5.10) and the proposition.

(v) Separating  $\psi^\delta$  in (5.4) one obtains the equation

$$\psi_{tt}^\delta + \frac{1}{\mu} \operatorname{rot} \frac{1}{\varepsilon} \operatorname{rot} \psi^\delta = 0 \quad \text{in } Q_\delta^{2T}$$

that may be written in the form

$$\psi_{tt}^\delta - \frac{1}{c^2} \Delta \psi^\delta + \dots = 0 \quad \text{in } Q_\delta^{2T} \tag{5.12}$$

taking into account  $\operatorname{div} \mu \psi^\delta = 0$  (the low order terms are omitted). So  $\psi^\delta$  turns out to be a solution of the hyperbolic system (5.12) with zero Cauchy data (5.6, 5.11) on the time-like noncharacteristic hypersurface  $\Sigma_\delta^{2T}$ . Applying the vectorial version [8] of the Holmgren-John-Tataru uniqueness theorem [16] and using the Russell's scheme [14] (see also [2]) one can conclude that  $\psi^\delta$  is continued by zero from  $\Sigma_\delta^{2T}$  into the subdomain

$$K_\delta^{2T} := \{ (x, t) \in Q_\delta^{2T} \mid \tau(x) + \delta < t < 2T - \tau(x) - \delta \}$$

bounded by characteristic surfaces:

$$\psi^\delta = 0 \quad \text{in } K_\delta^{2T}.$$

Therefore, by (5.4) we get

$$\operatorname{rot} \varphi^\delta = 0 \quad \text{in } K_\delta^{2T}. \tag{5.13}$$

(vi) As  $\delta \rightarrow 0$ , the convergence  $\varphi^\delta \rightarrow \varphi$  occurs in  $C([\delta_0, 2T - \delta_0]; J)$  for any fixed  $\delta_0 > 0$ ; in particular, one has  $\varphi^\delta(\cdot, T) \rightarrow \varphi(\cdot, T) = y$  in  $J$ .

Choose any field  $\rho \in C^\infty(\overline{\Omega})$ ,  $\operatorname{supp} \rho \subset \overline{\Omega}^\xi$  for  $\xi < T$ . By virtue of (5.5) and (5.13) the equalities

$$0 = (\operatorname{rot} \varphi^\delta(\cdot, T), \rho)_{L_2(\Omega)} = (\varphi^\delta, \operatorname{rot} \rho)_{L_2(\Omega)} \tag{5.14}$$

are valid. The limit passage  $\delta \rightarrow 0$  gives

$$(y, \operatorname{rot} \rho)_{L_2(\Omega)} = 0,$$

which means that  $y$  satisfies

$$\operatorname{rot} y = 0 \text{ in } \Omega^T, \quad \nu \times y = 0 \text{ on } \Gamma \tag{5.15}$$

in a weak sense (see *e.g.* [7]). Since the boundary  $\Gamma$  is smooth (5.15) and  $\operatorname{div} \varepsilon y = 0$  lead to  $C^\infty$ -smoothness of  $y$  in  $\Omega^T$  up to  $\Gamma$  by standard elliptic theory. Thus, we get  $y \in \mathcal{N}^T$  that proves the theorem.

6. APPROXIMATE CONTROLLABILITY

We continue to study the subspace  $\mathcal{N}^T$ . Let us say that subdomain  $\Omega^T$  satisfies the EP-condition (existence of potential) if any cycle (a simple smooth closed curve) in  $\Omega^T$  may be continuously deformed into a cycle lying on  $\Gamma$ .

**Theorem 4.** *If time  $T > 0$  is such that  $\Omega^T$  satisfies the EP-condition then*

$$\mathcal{N}^T = \{0\}.$$

*Proof.* Choose  $y \in \mathcal{N}^T$ . In accordance with Theorem 3 one has

$$\operatorname{rot} y = 0 \text{ in } \Omega^T; \quad \nu \times y = 0 \text{ on } \Gamma. \tag{6.1}$$

Due to the EP-condition (6.1) ensures existence of a scalar function (potential)  $p$ , such that

$$\nabla p = y \text{ in } \Omega^T, \quad p = 0 \text{ on } \Gamma; \tag{6.2}$$

the inclusion  $y \in J^T$  implies

$$\operatorname{div} \varepsilon \nabla p = 0 \text{ in } \Omega^T. \tag{6.3}$$

Since  $\operatorname{supp} y \subset \overline{\Omega}^T$  and  $\operatorname{div} \varepsilon y = 0$  in the whole of  $\Omega$ , the equality  $\nu \cdot y = 0$  holds on  $\Gamma^T$  in appropriate (weak) sense that implies

$$\frac{\partial p}{\partial \nu} = 0 \text{ on } \Gamma^T. \tag{6.4}$$

Let  $B_r(x_0) := \{x \in \overline{\Omega} \mid \operatorname{dist}_c(x, x_0) \leq r\}$  be a “ball”; representing

$$\Omega^T = \bigcup_{\gamma \in \Gamma} B_T(\gamma)$$

one can easily show that subdomain  $\Omega^T$  satisfies the cone condition (see *e.g.* [12]).

In this case the elliptic equation (6.3) has a unique solution  $p \in H^1(\Omega^T)$  satisfying boundary conditions (6.2, 6.4). Hence,  $p = 0$  and  $y = \nabla p = 0$  that proves the theorem.

As a corollary, we conclude: for time  $T > 0$  such that  $\Omega^T$  satisfies the EP-condition the relation

$$\operatorname{clos} \mathcal{E}^T = J^T \tag{6.5}$$

holds, *i.e.* electric subsystem of the Maxwell system turns out to be approximately controllable.

The EP-condition is realized for small enough  $T$  or in the case of  $\Omega \setminus \Omega^T = \bigcup_{j=1}^m B_j$  where  $B_i \cap B_j = \emptyset$ , each  $B_j$  is homeomorphic to a closed ball. In both cases approximate controllability occurs.

7. LACK OF CONTROLLABILITY

Comparing controllability properties of the Maxwell system with ones of the system governed by the wave equation (2) the following peculiarity could be noted. In the case of the wave equation, the Holmgren-John-Tataru uniqueness theorem gives the implication

$$y \in \{ \text{unreachable subspace} \} \Rightarrow y = 0,$$



whereas for system (2.1–2.3) it leads to conditions

$$\operatorname{rot} y = 0, \quad \operatorname{div} \varepsilon y = 0 \quad \text{in } \Omega^T; \tag{7.1}$$

$$\nu \times y = 0 \quad \text{on } \Gamma; \tag{7.2}$$

$$\nu \cdot y = 0 \quad \text{on } \Gamma^T. \tag{7.3}$$

The known fact is that, depending on topology of  $\Omega^T$ , problem (7.1–7.3) may have nontrivial solutions (see [6, 15]). Consider an example, assuming for simplicity  $\varepsilon = \mu = 1$ .

**Lemma 2.** *Let  $\Omega$  be homeomorphic to a ball,  $\Omega \setminus \Omega^T$  homeomorphic to a torus; then*

$$\dim \mathcal{N}^T = 1.$$

*Proof.* At first, let us note that the case under consideration is realizable. As example, one can consider a rotation body  $\Omega$  having dumbbell shaped cross-section and take large enough  $T$ .

Denote  $D := \{(x^1, x^2) \in \mathbf{R}^2 \mid (x^1)^2 + (x^2)^2 \leq 1\}$ ,  $S := \partial D$ ; let  $D \times S$  be the torus,  $\varphi$  a homeomorphism from  $D \times S$  onto  $\Omega \setminus \Omega^T$ . The curve  $\gamma := \varphi[\{(0, 0)\} \times S]$  is a cycle lying in  $\Omega \setminus \Omega^T$ .

Choose a cycle  $l \subset \Omega^T$  which envelopes  $\Omega \setminus \overline{\Omega}^T$  and can't be deformed into a cycle lying on  $\Gamma$ . Define the circulation of a field  $y$ :

$$C_l[y] := \int_l y \cdot dl.$$

The Biot-Savart field

$$b(x) := \alpha \int_\gamma \frac{(x - \xi) \times dl_\xi}{|x - \xi|^3}$$

( $\alpha = \text{const}$ ) satisfies (7.1) and has nonzero circulation; assume  $\alpha$  to be such that

$$C_l[b] = 1. \tag{7.4}$$

For any cycle  $\lambda \subset \Gamma$  one has

$$C_\lambda[b] = C_\lambda[b_\theta] = 0,$$

where  $b_\theta := b - (b \cdot \nu)\nu$ ; therefore, the field  $b_\theta$  has a surface potential on  $\Gamma$ : there exists smooth  $\pi$  such that  $\nabla_\Gamma \pi = b_\theta$  on  $\Gamma$ .

Find  $p$  as a (unique) solution of the Neumann-Dirichlet problem:

$$\begin{aligned} \Delta p &= 0 \quad \text{in } \Omega^T; \\ \frac{\partial p}{\partial \nu} &= b \cdot \nu \quad \text{on } \Gamma^T; \\ p &= \pi \quad \text{on } \Gamma. \end{aligned}$$

As is easy to check, the field

$$a := b - \nabla p$$

satisfies (7.1–7.3) and is nontrivial due to (7.4); thus,  $a \in \mathcal{N}^T$ , and  $\dim \mathcal{N}^T \geq 1$ .

Take  $y \in \mathcal{N}^T$  and denote  $g = y - C_l[y]a$ . For any cycle  $l$  lying in  $\Omega^T$  one has  $C_l[g] = 0$ . This, together with  $\operatorname{rot} g = 0$ , leads to existence of a potential  $q$ :  $\nabla q = g$  in  $\Omega^T$ . By (7.1–7.3), we obtain:

$$\begin{aligned} \Delta q &= 0 \quad \text{in } \Omega^T; \\ \frac{\partial q}{\partial \nu} &= 0 \quad \text{on } \Gamma^T; \\ q &= \text{const} \quad \text{on } \Gamma, \end{aligned}$$

that implies  $q = \text{const}$  and  $g = 0$ . Hence,  $y = C_l[y]a$  that leads to  $\dim \mathcal{N}^T = 1$ . The lemma is proved.

Note that idea of the proof is taken from [6].

The obtained result may be simply generalized as follows: if  $\Omega$  is homeomorphic to a ball whereas  $\Omega \setminus \Omega^T$  is homeomorphic to a ball with  $n$  handles then  $\dim \mathcal{N}^T = n$ .

Denote  $\mathcal{H}^T := \{y \in J^T \mid \text{rot } y = 0 \text{ in } \Omega^T\}$ ,  $\mathcal{G}^T := \{y \in J^T \mid y = \nabla p, p \in H^1(\Omega^T)\}$ . A simple analysis of the proof of Lemma 3 and its generalization mentioned above leads to the equality

$$\dim \mathcal{N}^T = \dim \mathcal{H}^T / \mathcal{G}^T;$$

relations of this kind are well-known in the Hodge Theory (see [15]).

In conclusion let us consider an example demonstrating a curious behaviour of subspace  $\mathcal{N}^T$ . Let  $\Omega_1$  be a rotation body with a dumbbell crosssection,  $\Omega_2$  a big ball,  $\Omega_3$  a narrow cylindrical channel connecting  $\Omega_1$  with  $\Omega_2$ , so that the domain  $\Omega := \Omega_1 \cup \Omega_2 \cup \Omega_3$  is homeomorphic to a ball.

- (i) If  $T$  is small enough,  $\Omega^T$  satisfies the EP-condition; hence,  $\mathcal{N}^T = \{0\}$ ;
- (ii) if  $T$  is such that  $\Omega_3 \subset \Omega^T$  (the channel is captured by waves) but  $\Omega \setminus \overline{\Omega}^T$  contains a torus lying in  $\Omega_1$ , we have  $\mathcal{N}^T \neq \{0\}$ ;
- (iii) for large enough  $T < T_*$  one has  $\Omega_1 \cup \Omega_3 \subset \Omega^T$  ( $\Omega_1$  and the channel are captured) whereas  $\Omega^T$  turns out to be homeomorphic to a spherical layer satisfying the EP-condition; hence,  $\mathcal{N}^T = \{0\}$  holds again.

## 8. REMARKS AND ACKNOWLEDGMENTS

- (i) The version of the BCP studied in the paper differs from traditional ones (see *e.g.* [9, 13, 17]). A reason of our interest is that it is the version which works in an approach to the inverse problems based upon their relations to the boundary control theory (the BC-method [2, 3, 5]).
- (ii) Lack of controllability discussed above was first noticed in [4] and mentioned in [3] in connection with the inverse problem for system (2.1–2.3): the presence of nontrivial  $\mathcal{N}^T$  creates complications there.
- (iii) We are grateful to our colleagues for fruitful discussions and kind help: control problems for the Maxwell system were discussed with C. Bardos; S. Kichenassamy explains the relationship between problem (7.1–7.3) and the Hodge theory.
- (iv) We would like to thank Referee 1 for very useful criticism: the paper has been thoroughly revised under his recommendations.
- (v) In the paper [17] by N. Weck an analogous effect (lack of controllability) is exhibited and studied. The author deals with more delicate problem of exact controllability, a description of an unreachable subspace being given in natural topological terms (the Betti numbers of  $\Omega$ ).

## REFERENCES

- [1] S. Avdonin, M. Belishev and S. Ivanov, The controllability in the filled domain for the multidimensional wave equation with a singular boundary control. *J. Math. Sci.* **83** (1997).
- [2] M.I. Belishev, Boundary control in reconstruction of manifolds and metrics (the BC-method). *Inverse Problems* **13** (1997) R1–R45. <http://www.iop.org/Journals/ip/>.
- [3] M. Belishev and A. Glasman, Boundary control and inverse problem for the dynamical maxwell system: the recovering of velocity in regular zone. *Preprint CMLA ENS Cachan* (1998) 9814. <http://www.cmla.ens-cachan.fr>
- [4] M. Belishev and A. Glasman, Visualization of waves in the Maxwell dynamical system (The BC-method). Preprint POMI (1997) 22. <http://www.pdmi.ras.ru/preprint/1997/>
- [5] M. Belishev, V. Isakov, L. Pestov and V. Sharafutdinov, On reconstruction of gravity field *via* external electromagnetic measurements. *Preprint PDMI* (1999) 10. <http://www.pdmi.ras.ru/preprint/1999/10-99.ps.gz>.
- [6] E.B. Bykhovskii and N.V. Smirnov, On an orthogonal decomposition of the space of square-summable vector- functions and operators of the vector analysis. *Proc. Steklov Inst. Math.* **59** (1960) 5–36, in Russian.
- [7] G. Duvaut and J.L. Lions, *Les inéquations en mécanique et en physique*, Vol. 21 of *Travaux et recherches mathématiques*. Paris: Dunod. XX (1972).
- [8] M. Eller, V. Isakov, G. Nakamura and D. Tataru, Uniqueness and stability in the Cauchy Problem for Maxwell and elasticity systems. *Nonlinear Partial Differential Equations and their applications. College de France Seminar. XIV* (1999) to appear.

- [9] J. Lagnese, Exact boundary controllability of Maxwell's equations in a general region. *SIAM J. Control Optim.* **27** (1989) 374–388.
- [10] I. Lasiecka and R. Triggiani, Recent advances in regularity of second-order hyperbolic mixed problems, and applications, K.R.T. Christopher *et al.*, Eds. Jones, editor. Springer-Verlag, Berlin, *Dynam. Report. Expositions Dynam. Systems (N.S.)* **3** (1994) 104–162.
- [11] R. Leis, *Initial boundary value problems in mathematical physics*. Teubner, Stuttgart (1972).
- [12] V.G. Maz'ya, *The Sobolev spaces*. Leningrad, Leningrad State University (1985), in Russian.
- [13] O. Nalin, Controlabilité exacte sur une partie du bord des équations de Maxwell. *C. R. Acad. Sci. Paris Sér. I Math.* **309** (1989) 811–815.
- [14] D.L. Russell, Boundary value control theory of the higher-dimensional wave equation. *SIAM J. Control Optim.* **9** (1971) 29–42.
- [15] G. Schwarz, *Hodge decomposition. A method for solving boundary value problems*. Springer Verlag, Berlin, *Lecture Notes in Math.* **1607** (1995).
- [16] D. Tataru, Unique continuation for solutions to PDE's; between Hoermander's theorem and Holmgren's theorem. *Comm. Partial Differential Equations* **20** (1995) 855–884.
- [17] N. Weck, Exact boundary controllability of a Maxwell problem. *SIAM J. Control Optim.* (to appear).