

## ON THE EXISTENCE OF VARIATIONS, POSSIBLY WITH POINTWISE GRADIENT CONSTRAINTS

SIMONE BERTONE<sup>1</sup> AND ARRIGO CELLINA<sup>1</sup>

**Abstract.** We propose a necessary and sufficient condition about the existence of variations, *i.e.*, of non trivial solutions  $\eta \in W_0^{1,\infty}(\Omega)$  to the differential inclusion  $\nabla\eta(x) \in -\nabla u(x) + \mathbf{D}$ .

**Mathematics Subject Classification.** 49K10, 35F30.

Received February 16, 2006.

### 1. A CONJECTURE

Purpose of the present paper is to derive conditions for the existence of (non trivial) solutions  $\eta \in W_0^{1,\infty}(\Omega)$  to the differential inclusion

$$\nabla\eta(x) \in -\nabla u(x) + \mathbf{D}$$

where  $\mathbf{D}$  is a given set and  $u$  is in  $W^{1,1}(\Omega)$  and satisfies

$$\nabla u(x) \in \text{co}(\mathbf{D});$$

(in the case  $\mathbf{D}$  is convex,  $\eta = 0$  is always a solution).

The problem of characterizing conditions for the existence of solutions is complex: in  $\mathbb{R}^2$ , consider the function  $v(x_1; x_2) = \sqrt{x_1^2 + x_2^2}$  whose gradient satisfies  $\|\nabla v(\cdot)\| = 1$ , let  $\mathbf{B}$  be the unit ball of  $\mathbb{R}^2$  and, on  $\Omega \subset \mathbb{R}^2$ , consider the inclusion

$$\nabla\eta \in -\nabla v + \mathbf{B}.$$

When  $\Omega$  is the open disk  $x_1^2 + x_2^2 < R^2$ , it is easy to see that non trivial solutions  $\eta$  do exist; however, when  $\Omega$  is the annulus  $r^2 < x_1^2 + x_2^2 < R^2$ , nontrivial solutions do *not* exist. Hence, the existence or non-existence of nontrivial solutions depends on the geometry of  $\Omega$ , and cannot be expressed by *local* conditions.

As a motivation for the problem, and for the name of *variations* proposed here for the solutions  $\eta$ , assume we are considering the problem of minimizing

$$\int_{\Omega} L(\nabla v(x)) \, dx$$

---

*Keywords and phrases.* Variations, differential inclusions, necessary conditions.

<sup>1</sup> Dipartimento di Matematica e Applicazioni, Università degli Studi di Milano-Bicocca, Via R. Cozzi 53, 20125 Milano, Italy; [simone.bertone@unimib.it](mailto:simone.bertone@unimib.it); [arrigo.cellina@unimib.it](mailto:arrigo.cellina@unimib.it)

© EDP Sciences, SMAI 2007

under given boundary conditions, where  $L$  is a convex function, for instance

$$L(\xi) = \begin{cases} 1 - \sqrt{1 - \|\xi\|^2} & \text{for } \|\xi\| \leq 1 \\ +\infty & \text{elsewhere;} \end{cases}$$

$L$  is finite for  $\xi$  in  $\mathbf{B}$ , the unit ball of  $\mathbb{R}^N$  equipped with the Euclidean norm. Let  $u$  be a solution to the minimum problem, and assume that we wish to derive the necessary conditions satisfied by  $u$ , hence to compare the values of the integral functional at  $u$  and at  $u + \eta$ . To find these conditions, we have to ask ourselves whether there are nontrivial variations  $\eta$ , such that  $\|\nabla u(x) + \nabla \eta(x)\| \leq 1$ , i.e., solutions to  $\nabla \eta(x) \in -\nabla u(x) + \mathbf{B}$ . In this case the function  $u$ , appearing in the differential inclusion we are investigating, is interpreted as the solution to a variational problem and the set  $\mathbf{D}$  as the effective domain of a convex Lagrangean.

We propose the following conjecture, on the existence of non trivial variations. In it, and in the remainder of the paper, by saying that a vector function  $p \in L^1_{loc}(\Omega)$  is such that  $\text{div}(p) = 0$  we mean that, for every  $\eta \in C^\infty_c(\Omega)$ , we have

$$\int_{\Omega} \langle p(x), \nabla \eta(x) \rangle dx = 0.$$

**Conjecture.** *Let  $\mathbf{D} \subset \mathbb{R}^N$ . Let  $u$  be a solution to*

$$\nabla u(x) \in \text{co}(\mathbf{D}).$$

*Then, the following a) and b) are in alternative:*

*a) there exists a nontrivial  $\eta \in W^{1,\infty}_0(\Omega)$ , solution to*

$$\nabla \eta(x) \in -\nabla u(x) + \mathbf{D} \tag{1}$$

*b) there exists a vector function  $p \in L^1_{loc}(\Omega)$ ,  $p(x) \neq 0$  a.e., such that  $\text{div}(p) = 0$ , and*

$$\langle p(x), \nabla u(x) \rangle = \sup_{k \in \mathbf{D}} \langle p(x), k \rangle \tag{2}$$

*for almost every  $x \in \Omega$ .*

**Examples.**

1) In the case  $\mathbf{D} = \mathbb{R}^N$ , condition b) is never satisfied and variations do always exist.

2) Consider again the function  $v(x_1; x_2) = \sqrt{x_1^2 + x_2^2} = \rho$ , whose gradient  $\nabla v(x_1; x_2) = \frac{1}{\rho}(x_1; x_2)$ . When  $\Omega$  is the annulus  $r^2 < x_1^2 + x_2^2 < R^2$ , non trivial solutions do not exist, hence a) is never satisfied. Let us show that b) is true: the vector function  $p(x_1; x_2) = \frac{1}{x_1^2 + x_2^2}(x_1; x_2)$  has pointwise divergence zero everywhere in  $\Omega$ ; moreover

$$\sup_{k \in B} \langle p(x_1; x_2), k \rangle = \frac{1}{\rho} = \langle p(x_1; x_2), \nabla v(x_1; x_2) \rangle.$$

Hence b) is satisfied.

When  $\Omega$  is the open disk  $x_1^2 + x_2^2 < R^2$ , non trivial  $\eta$  exist, so a) is satisfied. The vector  $p$  as used before has *not* weak divergence zero in  $\Omega$ , hence it does *not* prove that b) is satisfied. The fact that b) *cannot* be satisfied will be proved below.

In the present paper we prove the above conjecture under some additional regularity assumption.

2. THE CASE  $\nabla u = 0$

In this section we show that the Conjecture is verified in the case  $\nabla u = 0$ .

**Theorem 1.** *Let  $\mathbf{D} \subset \mathbb{R}^N$  and let  $u$  be such that  $\nabla u = 0 \in \text{co}(\mathbf{D})$ . Then, the following a) and b) are in alternative:*

- a) *there exists a nontrivial  $\eta \in W_0^{1,\infty}(\Omega)$ , solution to  $\nabla \eta(x) \in \mathbf{D}$ ;*
- b) *there exists a vector function  $p \in L_{loc}^1(\Omega)$ ,  $p(x) \neq 0$  a.e., such that  $\text{div}(p) = 0$ , and for a.e.  $x \in \Omega$ ,*

$$\sup_{k \in \mathbf{D}} \langle p(x), k \rangle = 0.$$

In the proof of Theorem 1 we will need the following lemma, whose proof is a consequence of a result appearing in [4].

**Lemma 1.** *Let  $\Omega \subset \mathbb{R}^N$  an open bounded set, and  $\mathbf{D} \subset \mathbb{R}^N$ . There exists a nontrivial function  $\eta \in W_0^{1,\infty}(\Omega)$  such that  $\nabla \eta(x) \in \mathbf{D}$  for a.e.  $x \in \Omega$ , if and only if  $0 \in \text{int}(\text{co}(\mathbf{D}))$ .*

*Proof.* When  $0 \in \text{int}(\text{co}(\mathbf{D}))$ , by Lemma 1, there exists  $\eta \in W_0^{1,\infty}(\Omega)$  such that, a.e.,  $\nabla \eta(x) \neq 0$ , hence  $\eta$  is non trivial and a) is always satisfied. We show that b) cannot be true: in fact, in this case, there must exist a ball  $B(0, r) \subset \text{co}(\mathbf{D})$  so that, for every non trivial vector function  $p$ , we have  $\langle p(x), \nabla u(x) \rangle \equiv 0$ , while  $\sup_{k \in \mathbf{D}} \langle p(x), k \rangle \geq r \|p(x)\|$ , that is positive on a set of positive measure.

When  $0 \notin \text{int}(\text{co}(\mathbf{D}))$ , again by Lemma 1, there is no  $\eta \in W_0^{1,\infty}(\Omega)$  apart from  $\eta = 0$ , so that a) is not satisfied. We show that b) is true: in fact, the convex sets  $0$  and  $\text{co}(\mathbf{D})$  can be weakly separated, i.e. there exists a non zero vector  $v$  such that  $\langle v, k \rangle \leq 0$  for every  $k \in \text{co}(\mathbf{D})$ , i.e., such that  $\sup_{k \in \text{co}(\mathbf{D})} \langle v, k \rangle \leq 0$ . This constant vector  $v$  is the required  $p$  : we have  $\sup_{k \in \text{co}(\mathbf{D})} \langle v, k \rangle \leq \langle v, 0 \rangle = 0$  while, since  $0 \in \text{co}(\mathbf{D})$ ,  $\sup_{k \in \text{co}(\mathbf{D})} \langle p, 0 \rangle \geq 0$ . This ends the proof.  $\square$

3. b) IMPLIES NOT a)

We prove that b) implies non a) under the additional assumption that  $p$  be locally Lipschitzian in  $\Omega$ , but no special assumptions on  $\mathbf{D}$ .

**Theorem 2.** *Let  $\Omega \subset \mathbb{R}^N$  be open,  $u \in W^{1,1}(\Omega)$  with  $\nabla u(x) \in \text{co}(\mathbf{D})$  for a.e.  $x \in \Omega$ . Assume that there exists a vector function  $p \in W_{loc}^{1,\infty}(\Omega)$ ,  $p(x) \neq 0$  for  $x \in \Omega$ , such that  $\text{div}(p) = 0$  and, for a.e.  $x \in \Omega$ ,*

$$\langle p(x), \nabla u(x) \rangle = \sup_{k \in \mathbf{D}} \langle p(x), k \rangle.$$

*Then the only solution  $\eta \in W_0^{1,\infty}(\Omega)$  to the differential inclusion*

$$\nabla \eta(x) \in -\nabla u(x) + \mathbf{D}$$

*is  $\eta \equiv 0$ .*

In the proof we will need the following lemma, a well known result (Liouville's Theorem) for the case of a differentiable  $p$ .

**Lemma 2.** *Let  $p$  as in Theorem 2. Let  $S(t; x)$  be the solution to the Cauchy problem  $\dot{x}(t) = p(x(t))$ ,  $x(0) = x$ . Then the map  $x \rightarrow S(t; x)$  is measure preserving.*

*Proof of Lemma 2.* Let  $\tilde{\Omega} \subset \Omega$  and  $\delta > 0$  be such that solutions issuing from  $\tilde{\Omega}$  are defined on the interval  $[0, \delta]$ . We wish to prove that for  $t \in [0, \delta]$  and  $x \in \tilde{\Omega}$ ,  $J(t; x)$ , the Jacobian of the transformation  $x \rightarrow S(t; x)$ , equals

1 *a.e.* By Rademacher’s Theorem, for *a.e.*  $x$ ,  $(Dp)$ , the matrix of (pointwise) partial derivatives of  $p$  exists. By a result of Tsuji [9], for *a.e.*  $x$ ,

$$J(t; x) = e^{\int_0^t \text{tr}((Dp)(\tau)) d\tau}$$

where the matrix  $(Dp)$  is computed along the solution  $S(\tau; x)$ . We wish to show that for *a.e.*  $x \in \Omega$ , for *a.e.*  $t \in [0, \delta]$ , we have  $\text{tr}((Dp)(\tau)) = 0$ . Let  $g$  be any of the components of the vector  $p$ ; fix  $\eta \in C_c^\infty(\Omega)$ . The sequence  $\frac{g(x+h_n e_i) - g(x)}{h_n}$  converges pointwise *a.e.* to  $\frac{\partial g(x)}{\partial x_i}$  and it is (locally) uniformly bounded, so that

$$\frac{1}{h_n} \int [g(x + h_n e_i) - g(x)] \eta(x) dx$$

converges both to  $\int \frac{\partial g(\xi)}{\partial \xi_i} \eta(\xi) d\xi$  and ([6], p. 132), to  $\int g_i(x) \eta(x) dx$ , with  $g_i$  the  $i$ th Sobolev partial derivative of  $g$ . So  $\int \left[ \frac{\partial g(x)}{\partial x_i} - g_i(x) \right] \eta(x) dx = 0$ , hence  $\frac{\partial g(x)}{\partial x_i} - g_i(x) = 0$ , for all components  $g$  and all  $i$ , with the exception of a set  $E \subset \Omega$  of  $N$  dimensional measure zero. In particular, on  $\Omega \setminus E$ , the pointwise divergence of  $S$  with respect to  $x$ ,  $\text{tr}(Dp)$  and the weak divergence  $\text{div}(p)$ , coincide and are zero.

For  $t$  in  $[0, \delta]$  and  $y$  in  $\{S(t; x) : x \in \tilde{\Omega}\}$  define the inverse map

$$S^{-1}(t; y) = (t; x).$$

$S^{-1}$  is locally Lipschitzian in its variables and sends the set  $[0, \delta] \times E$  into a set  $E^* \subset ([0, \delta] \times \tilde{\Omega})$  of  $N + 1$  dimensional measure zero. By Fubini’s Theorem, with the exception of a set  $X_{E^*}$  of  $N$  dimensional measure zero, the segments  $\{(t; x) : t \in [0, \delta]\}$  meet the set  $E^*$  on a set of 1 dimensional measure zero. This means that for  $x \notin X_{E^*}$ , for *a.e.*  $t \in [0, \delta]$ ,  $S(t; x) \notin ([0, \delta] \times E)$ , *i.e.*, that  $\text{tr}(Dp(x))$  and  $\text{div}(p)$ , computed along  $S(t; x)$ , coincide. □

*Proof of Theorem 2.*

a) We first notice that condition (2) implies that  $\nabla u(x) \in \text{co}(\mathbf{D})$  for *a.e.*  $x \in \Omega$ .

In fact, otherwise, we can find a set, of positive measure,  $\Omega_* \subset \Omega$  and  $\varepsilon > 0$  such that  $\nabla u(x) + \varepsilon p(x) \in \text{co}(\mathbf{D})$ . For  $x \in \Omega_*$ , we have

$$\sup_{k \in \text{co}(\mathbf{D})} \langle p(x), k \rangle \geq \langle p(x), \nabla u(x) + \varepsilon p(x) \rangle = \langle p(x), \nabla u(x) \rangle + \varepsilon \|p(x)\|^2 >$$

$$\langle p(x), \nabla u(x) \rangle = \sup_{k \in \mathbf{D}} \langle p(x), k \rangle.$$

Recalling that  $\sup_{k \in \text{co}(\mathbf{D})} \langle p(x), k \rangle = \sup_{k \in \mathbf{D}} \langle p(x), k \rangle$ , we obtain a contradiction.

b) To prove the theorem, suppose, by contradiction, that there exists a nontrivial  $\eta \in W_0^{1,\infty}(\Omega)$ , that verifies condition (1) almost everywhere.

In the case that  $\text{int}(\text{co}(\mathbf{D})) = \emptyset$ ,  $\mathbf{D}$  is contained in a hyperplane, and condition (1) implies that also  $\nabla \eta$  is in a hyperplane, a contradiction to Lemma 1. Hence, in what follows, we consider  $\text{int}(\text{co}(\mathbf{D})) \neq \emptyset$ .

c) *Claim.* For every  $x \in \Omega$ , there exists  $c$  such that  $\eta(S(t; x)) = c$  for  $t \in (\alpha_x, \beta_x)$ , the maximal interval of existence for the solution  $S$ .

*Proof of this claim.* By assumption, for almost every  $x \in \Omega$ ,

$$\langle \nabla u(x), p(x) \rangle = \sup_{k \in \mathbf{D}} \langle k, p(x) \rangle$$

and

$$\begin{aligned} \langle \nabla \eta(x), p(x) \rangle &= \langle -\nabla u(x), p(x) \rangle + \langle \nabla \eta(x) + \nabla u(x), p(x) \rangle \leq \\ &= -\langle \nabla u(x), p(x) \rangle + \sup_{k \in \mathbf{D}} \langle k, p(x) \rangle, \end{aligned}$$

so that

$$\langle \nabla \eta(x), p(x) \rangle \leq 0.$$

Since  $\eta \in W_0^{1,\infty}(\Omega)$ , the assumption on the divergence of  $p$  implies

$$\int_{\Omega} \langle \nabla \eta(x), p(x) \rangle \, dx = 0,$$

hence we obtain that, for almost every  $x \in \Omega$ ,

$$\langle \nabla \eta(x), p(x) \rangle = 0.$$

Fix  $x^* \in \Omega$ . Consider the  $N - 1$  dimensional affine space

$$V = x^* + p(x^*)^\perp.$$

There exists  $\delta > 0$  and  $r > 0$ , such that a solution  $S(t; v)$  to  $\dot{x} = p(x)$  and  $x(0) = v$  exists for  $v \in V \cap B(x^*, r)$  on an interval  $(-\delta, \delta)$ . The map  $(t; v) \rightarrow S(t; v)$  is Lipschitzian and invertible. Hence, by the coarea theorem, with the exception of a subset of  $V$  of  $N - 1$  dimensional measure zero,  $S(t; v)$  meets the set  $M$ , where  $\langle \nabla \eta(x), p(x) \rangle \neq 0$ , on a subset of  $(-\delta, \delta)$  of 1-dimensional measure zero, and, outside of this exceptional set, we have

$$\frac{d}{dt} \eta(S(t; v)) = \langle \nabla \eta(S(t; v)), p(S(t; v)) \rangle = 0.$$

Hence, there exists a sequence  $v_n \rightarrow x^*$  such that  $\eta(S(t; v_n)) \equiv c_n$  on  $(-\delta, \delta)$ . Since the limit of solutions is a solution and  $\eta$  is continuous, we have that  $\eta(S(t; x^*)) \equiv c$  on  $(-\delta, \delta)$ . This local reasoning can be extended to the maximal interval of existence, proving the claim.

d) Let  $\bar{x}$  be such that  $\eta(\bar{x}) > 0$ , and define

$$E = \left\{ x \in \Omega : \eta(x) \geq \frac{1}{2} \eta(\bar{x}) \right\} \subset \Omega.$$

The set  $E$  is nonempty, compact,  $\text{int}(E) \neq \emptyset$  and  $d(E, \partial\Omega) > 0$ . As a consequence of c), it cannot happen that there exists  $x \in E$  such that, for some  $t \in (\alpha_x, \beta_x)$ ,  $S(t; x) \notin E$ . Hence for every  $x \in E$  and every  $t \in (\alpha_x, \beta_x)$ ,  $S(t; x) \in E$ . By the basic theorems on the prolongability of solutions to ordinary differential equations, it follows then that the solution  $S(t; x)$  must be defined for every  $t \in \mathbb{R}$ , since  $d(E, \partial\Omega) > 0$ . Hence, for every  $t \in \mathbb{R}$ , the map  $S(t; \cdot)$  is a bijection of  $E$  into itself and, in addition, by Lemma 1, it is measure preserving.

e) We wish to apply the following Poincaré recurrence Theorem to the map  $S(t; \cdot)$ , (see for instance [1] for the proof).

**Lemma 3** (Poincaré). *Let  $E$  be a compact, nonempty set such that  $\text{int}(E) \neq \emptyset$ , and let  $\psi : E \rightarrow E$  a bijective, measure preserving function. Then, for every  $x_0 \in \text{int}(E)$  and every  $\varepsilon > 0$ , there exists an integer  $k > 0$  such that*

$$\psi^k(B(x_0, \varepsilon)) \cap B(x_0, \varepsilon) \neq \emptyset.$$

Going back to the proof, let  $r^0 > 0$  and  $x^0$  be such that  $B(x^0, r^0) \subset E$  and let  $t_0 > 0$  be such that  $S(t_0; x^0) \neq x^0$ . Let  $V \subset \Omega$  be a neighborhood of the trajectory

$$\{S(t; x^0) : t \in [0, t_0]\}$$

and let  $p^0 > 0$  be such that  $\|p(x)\| \geq p^0$  for  $x$  in  $V$ . Let  $r \leq r^0$  be so small that:

$$S(t_0; B(x^0, r)) \cap B(x^0, r) = \emptyset$$

and, for every  $\xi \in B(x^0, r)$ , the solution  $S(t; \xi) \in V$  for  $t \in [0, t_0]$ . Applying Poincaré’s method we obtain that, for every  $\rho < r$ , there exist  $\xi_\rho \in B(x^0, \rho)$  and an integer  $\nu_\rho > 1$ , such that

$$|S(t_0\nu_\rho; \xi_\rho) - x^0| \leq \rho.$$

f) Choose  $v \in \text{int}(\mathbf{D})$  and let  $s > 0$  be such that  $B(v, s) \subset \mathbf{D}$ . Consider the function  $u_0$  defined by

$$u_0(x) = u(x) - \langle v, x \rangle.$$

Condition (2) implies that  $u_0$ , computed along  $S(t; x)$ , for  $x \in B(x^0, r)$ , is strictly increasing:

$$\begin{aligned} \frac{d}{dt}u_0(S(t; x)) &= \langle \nabla u(S(t; x)) - v, p(S(t; x)) \rangle = \sup_{k \in D} \langle k - v, p(S(t; x)) \rangle \\ &\geq s \|p(S(t; x))\| > 0; \end{aligned}$$

in particular, for  $\xi \in B(x^0, \rho)$ , with  $\rho \leq r$ , we obtain

$$u_0(S(t_0\nu_\rho; \xi)) - u_0(\xi) \geq t_0\nu_\rho s p^0 \geq t_0 s p^0.$$

This last estimate is independent of  $\rho$ .

Apply this estimate to  $\xi_\rho$ ; we have that both  $\xi_\rho$  and  $S(t_0\nu_\rho; \xi_\rho)$  are in  $B(x^0, \rho)$ . By the continuity of  $u_0$  at  $x^0$ , the difference  $u_0(S(t_0\nu_\rho; \xi_\rho)) - u_0(\xi_\rho)$  can be made arbitrarily small by decreasing  $\rho$ , a contradiction.  $\square$

The following result completes the discussion of the example in Section 2.

**Theorem 3.** *Let  $\Omega \subset \mathbb{R}^2$  be the open disk  $x_1^2 + x_2^2 < 1$  and  $v(x_1; x_2) = \sqrt{x_1^2 + x_2^2}$ . There is no vector function  $p \in L^1_{loc}(\Omega)$ ,  $p(x) \neq 0$  a.e., such that  $\text{div}(p) = 0$ , and*

$$\langle p(x), \nabla v(x) \rangle = \sup_{k \in \mathbf{B}} \langle p(x), k \rangle$$

for almost every  $x \in \Omega$ .

*Proof of Theorem 3.* The function  $\eta = -\sqrt{x_1^2 + x_2^2} + 1$  is in  $W^{1,\infty}(\Omega)$  and is a solution to the differential inclusion

$$\nabla \eta(x) \in -\nabla v(x) + \mathbf{B}.$$

Assume that  $p$  exists. By assumption we must have, for almost every  $x \in \Omega$ ,

$$\langle p(x), \nabla v(x) \rangle = \|p(x)\|$$

so that  $p(x) = \alpha(x) \frac{x}{\|x\|}$ , and  $\alpha \geq 0$ . On the other hand, in c) of the proof of the previous theorem we have obtained that, for almost every  $x \in \Omega$ ,

$$\langle \nabla \eta(x), p(x) \rangle = 0,$$

so that  $\alpha(x) = 0$  a.e. in  $\Omega$ .  $\square$

4. WHEN  $\mathbf{D} = \mathbf{B}$ , NOT a) IMPLIES b)

We prove this part of the conjecture in the case  $\mathbf{D} = \mathbf{B}$ .

**Theorem 4.** *Let  $u \in W^{1,\infty}(\Omega)$  be a solution to  $\nabla u(x) \in \mathbf{B}$  and assume that there exist no nontrivial  $\eta \in W_0^{1,\infty}(\Omega)$ , solution to the differential inclusion*

$$\nabla \eta(x) \in -\nabla u(x) + \mathbf{B}.$$

Then:

- i) the solution  $u$  belongs to  $C^1(\Omega) \cap W_{loc}^{2,\infty}(\Omega)$ ;
- ii) there exists  $p \in L_{loc}^1(\Omega)$ ,  $p(x) \neq 0$  for almost every  $x \in \Omega$ , such that  $\operatorname{div}(p) = 0$ , and

$$\langle p(x), \nabla u(x) \rangle = \sup_{k \in \mathbf{B}} \langle p(x), k \rangle$$

for almost every  $x \in \Omega$ .

**Remark 1.** In the proof of Theorem 4, we will construct a function  $p$  that verifies ii). This function  $p$  can be interpreted as a mass-transfer vector field, and from condition (2) we see that  $\nabla u$  determines the optimal direction for  $p$ . Hence, we expect  $p$  to be of the form  $p = \lambda \nabla u$ , for a suitable function  $\lambda(x)$ , and we compute  $\lambda$  by the equation  $\operatorname{div}(\lambda \nabla u) = 0$ . As appears in [7], this equation is related to the Monge-Kantorovich transport problem. In particular,  $\lambda$  plays the role of a transport density, and is the Lagrange multiplier for the constraint  $\nabla u \in \mathbf{B}$ .

*Proof of Theorem 4.* The proof makes use of some results and techniques developed in [5].

a) Fix any point  $x^0 \in \Omega$ . Using Lemmas 2.2 and 2.3 of [5], from the fact that there is no variation  $\eta$  such that  $u(x^0) + \eta(x^0) < u(x^0)$ , we infer the existence of at least one unit vector, a direction,  $d^+$ , with the property that, for every  $r$  such that the ball  $B(x^0, r)$  is contained in  $\Omega$ , we have  $u(x^0 + rd^+) - u(x^0) = r$ . Such a direction will be called a direction of maximal growth. By the same reasons, since there is no variation  $\eta$  such that  $u(x^0) + \eta(x^0) > u(x^0)$ , we infer the existence of at least one direction,  $d^-$ , such that  $u(x^0) - u(x^0 + rd^-) = r$ . However we must have that  $d^+ = d^-$ , in fact, since  $u$  is Lipschitzian of constant 1, we have

$$r \|d^+ + d^-\| \geq |u(x^0 + rd^+) - u(x^0 - rd^-)| = |u(x^0 + rd^+) - u(x^0) - u(x^0 + rd^-) + u(x^0)| = 2r,$$

*i.e.*  $\|d^+ + d^-\| = \|d^+\| + \|d^-\|$ , that implies  $d^+ = d^-$ . Notice that this result implies that  $d^+$  and  $d^-$  are unique. Hence, from the assumption that there is no variation  $\eta$ , to each  $x \in \Omega$  we associate a unique direction  $d(x)$  such that  $u(x + rd(x)) - u(x) = r$  as long as  $x + rd(x) \in \Omega$ ; *i.e.*, there exists a unique segment  $(x + \alpha(x)d(x), x + \beta(x)d(x))$ ,  $\alpha(x) < 0 < \beta(x)$ , such that:  $x + \alpha(x)d(x) \in \partial\Omega$ ,  $x + \beta(x)d(x) \in \partial\Omega$  and  $u(x + \lambda_1 d(x)) - u(x + \lambda_2 d(x)) = \lambda_1 - \lambda_2$  for every  $\lambda_1, \lambda_2 \in (\alpha(x), \beta(x))$ . The direction  $d$  has the following interpretation: at every point  $x^0$  such that  $\nabla u(x^0)$  exists, we have that  $\nabla u(x^0) = d(x^0)$ . In fact, from

$$u(x) - u(x^0) = \langle \nabla u(x^0), x - x^0 \rangle + \|x - x^0\| o(\|x - x^0\|),$$

we obtain

$$r = r \langle \nabla u(x^0), d(x^0) \rangle + ro(r),$$

that implies  $\nabla u(x^0) = d(x^0)$ . Moreover, the following property holds: for no  $y \in \Omega$  we can have

$$y = x + \lambda d(x) = x' + \lambda' d(x')$$

unless  $d(x') = d(x)$ . In fact, otherwise, both  $d(x)$  and  $d(x')$  would be directions of maximal growth at  $y$ , contradicting the uniqueness of  $d(y)$ .

b) *Claim.* Let  $\rho$  be such that  $B(x^0, \rho) \subset \Omega$ . Then, on  $B(x^0, \frac{\rho}{6})$ , the map  $x \rightarrow d(x)$  is Lipschitzian of constant  $\Lambda = \frac{3}{\rho}$ .

*Proof of this Claim.* Let  $P$  and  $P'$  in  $B(x^0, \frac{\rho}{6})$ , so that  $\|P - P'\| \leq \frac{\rho}{3}$ . Set  $d = d(P)$  and  $d' = d(P')$ ; let  $O$  on the line  $r = \{P + \lambda d\}$  and  $O'$  on the line  $r' = \{P' + \lambda d'\}$  be such that  $\|O - O'\| = \inf_{Q \in r, Q' \in r'} \|Q - Q'\|$ . We have that  $(O - O')$  is orthogonal both to  $r$  and to  $r'$ . Two cases are possible: either, a),  $\inf \{\|P - O\|, \|P' - O'\|\} > \frac{\rho}{3}$  or, b),  $\inf \{\|P - O\|, \|P' - O'\|\} \leq \frac{\rho}{3}$ .

Consider case a). Call  $P$  the point such that  $\|P - O\| \leq \|P' - O'\|$ . We will need the line  $r'' = r' + (O - O')$ : it is the parallel to  $r'$  in the plane containing  $r$  and orthogonal to  $(O - O')$ . Let  $P''$  be the projection of  $P'$  on  $r''$ . Since  $\|P'' - O\| = \|P' - O'\| \geq \|P - O\|$ , on the segment  $[O, P'']$  choose  $P_i$  such that  $\|P_i - O\| = \|P - O\|$  and consider the isosceles triangle  $O, P', P_i$ : we have

$$\frac{\|d - d'\|}{1} = \frac{\|P_i - P\|}{\|P - O\|},$$

so that  $\|P - O\| \geq \frac{\rho}{3}$  implies

$$\|d - d'\| \leq \frac{3}{\rho} \|P_i - P\|.$$

We claim that  $\|P'' - P\| \geq \|P_i - P\|$ . In fact, the angle  $P, P_i, P''$  is larger than  $\frac{\pi}{2}$ , being the triangle  $O, P', P_i$  isosceles, so that

$$\begin{aligned} \|P'' - P\|^2 &= \|P - P_i\|^2 + \|P_i - P''\|^2 + 2 \langle P - P_i, P_i - P'' \rangle \\ &\geq \|P - P_i\|^2 + \|P_i - P''\|^2 \geq \|P - P_i\|^2. \end{aligned}$$

We have shown that

$$\|d - d'\| \leq \frac{3}{\rho} \|P_i - P\| \leq \frac{3}{\rho} \|P'' - P\| \leq \frac{3}{\rho} \|P' - P\|.$$

Consider case b). Consider the two points  $O$  and  $O'$ ; since  $\|O - O'\| \leq \|P - P'\|$ , we obtain that both points  $O$  and  $O'$  are in  $B(x^0, \rho)$ , so that  $u$  is defined at  $O$  and  $O'$ . For case b), we assign names to the points  $P$  and  $P'$  by assuming that  $u(O') \geq u(O)$ . With this choice of names, consider again the lines  $r, r'$  and set again  $r'' = r' + (O - O')$ . On  $r$  consider the segment  $[A, D]$ , centered at  $O$ , such that  $\|A - O\| = \|D - O\| = \frac{\rho}{3}$ ; on  $r'$ , the segment  $[B', C']$ , centered at  $O'$ , such that  $\|B' - O'\| = \|C' - O'\| = \frac{\rho}{3}$ ; orientations are chosen so that  $A = O + \frac{\rho}{3}d$  and  $B' = O' + \frac{\rho}{3}d'$ . Call  $B$  and  $C$  the projections of  $B'$  and  $C'$  on the line  $r''$ . We obtain

$$\begin{aligned} \|B' - D\| &\geq u(B') - u(D) = u(B') - u(O') + u(O') - u(O) + u(O) - u(D) \\ &\geq u(B') - u(O') + u(O) - u(D) = \frac{\rho}{3} + \frac{\rho}{3}. \end{aligned}$$

Set  $H = \frac{1}{2}A + \frac{1}{2}B$ . We have:

$$\begin{aligned} \|H - O\|^2 &= \left\| \frac{1}{2}(A - O) + \frac{1}{2}(B - O) \right\|^2 = \left\| \frac{1}{2}(O - D) + \frac{1}{2}(B - O) \right\|^2 \\ &= \left\| \frac{1}{2}(B - D) \right\|^2 = \frac{1}{4} (\|B' - D\|^2 - \|O - O'\|^2), \end{aligned}$$

the last equality deriving from the Pythagorean Theorem applied to the triangle  $D, B, B'$ . Hence we have:

$$\begin{aligned} \frac{1}{4} \|A - B\|^2 &= \|B - H\|^2 = \|B - O\|^2 - \|H - O\|^2 \\ &= \left(\frac{\rho}{3}\right)^2 - \frac{1}{4} (\|B' - D\|^2 - \|O - O'\|^2) \leq \left(\frac{\rho}{3}\right)^2 - \frac{1}{4} \left(\frac{2\rho}{3}\right)^2 + \frac{1}{4} \|O - O'\|^2 \\ &= \frac{1}{4} \|O - O'\|^2. \end{aligned}$$

We obtain

$$\|A - B\| = \left\| O + \frac{\rho}{3}d - \left( O + \frac{\rho}{3}d' \right) \right\| \leq \|O - O'\| \leq \|P - P'\|.$$

We conclude that, for case b) as well, we have

$$\|d - d'\| \leq \frac{3}{\rho} \|P - P'\|$$

proving the claim.

c) We claim that, as a consequence of the Lipschitzianity of  $d$ , we have that  $u \in C^1(\Omega) \cap W_{loc}^{2,\infty}(\Omega)$ . The directions of the coordinate axis are denoted by  $e_i$ .

Fix  $x$ ; let  $B(x, r) \subset \Omega$  and let  $\Lambda$  be a Lipschitz constant for  $d$  in  $B(x, r)$ . We first notice that if it happens that on the intersection of the line  $\{x + te_i : t \in \mathbb{R}\}$  with  $B(x, r)$ ,  $u$  is differentiable at  $x + te_i$  for almost every  $t$ , then we must have

$$|u(x + he_i) - u(x) - h \langle d(x), e_i \rangle| \leq h^2 \Lambda.$$

In fact, the Lipschitzian map  $t \rightarrow u(x + te_i)$  is the integral of its derivative, that coincides, for *a.e.*  $t$ , with  $\langle d(x + te_i), e_i \rangle$ , so that

$$|u(x + he_i) - u(x) - h \langle d(x), e_i \rangle| = \left| h \int_0^1 \langle d(x + she_i) - d(x), e_i \rangle ds \right| \leq h^2 \Lambda.$$

Notice next that, since  $\nabla u(x)$  exists for *a.e.*  $x \in \Omega$ , there must exist a sequence  $x_n \rightarrow x$  such that, on the intersection of the line  $\{x_n + te_i : t \in \mathbb{R}\}$  with  $B(x, r)$ ,  $\nabla u(x_n + te_i)$  exists for *a.e.*  $t$ . Then we have:

$$\begin{aligned} |u(x + he_i) - u(x) - h \langle d(x), e_i \rangle| &= |u(x_n + he_i) - u(x_n) - h \langle d(x_n), e_i \rangle + h \langle d(x_n) - d(x), e_i \rangle \\ &\quad + u(x + he_i) - u(x_n + he_i) + u(x_n) - u(x)| \\ &\leq h^2 \Lambda + h \Lambda |x_n - x| + 2|x_n - x|. \end{aligned}$$

Letting  $n \rightarrow \infty$  we obtain that  $\frac{\partial u}{\partial x_i}$  exists at  $x$  and equals  $\langle d(x), e_i \rangle$ . Since the gradient is continuous, we obtain that  $u$  is differentiable and that  $u \in C^1(\Omega)$ .

Fix  $\eta \in C_c^\infty(\Omega)$ . Then on  $\text{supp}(\eta)$ ,  $\nabla u(x) = d(x)$  is uniformly Lipschitzian: hence, see [8], for each component  $d^i$  and each  $j$  there is  $g_j^i$  such that

$$\int_{\Omega} g_j^i \eta \, dx = - \int_{\Omega} d^i \frac{\partial \eta}{\partial x_j} \, dx.$$

This proves i).

d) As established in the Remark, the map  $p$ , as required in ii), will be of the form  $\lambda(x)d(x)$ . To find  $\lambda$  amounts to finding a weak solution to the equation  $\text{div}(\lambda(x)d(x)) = 0$ , where  $\text{div}(d(x)) \in L_{loc}^\infty(\Omega)$ .

Fix  $x^* \in \Omega$  and consider the corresponding level set for the function  $u$ , *i.e.*  $\{x : u(x) = u(x^*)\}$ . We claim that we can parametrize locally this set by a differentiable and invertible map  $\phi_{x^*}$  from an open set  $V_{x^*}$  in

a  $N - 1$  space, to  $\Omega$ , *i.e.* that there exists  $V_{x^*}$ ,  $\phi_{x^*}$ ,  $r^*$  such that  $u(\phi_{x^*}(\xi)) \equiv u(x^*)$ , for every  $\xi \in V_{x^*}$  and  $\phi(V_{x^*}) = \{u(x) = u(x^*)\} \cap B(x^*, r^*)$ .

*Proof of this Claim.* Consider the  $N - 1$  dimensional space  $d(x^*)^\perp$ , defined by the equation  $\langle d(x^*), x \rangle = \langle d(x^*), x^* \rangle$ ; let  $d_i(x^*) \neq 0$ , set the  $N - 1$  vector  $\xi$  be  $\xi_j = x_j$ ,  $j \neq i$ , and set  $\xi^*$  be  $\xi_j^* = x_j^*$ ,  $j \neq i$ , so that  $d(x^*)^\perp$  is the image of the affine map  $\ell$ , given by  $\ell(\xi)_j = x_j$ ,  $j \neq i$ , and

$$\ell(\xi)_i = \frac{\langle d(x^*), x^* \rangle - \sum_{j \neq i} d_j(x^*) \xi_j}{d_i(x^*)}.$$

The map  $\ell$  is one to one from  $\mathbb{R}^{N-1}$  to  $\mathbb{R}^N$ . For  $\xi$  in a sufficiently small neighborhood  $V_{x^*}$  of  $\xi^*$ , so that the maps are defined, we have that  $u(\ell(\xi) + td(\ell(\xi))) = u(\ell(\xi)) + t$  and  $u(\ell(\xi) + td(\ell(\xi)))$  assumes the value  $u(x^*)$  for  $u(x^*) - u(\ell(\xi))$ . The required parametrization is given by the (differentiable) map  $\phi_{x^*}(\xi) = \ell(\xi) + (u(x^*) - u(\ell(\xi)))d(\ell(\xi))$ . The map  $\phi_{x^*}$  is invertible: assume that  $\ell(\xi) + t(\xi)d(\ell(\xi)) = \ell(\xi') + t(\xi')d(\ell(\xi')) = P$ ; then  $u(P) - u(x^*) = t(\xi)$ ,  $u(P) - u(x^*) = t(\xi')$  and, by the results of a),  $d(\ell(\xi)) = d(\ell(\xi'))$ , so that  $\ell(\xi) = \ell(\xi')$  and  $\xi = \xi'$ .

e) Consider the flow  $S(t; x) = x + td(x)$ : it is a solution to the Cauchy problem

$$\frac{d}{dt}S(t; x) = d(S(t; x)), \quad S(0; x) = x.$$

In particular, consider the map  $(t; \xi) \rightarrow S(t; \phi_{x^*}(\xi))$ : by the basic theorems on uniqueness for ordinary differential equations, and by the invertibility of  $\phi_{x^*}$ , it is an invertible map.

We will denote by  $D$  the square matrix of partial derivatives of the vector field  $d(x)$  and by  $M_x(t)$  the square matrix of partial derivatives of  $S(t; x)$  with respect to the space variables, computed at  $x$ , *i.e.*  $M_x(t) = I + tD(x)$ . Since the vector field  $d$  is autonomous, we have the basic identity

$$M_x(t)d(x) = d(S(t; x)).$$

In addition, Lindelöf's Theorem on differentiability with respect to initial conditions implies that

$$\det(M_x(t)) = e^{\int_0^t \text{tr}D(s) ds}$$

where the trace of  $D$  appearing at the right hand side is computed along  $S(s; x)$ . As a consequence of the uniform Lipschitzianity of  $d$  on compact subsets of  $\Omega$ , we have that on a compact set, there exists  $k$  such that  $\det(M_x(t)) \geq k > 0$ . Denote by  $\Phi_\xi$  the  $N \times (N - 1)$  matrix of partial derivatives of  $\phi$  with respect to  $\xi$ . We obtain that

$$D_{(t; \xi)}(S(t; \phi(\xi))) = (d(S(t; \phi(\xi))); M_{\phi(\xi)}(t)\Phi_\xi)$$

and, recalling that  $d(S(t; \phi(\xi))) = d(\phi(\xi)) = M_{\phi(\xi)}(t)d(\phi(\xi))$ , we obtain

$$\det(D_{(t; \xi)}(S(t; \phi(\xi)))) = \det(M_{\phi(\xi)}(t)) \det(d(\phi(\xi)); \phi_{\xi_1}; \dots; \phi_{\xi_{N-1}}).$$

f) An easy contradiction argument shows that the set

$$O_{x^*} = \{(t; \xi) : \alpha(\phi_{x^*}(\xi)) < t < \beta(\phi_{x^*}(\xi)); \xi \in V_{x^*}\}$$

is an open subset of  $\mathbb{R} \times \mathbb{R}^{N-1}$  and, being the continuous map  $S(t; \phi_{x^*}(\xi))$  one to one, its image  $S_{x^*}$  is an open subset of  $\Omega$ .

Consider a countable covering of  $\Omega$  by sets  $S_{x_n}$ ,  $n = 1, \dots$  (for brevity we will set  $S_{x_n} = S_n$ ,  $V_{x_n} = V_n$  and  $\phi_{x_n} = \phi_n$ ). Fix  $x \in S_n$ ; let  $t$  and  $\xi$  be such that  $x = S(t; \phi_n(\xi))$  and set

$$\lambda_n(x) = \frac{1}{\det M_{\phi_n(\xi)}(t)}.$$

This definition sets (arbitrarily)  $\lambda_n$  to be 1 on the level set  $\{x : u(x) = u(x_n)\} \cap S_n$ . Set  $E_1 = \Omega \cap S_1$ ;  $E_{n+1} = \Omega \cap [S_{n+1} \setminus E_n]$ , so that  $\Omega = \bigcup E_n$ , and the  $E_n$  are disjoint.

In general, define  $\lambda(x) = \sum \lambda_n(x) \chi_{E_n}$ . On a compact subset of  $\Omega$ , we have that  $\lambda_n(x) \leq h$  where  $h$  does not depend on  $n$ , so that  $\lambda \in L^\infty_{loc}(\Omega)$ . We claim that, for every  $\eta \in C^\infty_c(\Omega)$ , we have

$$\int_\Omega \lambda(x) \langle d(x), \nabla \eta(x) \rangle \, dx = \sum_n \int_{E_n} \lambda_n(x) \langle d(x), \nabla \eta(x) \rangle \, dx = 0$$

i.e. that the map  $p(x) = \lambda(x)d(x)$  has divergence zero.

On  $E_n$  consider the change of variables given by  $x = S(t; \phi_n(\xi))$ , with Jacobian  $J_n(t; \xi) = |\det D_{(t;\xi)}(S(t; \phi(\xi)))|$ . We have

$$\begin{aligned} \lambda_n(S(t; \phi_n(\xi))) J_n(t; \xi) &= \frac{1}{\det M_{\phi_n(\xi)}(t)} |\det M_{\phi(\xi)}(t) \det (d(\phi(\xi)); \phi_{\xi_1}; \dots; \phi_{\xi_{n-1}})| \\ &= |\det (d(\phi(\xi)); \phi_{\xi_1}; \dots; \phi_{\xi_{n-1}})|, \end{aligned}$$

so that

$$\begin{aligned} \int_{E_n} \lambda_n(x) \langle d(x), \nabla \eta(x) \rangle \, dx &= \int_{E_n} \lambda_n(S(t; \phi_n(\xi))) \langle d(S(t; \phi_n(\xi))), \nabla \eta(S(t; \phi_n(\xi))) \rangle J_n(t; \xi) \, d(t; \xi) \\ &= \int \left( \int_{\alpha(\phi_n(\xi))}^{\beta(\phi_n(\xi))} \lambda_n(S(t; \phi_n(\xi))) \langle d(S(t; \phi_n(\xi))), \nabla \eta(S(t; \phi_n(\xi))) \rangle J_n(t; \xi) \, dt \right) \, d\xi \\ &= \int \left( \int_{\alpha(\phi_n(\xi))}^{\beta(\phi_n(\xi))} \frac{d}{dt} \eta(S(t; \phi_n(\xi))) \, dt \right) |\det (d(\phi(\xi)); \phi_{\xi_1}; \dots; \phi_{\xi_{n-1}})| \, d\xi. \end{aligned}$$

Since, for every  $\xi$ ,  $S(\alpha(\phi_n(\xi)); \phi_n(\xi))$  and  $S(\beta(\phi_n(\xi)); \phi(\xi))$  belong to  $\partial\Omega$ , we obtain that  $\eta(S(\alpha(\phi(\xi)); \phi(\xi))) = \eta(S(\beta(\phi(\xi)); \phi(\xi))) = 0$  for every  $\xi$ , so that

$$\int_\Omega \lambda(x) \langle d(x), \nabla \eta(x) \rangle \, dx = 0.$$

g) We have

$$\langle p(x), \nabla u(x) \rangle = \langle \lambda(x)d(x), \nabla u(x) \rangle = \lambda(x) = \sup_{k \in B} \langle p(x), k \rangle,$$

concluding the proof. □

**Remark 2.** The vector  $p(\cdot)$  admits a divergence in the integral sense, but need not belong to  $W^{1,1}_{loc}(\Omega)$ .

In fact, in  $\mathbb{R}^2$  consider

$$\Omega = \{(x; y) : x^2 + y^2 < 1, x \leq 0, y > 0\} \cup \{(x; y) : x^2 + (y - 1)^2 < 1, x \geq 0, y < 1\}.$$

On  $\Omega$  set  $P = (x; y)$  and

$$u(P) = \begin{cases} \sqrt{x^2 + y^2} & \text{if } x \leq 0 \\ 1 - \sqrt{x^2 + (y - 1)^2} & \text{otherwise.} \end{cases}$$

Then

$$\nabla u(P) = \begin{cases} \frac{P}{\|P\|} & \text{if } x \leq 0 \\ \frac{(0;1)-P}{\|P-(0;1)\|} & \text{otherwise} \end{cases}$$

and

$$\Delta u(P) = \begin{cases} \frac{1}{\|P\|} & \text{if } x \leq 0 \\ \frac{-1}{\|P-(0;1)\|} & \text{otherwise.} \end{cases}$$

One verifies that the differential equation for  $\lambda$

$$\langle \nabla \lambda(P), \nabla u(P) \rangle + \lambda(P) \Delta u(P) = 0$$

admits the solution

$$\lambda(P) = \begin{cases} \frac{1}{\|P\|} & \text{if } x \leq 0 \\ \frac{1}{\|P-(0;1)\|} & \text{otherwise.} \end{cases}$$

Hence

$$p(P) = \begin{cases} \frac{P}{\|P\|^2} & \text{if } x \leq 0 \\ \frac{(0;1)-P}{\|P-(0;1)\|^2} & \text{otherwise,} \end{cases}$$

that has a jump discontinuity through the line  $x = 0$ . Hence  $p$  cannot belong to  $W^{1,1}(\Omega)$ .

## REFERENCES

- [1] V.I. Arnold, *Mathematical methods of classical mechanics*, *Graduate Texts in Mathematics* **60**, Springer-Verlag, New York, Heidelberg, Berlin.
- [2] H. Brezis, *Analyse fonctionnelle, théorie et applications*. Masson, Paris (1983).
- [3] A. Cellina, On minima of a functional of the gradient: necessary conditions. *Nonlinear Anal.* **20** (1993) 337–341.
- [4] A. Cellina, On minima of a functional of the gradient: sufficient conditions. *Nonlinear Anal.* **20** (1993) 343–347.
- [5] A. Cellina and S. Perrotta, On the validity of the maximum principle and of the Euler-Lagrange equation for a minimum problem depending on the gradient. *SIAM J. Control Optim.* **36** (1998) 1987–1998.
- [6] L.C. Evans, *Partial Differential Equations*, *Graduate Studies in Mathematics* **19**, American Mathematical Society, Providence, Rhode Island (1998).
- [7] L.C. Evans and W. Gangbo, Differential equations methods for the Monge-Kantorovich mass transfer problem, *Mem. Amer. Math. Soc.* **137** (1999) 653.
- [8] L.C. Evans and R.F. Gariepy, *Measure Theory and Fine Properties of Functions*, *Studies in Advanced Mathematics*, CRC Press, Boca Raton, FL (1992).
- [9] M. Tsuji, On Lindelöf's theorem in the theory of differential equations. *Jap. J. Math.* **16** (1939) 149–161.