

HAMILTON-JACOBI EQUATIONS FOR OPTIMAL CONTROL ON NETWORKS WITH ENTRY OR EXIT COSTS

MANH KHANG DAO*

Abstract. We consider an optimal control on networks in the spirit of the works of Achdou *et al.* [*NoDEA Nonlinear Differ. Equ. Appl.* **20** (2013) 413–445] and Imbert *et al.* [*ESAIM: COCV* **19** (2013) 129–166]. The main new feature is that there are entry (or exit) costs at the edges of the network leading to a possible discontinuous value function. We characterize the value function as the unique viscosity solution of a new Hamilton-Jacobi system. The uniqueness is a consequence of a comparison principle for which we give two different proofs, one with arguments from the theory of optimal control inspired by Achdou *et al.* [*ESAIM: COCV* **21** (2015) 876–899] and one based on partial differential equations techniques inspired by a recent work of Lions and Souganidis [*Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **27** (2016) 535–545].

Mathematics Subject Classification. 34H05, 35F21, 49L25, 49J15, 49L20, 93C30

Received June 27, 2017. Accepted January 8, 2018.

1. INTRODUCTION

A network (or a graph) is a set of items, referred to as vertices or nodes, which are connected by edges (see Fig. 1 for example). Recently, several research projects have been devoted to dynamical systems and differential equations on networks, in general or more particularly in connection with problems of data transmission or traffic management (see for example Garavello and Piccoli [14] and Engel *et al.* [12]).

An optimal control problem is an optimization problem where an agent tries to minimize a cost which depends on the solution of a controlled ordinary differential equation (ODE). The ODE is controlled in the sense that it depends on a function called the control. The goal is to find the best control in order to minimize the given cost. In many situations, the optimal value of the problem as a function of the initial state (and possibly of the initial time when the horizon of the problem is finite) is a viscosity solution of a Hamilton-Jacobi-Bellman partial differential equation (HJB equation). Under appropriate conditions, the HJB equation has a unique viscosity solution characterizing by this way the value function. Moreover, the optimal control may be recovered from the solution of the HJB equation, at least if the latter is smooth enough.

The first articles about optimal control problems in which the set of admissible states is a network (therefore the state variable is a continuous one) appeared in 2012: in [2], Achdou *et al.* derived the HJB equation associated to an infinite horizon optimal control on a network and proposed a suitable notion of viscosity solution. Obviously, the main difficulties arise at the vertices where the network does not have a regular differential

Keywords and phrases: Optimal control, networks, Hamilton-Jacobi equation, viscosity solutions, uniqueness, switching cost.
IRMAR, Université de Rennes 1, 35000 Rennes, France.

* Corresponding author: manh-khang.dao@univ-rennes1.fr

structure. As a result, the new admissible test-functions whose restriction to each edge is C^1 are applied. Independently and at the same time, Imbert *et al.* [17] proposed an equivalent notion of viscosity solution for studying a Hamilton-Jacobi approach to junction problems and traffic flows. Both [2] and [17] contain first results on comparison principles which were improved later. It is also worth mentioning the work by Schieborn and Camilli [22], in which the authors focus on eikonal equations on networks and on a less general notion of viscosity solution. In the particular case of eikonal equations, Camilli and Marchi established in [10] the equivalence between the definitions given in [2, 17, 22].

Since 2012, several proofs of comparison principles for HJB equations on networks, giving uniqueness of the solution, have been proposed.

1. In [3], Achdou *et al.* give a proof of a comparison principle for a stationary HJB equation arising from an optimal control with infinite horizon (therefore the Hamiltonian is convex) by mixing arguments from the theory of optimal control and PDE techniques. Such a proof was much inspired by works of Barles *et al.* [6, 7], on regional optimal control problems in \mathbb{R}^d (with discontinuous dynamics and costs).
2. A different and more general proof, using only arguments from the theory of PDEs was obtained by Imbert and Monneau in [16]. The proof works for quasi-convex Hamiltonians, and for stationary and time-dependent HJB equations. It relies on the construction of suitable *vertex test functions*.
3. A very simple and elegant proof, working for non convex Hamiltonians, has been very recently given by Lions and Souganidis [19, 20].

The goal of this paper is to consider an optimal control problem on a network in which there are entry (or exit) costs at each edge of the network and to study the related HJB equations. The effect of the entry/exit costs is to make the value function of the problem discontinuous. Discontinuous solutions of Hamilton-Jacobi equation have been studied by various authors, see for example Barles [4], Frankowska and Mazzola [13], and in particular Graber *et al.* [15] for different HJB equations on networks with discontinuous solutions.


To simplify the problem, we will first study the case of junction, *i.e.*, a network of the form $\mathcal{G} = \cup_{i=1}^N \Gamma_i$ with N edges Γ_i (Γ_i is the closed half line $\mathbb{R}^+ e_i$) and only one vertex O , where $\{O\} = \cap_{i=1}^N \Gamma_i$. Later, we will generalize our analysis to networks with an arbitrary number of vertices. In the case of the junction described above, our assumptions about the dynamics and the running costs are similar to those made in [3], except that additional costs c_i for entering the edge Γ_i at O or d_i for exiting Γ_i at O are added in the cost functional. Accordingly, the value function is continuous on $\mathcal{G} \setminus \{O\}$, but is in general discontinuous at the vertex O . Hence, instead of considering the value function \mathbf{v} , we split it into the collection $(v_i)_{1 \leq i \leq N}$, where v_i is continuous function defined on the edge Γ_i . More precisely,

$$v_i(x) = \begin{cases} \mathbf{v}(x) & \text{if } x \in \Gamma_i \setminus \{O\}, \\ \lim_{\delta \rightarrow 0^+} \mathbf{v}(\delta e_i) & \text{if } x = O. \end{cases}$$

Our approach is therefore reminiscent of optimal switching problems (impulsional control): in the present case the switches can only occur at the vertex O . Note that our assumptions will ensure that $\mathbf{v}|_{\Gamma_i \setminus \{O\}}$ is Lipschitz continuous near O and that $\lim_{\delta \rightarrow 0^+} \mathbf{v}(\delta e_i)$ does exist. In the case of entry costs for example, our first main result will be to find the relation between $\mathbf{v}(O)$, $v_i(O)$ and $v_j(O) + c_j$ for $i, j = \overline{1, N}$.

This will show that the functions $(v_i)_{1 \leq i \leq N}$ are (suitably defined) viscosity solutions of the following system

$$\begin{aligned} \lambda u_i(x) + H_i\left(x, \frac{du_i}{dx_i}(x)\right) &= 0 \quad \text{if } x \in \Gamma_i \setminus \{O\}, \\ \lambda u_i(O) + \max\left\{-\lambda \min_{j \neq i} \{u_j(O) + c_j\}, H_i^+\left(O, \frac{du_i}{dx_i}(O)\right), H_O^T\right\} &= 0 \quad \text{if } x = O. \end{aligned} \quad (1.1)$$

FIGURE 1. The network \mathcal{G} ($N = 5$).

Here H_i is the Hamiltonian corresponding to edge Γ_i . At vertex O , the definition of the Hamiltonian has to be particular, in order to consider all the possibilities when x is close to O . More specifically, if x is close to O and belongs to Γ_i then:

- The term $\min_{j \neq i} \{u_j(O) + c_j\}$ accounts for situations in which the trajectory enters Γ_{i_0} where $u_{i_0}(O) + c_{i_0} = \min_{j \neq i} \{u_j(O) + c_j\}$.
- The term $H_i^+ \left(O, \frac{du_i}{dx_i}(O) \right)$ accounts for situations in which the trajectory does not leave Γ_i .
- The term H_O^T accounts for situations in which the trajectory stays at O .

The most important part of the paper will be devoted to two different proofs of a comparison principle leading to the well-posedness of (1.1): the first one uses arguments from optimal control theory coming from Barles *et al.* [6, 7] and Achdou *et al.* [3]; the second one is inspired by Lions and Souganidis [19] and uses arguments from the theory of PDEs.

The paper is organized as follows: Section 2 deals with the optimal control problems with entry and exit costs: we give a simple example in which the value function is discontinuous at the vertex O , and also prove results on the structure of the value function near O . In Section 3, the new system of (1.1) is defined and a suitable notion of viscosity solutions is proposed. In Section 4, we prove our value functions are viscosity solutions of the above mentioned system. In Section 5, some properties of viscosity sub and super-solution are given and used to obtain the comparison principle. Finally, optimal control problems with entry costs which may be zero and related HJB equations are considered in Section 6.

2. OPTIMAL CONTROL PROBLEM ON JUNCTION WITH ENTRY/EXIT COSTS

2.1. The geometry

We consider the model case of the junction in \mathbb{R}^d with N semi-infinite straight edges, $N > 1$. The edges are denoted by $(\Gamma_i)_{i=\overline{1,N}}$ where Γ_i is the closed half-line \mathbb{R}^+e_i . The vectors e_i are two by two distinct unit vectors in \mathbb{R}^d . The half-lines Γ_i are glued at the vertex O to form the junction \mathcal{G}

$$\mathcal{G} = \bigcup_{i=1}^N \Gamma_i.$$

The geodetic distance $d(x, y)$ between two points x, y of \mathcal{G} is

$$d(x, y) = \begin{cases} |x - y| & \text{if } x, y \text{ belong to the same edge } \Gamma_i, \\ |x| + |y| & \text{if } x, y \text{ belong to different edges } \Gamma_i \text{ and } \Gamma_j. \end{cases}$$

2.2. The optimal control problem

We consider infinite horizon optimal control problems which have different dynamic and running costs for each and every edge. For $i = \overline{1, N}$,

- the set of control on Γ_i is denoted by A_i
- the system is driven by a dynamics f_i
- there is a running cost ℓ_i .

Our main assumptions, referred to as $[H]$ hereafter, are as follows:

$[H0]$ (**Control sets**) Let A be a metric space (one can take $A = \mathbb{R}^d$). For $i = \overline{1, N}$, A_i is a nonempty compact subset of A and the sets A_i are disjoint.

$[H1]$ (**Dynamics**) For $i = \overline{1, N}$, the function $f_i : \Gamma_i \times A_i \rightarrow \mathbb{R}$ is continuous and bounded by M . Moreover, there exists $L > 0$ such that

$$|f_i(x, a) - f_i(y, a)| \leq L|x - y| \quad \text{for all } x, y \in \Gamma_i, a \in A_i.$$

Hereafter, we will use the notation $F_i(x)$ for the set $\{f_i(x, a)e_i : a \in A_i\}$.

$[H2]$ (**Running costs**) For $i = \overline{1, N}$, the function $\ell_i : \Gamma_i \times A_i \rightarrow \mathbb{R}$ is a continuous function bounded by $M > 0$. There exists a modulus of continuity ω such that

$$|\ell_i(x, a) - \ell_i(y, a)| \leq \omega(|x - y|) \quad \text{for all } x, y \in \Gamma_i, a \in A_i.$$

$[H3]$ (**Convexity of dynamic and costs**) For $x \in \Gamma_i$, the following set

$$FL_i(x) = \{(f_i(x, a)e_i, \ell_i(x, a)) : a \in A_i\}$$

is non-empty, closed and convex.

$[H4]$ (**Strong controllability**) There exists a real number $\delta > 0$ such that

$$[-\delta e_i, \delta e_i] \subset F_i(O) = \{f_i(O, a)e_i : a \in A_i\}.$$

Remark 2.1. The assumption that the sets A_i are disjoint is not restrictive. Indeed, if A_i are not disjoint, then we define $\tilde{A}_i = A_i \times \{i\}$ and $\tilde{f}_i(x, \tilde{a}) = f_i(x, a)$, $\tilde{\ell}_i(x, \tilde{a}) = \ell_i(x, a)$ with $\tilde{a} = (a, i)$ with $a \in A_i$. The assumption

[H3] is made to avoid the use of relaxed control. With assumption [H4], one gets that the Hamiltonian which will appear later is coercive for x close to the O . Moreover, [H4] is an important assumption to prove Lemmas 2.7 and 5.3.

Let

$$\mathcal{M} = \{(x, a) : x \in \mathcal{G}, a \in A_i \text{ if } x \in \Gamma_i \setminus \{O\}, \text{ and } a \in \cup_{i=1}^N A_i \text{ if } x = O\}.$$

Then \mathcal{M} is closed. We also define the function on \mathcal{M} by

$$\text{for all } (x, a) \in \mathcal{M}, \quad f(x, a) = \begin{cases} f_i(x, a) e_i & \text{if } x \in \Gamma_i \setminus \{O\} \text{ and } a \in A_i, \\ f_i(O, a) e_i & \text{if } x = O \text{ and } a \in A_i. \end{cases}$$

The function f is continuous on \mathcal{M} since the sets A_i are disjoint.

Definition 2.2 (*The speed set and the admissible control set*). The set $\tilde{F}(x)$ which contains all the “possible speeds” at x is defined by

$$\tilde{F}(x) = \begin{cases} F_i(x) & \text{if } x \in \Gamma_i \setminus \{O\}, \\ \cup_{i=1}^N F_i(O) & \text{if } x = O. \end{cases}$$

For $x \in \mathcal{G}$, the set of admissible trajectories starting from x is

$$Y_x = \left\{ y_x \in Lip(\mathbb{R}^+; \mathcal{G}) : \begin{cases} \dot{y}_x(t) \in \tilde{F}(y_x(t)) & \text{for a.e. } t > 0 \\ y_x(0) = x \end{cases} \right\}.$$

According to Theorem 1.2 from [3], a solution y_x can be associated with several control laws. We introduce the set of admissible controlled trajectories starting from x

$$\mathcal{T}_x = \left\{ (y_x, \alpha) \in L_{loc}^\infty(\mathbb{R}^+; \mathcal{M}) : y_x \in Lip(\mathbb{R}^+; \mathcal{G}) \text{ and } y_x(t) = x + \int_0^t f(y_x(s), \alpha(s)) ds \right\}.$$

Notice that, if $(y_x, \alpha) \in \mathcal{T}_x$ then $y_x \in Y_x$. Hereafter, we will denote y_x by $y_{x,\alpha}$ if $(y_x, \alpha) \in \mathcal{T}_x$. For any $y_{x,\alpha}$, we can define the closed set $T_O = \{t \in \mathbb{R}^+ : y_{x,\alpha}(t) = O\}$ and the open set T_i in $\mathbb{R}^+ = [0, +\infty)$ by $T_i = \{t \in \mathbb{R}^+ : y_{x,\alpha}(t) \in \Gamma_i \setminus \{O\}\}$. The set T_i is a countable union of disjoint open intervals

$$T_i = \bigcup_{k \in K_i \subset \mathbb{N}} T_{ik} = \begin{cases} [0, \eta_{i0}) \cup \bigcup_{k \in K_i \subset \mathbb{N}^*} (t_{ik}, \eta_{ik}) & \text{if } x \in \Gamma_i \setminus \{O\}, \\ \bigcup_{k \in K_i \subset \mathbb{N}^*} (t_{ik}, \eta_{ik}) & \text{if } x \notin \Gamma_i \setminus \{O\}, \end{cases}$$

where $K_i = \overline{1, n}$ if the trajectory $y_{x,\alpha}$ enters Γ_i n times and $K_i = \mathbb{N}$ if the trajectory $y_{x,\alpha}$ enters Γ_i infinite times.

Remark 2.3. From the above definition, one can see that t_{ik} is an entry time in $\Gamma_i \setminus \{O\}$ and η_{ik} is an exit time from $\Gamma_i \setminus \{O\}$. Hence

$$y_{x,\alpha}(t_{ik}) = y_{x,\alpha}(\eta_{ik}) = O.$$

Let $C = \{c_1, c_2, \dots, c_N\}$ be a set of **entry costs** and $D = \{d_1, d_2, \dots, d_N\}$ be a set of **exit costs**. We underline that, except in Section 6, entry and exist costs are positive.

In the sequel, we define two different *cost functionals* (the first one corresponds to the case when there is a cost for entering the edges and the second one corresponds to the case when there is a cost for exiting the edges):

Definition 2.4 (The cost functionals and value functions with entry/exit costs). The costs associated to trajectory $(y_{x,\alpha}, \alpha) \in \mathcal{T}_x$ are defined by

$$J(x; (y_{x,\alpha}, \alpha)) = \int_0^{+\infty} \ell(y_{x,\alpha}(t), \alpha(t)) e^{-\lambda t} dt + \sum_{i=1}^N \sum_{k \in K_i} c_i e^{-\lambda t_{ik}} \quad (\text{cost functional with entry cost}),$$

and

$$\widehat{J}(x; (y_{x,\alpha}, \alpha)) = \int_0^{+\infty} \ell(y_{x,\alpha}(t), \alpha(t)) e^{-\lambda t} dt + \sum_{i=1}^N \sum_{k \in K_i} d_i e^{-\lambda \eta_{ik}} \quad (\text{cost functional with exit cost}),$$

where the running cost $\ell : \mathcal{M} \rightarrow \mathbb{R}$ is

$$\ell(x, a) = \begin{cases} \ell_i(x, a) & \text{if } x \in \Gamma_i \setminus \{O\} \text{ and } a \in A_i, \\ \ell_i(O, a) & \text{if } x = 0 \text{ and } a \in A_i. \end{cases}$$

Hereafter, to simplify the notation, we will use $J(x, \alpha)$ and $\widehat{J}(x, \alpha)$ instead of $J(x; (y_{x,\alpha}, \alpha))$ and $\widehat{J}(x; (y_{x,\alpha}, \alpha))$, respectively.

The value functions of the infinite horizon optimal control problem are defined by:

$$v(x) = \inf_{(y_{x,\alpha}, \alpha) \in \mathcal{T}_x} J(x; (y_{x,\alpha}, \alpha)) \quad (\text{value function with entry cost}),$$

and

$$\widehat{v}(x) = \inf_{(y_{x,\alpha}, \alpha) \in \mathcal{T}_x} \widehat{J}(x; (y_{x,\alpha}, \alpha)) \quad (\text{value function with exit cost}).$$

Remark 2.5. By the definition of the value function, we are mainly interested in a control law α such that $J(x, \alpha) < +\infty$. In such a case, if $|K_i| = +\infty$, then we can order $\{t_{ik}, \eta_{ik} : k \in \mathbb{N}\}$ such that

$$t_{i1} < \eta_{i1} < t_{i2} < \eta_{i2} < \cdots < t_{ik} < \eta_{ik} < \cdots,$$

and

$$\lim_{k \rightarrow \infty} t_{ik} = \lim_{k \rightarrow \infty} \eta_{ik} = +\infty.$$

Indeed, assuming if $\lim_{k \rightarrow \infty} t_{ik} = \bar{t} < +\infty$, then

$$J(x, \alpha) \geq -\frac{M}{\lambda} + \sum_{k=1}^{+\infty} e^{-\lambda t_{ik}} c_i = -\frac{M}{\lambda} + c_i \sum_{k=1}^{+\infty} e^{-\lambda t_{ik}} = +\infty,$$

in contradiction with $J(x, \alpha) < +\infty$. This means that the state cannot switch edges infinitely many times in finite time, otherwise the cost functional is obviously infinite.

The following example shows that the value function with entry costs is possibly discontinuous (the same holds for the value function with exit costs).

Example 2.6. Consider the network $\mathcal{G} = \Gamma_1 \cup \Gamma_2$ where $\Gamma_1 = \mathbb{R}^+ e_1 = (-\infty, 0]$ and $\Gamma_2 = \mathbb{R}^+ e_2 = [0, +\infty)$. The control sets are $A_i = [-1, 1] \times \{i\}$ with $i \in \{1, 2\}$. Set

$$(f(x, a), \ell(x, a)) = \begin{cases} (f_i(x, (a_i, i)) e_i, \ell_i(x, (a_i, i))) & \text{if } x \in \Gamma_i \setminus \{O\} \text{ and } a = (a_i, i) \in A_i, \\ (f_i(O, (a_i, i)) e_i, \ell_i(O, (a_i, i))) & \text{if } x = O \text{ and } a = (a_i, i) \in A_i, \end{cases}$$

where $f_i(x, (a_i, i)) = a_i$ and $\ell_1 \equiv 1, \ell_2(x, (a_2, 2)) = 1 - a_2$. For $x \in \Gamma_2 \setminus \{O\}$, then $v(x) = v_2(x) = 0$ with optimal strategy consists in choosing $\alpha(t) \equiv (1, 2)$. For $x \in \Gamma_1$, we can check that $v(x) = \min \left\{ \frac{1}{\lambda}, \frac{1 - e^{-\lambda|x|}}{\lambda} + c_2 e^{-\lambda|x|} \right\}$. More precisely, for all $x \in \Gamma_1$, we have

$$v(x) = \begin{cases} \frac{1}{\lambda} & \text{if } c_2 \geq \frac{1}{\lambda}, \text{ with the optimal control } \alpha(t) \equiv (-1, 1), \\ \frac{1 - e^{-\lambda|x|}}{\lambda} + c_2 e^{-\lambda|x|} & \text{if } c_2 < \frac{1}{\lambda}, \text{ with the optimal control } \alpha(t) = \begin{cases} (1, 1) & \text{if } t \leq |x|, \\ (1, 2) & \text{if } t \geq |x|. \end{cases} \end{cases}$$

Summarizing, we have the two following cases

1. If $c_2 \geq \frac{1}{\lambda}$, then

$$v(x) = \begin{cases} 0 & \text{if } x \in \Gamma_2 \setminus \{O\}, \\ \frac{1}{\lambda} & \text{if } x \in \Gamma_1. \end{cases}$$

The graph of the value function with entry costs $c_2 \geq \frac{1}{\lambda} = 1$ is plotted in Figure 2a.

2. If $c_2 < \frac{1}{\lambda}$, then

$$v(x) = \begin{cases} 0 & \text{if } x \in \Gamma_2 \setminus \{O\}, \\ \frac{1 - e^{-\lambda|x|}}{\lambda} + c_2 e^{-\lambda|x|} & \text{if } x \in \Gamma_1. \end{cases}$$

The graph of the value function with entry costs $c_2 = \frac{1}{2} < 1 = \frac{1}{\lambda}$ is plotted in Figure 2b.

Lemma 2.7. Under assumptions [H1] and [H4], there exist two positive numbers r_0 and C such that for all $x_1, x_2 \in B(O, r_0) \cap \mathcal{G}$, there exists $(y_{x_1, \alpha_{x_1, x_2}}, \alpha_{x_1, x_2}) \in \mathcal{T}_{x_1}$ and $\tau_{x_1, x_2} \leq Cd(x_1, x_2)$ such that $y_{x_1}(\tau_{x_1, x_2}) = x_2$.

Proof of Lemma 2.7. This proof is classical. It is sufficient to consider the case when x_1 and x_2 belong to same edge Γ_i , since in the other cases, we will use O as a connecting point between x_1 and x_2 . According to Assumption [H4], there exists $a \in A_i$ such that $f_i(O, a) = \delta$. Additionally, by the Lipschitz continuity of f_i ,

$$|f_i(O, a) - f_i(x, a)| \leq L|x|,$$

hence, if we choose $r_0 := \frac{\delta}{2L} > 0$, then $f_i(x, a) \geq \frac{\delta}{2}$ for all $x \in B(O, r_0) \cap \Gamma_i$. Let x_1, x_2 be in $B(O, r_0) \cap \Gamma_i$ with $|x_1| < |x_2|$: there exist a control law α and $\tau_{x_1, x_2} > 0$ such that $\alpha(t) = a$ if $0 \leq t \leq \tau_{x_1, x_2}$ and




FIGURE 2. An example of value function with entry cost.

$y_{x_1, \alpha}(\tau_{x_1, x_2}) = x_2$. Moreover, since the velocity $f_i(y_{x_1, \alpha}(t), \alpha(t))$ is always greater than $\frac{\delta}{2}$ when $t \leq \tau_{x_1, x_2}$, then $\tau_{x_1, x_2} \leq \frac{2}{\delta}d(x_1, x_2)$. If $|x_1| > |x_2|$, the proof is achieved by replacing $a \in A_i$ by $\bar{a} \in A_i$ such that $f_i(O, \bar{a}) = -\delta$ and applying the same argument as above. \square

2.3. Some properties of value function at the vertex

Lemma 2.8. *Under assumption [H], $v|_{\Gamma_i \setminus \{O\}}$ and $\widehat{v}|_{\Gamma_i \setminus \{O\}}$ are continuous for any $i = \overline{1, N}$. Moreover, there exists $\varepsilon > 0$ such that $v|_{\Gamma_i \setminus \{O\}}$ and $\widehat{v}|_{\Gamma_i \setminus \{O\}}$ are Lipschitz continuous in $(\Gamma_i \setminus \{O\}) \cap B(O, \varepsilon)$. Hence, it is possible to extend $v|_{\Gamma_i \setminus \{O\}}$ and $\widehat{v}|_{\Gamma_i \setminus \{O\}}$ at O into Lipschitz continuous functions in $\Gamma_i \cap B(O, \varepsilon)$. Hereafter, v_i and \widehat{v}_i denote these extensions.*

Proof of Lemma 2.8. The proof of continuity inside the edge is classical by using [H4], see [1] for more details. The proof of Lipschitz continuity is a consequence of Lemma 2.7. Indeed, for x, y belong to $\Gamma_i \cap B(O, \varepsilon)$, by Lemma 2.7 and the definition of value function, we have

$$v(x) - v(z) = v_i(x) - v_i(z) \leq \int_0^{\tau_{x,z}} \ell_i(y_{x, \alpha_{x,z}}(t), \alpha_{x,z}(t)) e^{-\lambda t} dt + v_i(z) (e^{-\lambda \tau_{x,z}} - 1).$$

Since ℓ_i is bounded by M (by [H2]), v_i is bounded in $\Gamma_i \cap B(O, \varepsilon)$ and $e^{-\lambda \tau_{x,z}} - 1$ is bounded by $\tau_{x,y}$, there exists a constant \overline{C} such that

$$v_i(x) - v_i(z) \leq \overline{C} \tau_{x,z} \leq \overline{C} C |x - z|.$$

The last inequality follows from the Lemma 2.7. The inequality $v_i(z) - v_i(x) \leq \overline{C} C |x - z|$ is obtained in a similar way. The proof is done. \square

Let us define the tangential Hamiltonian H_O^T at vertex O by

$$H_O^T = \max_{i=1, \overline{N}} \max_{a_i \in A_i^O} \{-\ell_j(O, a_j)\} = - \min_{i=1, \overline{N}} \min_{a_i \in A_i^O} \{\ell_j(O, a_j)\}, \quad (2.1)$$

where $A_i^O = \{a_i \in A_i : f_i(O, a_i) = 0\}$. The relationship between the values $v(O)$, $v_i(O)$ and H_O^T will be given in the next theorem. Hereafter, the proofs of the results will be supplied only for the value function with entry costs v , the proofs concerning the value function with exit costs \widehat{v} are totally similar.

Theorem 2.9. *Under assumption [H], the value functions v and \widehat{v} satisfy*

$$v(O) = \min \left\{ \min_{i=1, \overline{N}} \{v_i(O) + c_i\}, -\frac{H_O^T}{\lambda} \right\},$$

and

$$\widehat{v}(O) = \min \left\{ \min_{i=1, \overline{N}} \{\widehat{v}_i(O)\}, -\frac{H_O^T}{\lambda} \right\}.$$

Remark 2.10. Theorem 2.9 gives us the characterization of the value function at vertex O .

The proof of Theorem 2.9, makes use of Lemmas 2.11 and 2.12.

Lemma 2.11 (Value functions v and \widehat{v} at O). *Under assumption [H], then*

$$\max_{i=1, \overline{N}} \{v_i(O)\} \leq v(O) \leq \min_{i=1, \overline{N}} \{v_i(O) + c_i\},$$

and

$$\max_{i=1, \overline{N}} \{\widehat{v}_i(O) - d_i\} \leq \widehat{v}(O) \leq \min_{i=1, \overline{N}} \{\widehat{v}_i(O)\}.$$

Proof of Lemma 2.11. We divide the proof into two parts.

Prove that $\max_{i=1, \overline{N}} \{v_i(O)\} \leq v(O)$. First, we fix $i \in \{1, \dots, N\}$ and any control law $\bar{\alpha}$ such that $(y_{O, \bar{\alpha}}, \bar{\alpha}) \in \mathcal{T}_O$. Let $x \in \Gamma_i \setminus \{O\}$ such that $|x|$ is small. From Lemma 2.7, there exists a control law $\alpha_{x, O}$ connecting x and O and we consider

$$\alpha(s) = \begin{cases} \alpha_{x, O}(s) & \text{if } s \leq \tau_{x, O}, \\ \bar{\alpha}(s - \tau_{x, O}) & \text{if } s > \tau_{x, O}. \end{cases}$$

It means that the trajectory goes from x to O with the control law $\alpha_{x, O}$ and then proceeds with the control law $\bar{\alpha}$. Therefore

$$v(x) = v_i(x) \leq J(x, \alpha) = \int_0^{\tau_{x, O}} \ell_i(y_{x, \alpha}(s)) e^{-\lambda s} ds + e^{-\lambda \tau_{x, O}} J(O, \bar{\alpha}).$$

Since $\bar{\alpha}$ is chosen arbitrarily and ℓ_i is bounded by M , we get

$$v_i(x) \leq M\tau_{x, O} + e^{-\lambda \tau_{x, O}} v(O).$$

Let x tend to O then $\tau_{x,O}$ tend to 0 from Lemma 2.7. Therefore, $v_i(O) \leq v(O)$. Since the above inequality holds for $i = \overline{1, N}$, we obtain that

$$\max_{i=\overline{1, N}} \{v_i(O)\} \leq v(O).$$

Prove that $v(O) \leq \min_{i=\overline{1, N}} \{v_i(O) + c_i\}$. For $i = \overline{1, N}$; we claim that $v(O) \leq v_i(O) + c_i$. Consider $x \in \Gamma_i \setminus \{O\}$ with $|x|$ small enough and any control law $\bar{\alpha}_x$ such that $(y_{x, \bar{\alpha}_x}, \bar{\alpha}_x) \in \mathcal{T}_x$. From Lemma 2.7, there exists a control law $\alpha_{O,x}$ connecting O and x and we consider

$$\alpha(s) = \begin{cases} \alpha_{O,x}(s) & \text{if } s \leq \tau_{O,x}, \\ \bar{\alpha}_x(s - \tau_{O,x}) & \text{if } s > \tau_{O,x}. \end{cases}$$

It means that the trajectory goes from O to x using the control law $\alpha_{O,x}$ then proceeds with the control law $\bar{\alpha}_x$. Therefore

$$v(O) \leq J(O, \alpha) = c_i + \int_0^{\tau_{O,x}} \ell_i(y_{O,\alpha}(s)) e^{-\lambda s} ds + e^{-\lambda \tau_{O,x}} J(x, \bar{\alpha}_x).$$

Since $\bar{\alpha}_x$ is chosen arbitrarily and ℓ_i is bounded by M , we get

$$v(O) \leq c_i + M\tau_{O,x} + e^{-\lambda \tau_{O,x}} v_i(x)$$

Let x tend to O then $\tau_{O,x}$ tends to 0 from Lemma 2.7, then $v(O) \leq c_i + v_i(O)$. Since the above inequality holds for $i = \overline{1, N}$, we obtain that

$$v(O) \leq \min_{i=\overline{1, N}} \{v_i(O) + c_i\}.$$

□

Lemma 2.12. *The value functions v and \hat{v} satisfy*

$$v(O), \hat{v}(O) \leq -\frac{H_O^T}{\lambda} \tag{2.2}$$

where H_O^T is defined in (2.1).

Proof of Lemma 2.12. From (2.1), there exists $j \in \{1, \dots, N\}$ and $a_j \in A_j^O$ such that

$$H_O^T = -\min_{i=\overline{1, N}} \min_{a_i \in A_i^O} \{\ell_i(O, a_i)\} = -\ell_j(O, a_j)$$

Let the control law α be defined by $\alpha(s) \equiv a_j$ for all s , then

$$v(O) \leq J(O, \alpha) = \int_0^{+\infty} \ell_j(O, a_j) e^{-\lambda s} ds = \frac{\ell_j(O, a_j)}{\lambda} = -\frac{H_O^T}{\lambda}.$$

□

We are ready to prove Theorem 2.9.

Proof of Theorem 2.9. According to Lemma 2.11 and Lemma 2.12,

$$v(O) \leq \min \left\{ \min_{i=1, N} \{v_i(O) + c_i\}, -\frac{H_O^T}{\lambda} \right\}.$$

Assuming that

$$v(O) < \min_{i=1, N} \{v_i(O) + c_i\}, \quad (2.3)$$

it is sufficient to prove that $v(O) = -\frac{H_O^T}{\lambda}$. By (2.3), there exists a sequence $\{\varepsilon_n\}_{n \in \mathbb{N}}$ such that $\varepsilon_n \rightarrow 0$ and

$$v(O) + \varepsilon_n < \min_{i=1, N} \{v_i(O) + c_i\} \quad \text{for all } n \in \mathbb{N}.$$

On the other hand, there exists an ε_n -optimal control α_n , $v(O) + \varepsilon_n > J(O, \alpha_n)$. Let us define the first time that the trajectory y_{O, α_n} leaves O

$$t_n := \inf_{i=1, N} T_i^n,$$

where T_i^n is the set of times t for which $y_{O, \alpha_n}(t)$ belongs to $\Gamma_i \setminus \{O\}$. Notice that t_n is possibly $+\infty$, in which case $y_{O, \alpha_n}(s) = O$ for all $s \in [0, +\infty)$. Extracting a subsequence if necessary, we may assume that t_n tends to $\bar{t} \in [0, +\infty]$ when ε_n tends to 0.

If there exists a subsequence of $\{t_n\}_{n \in \mathbb{N}}$ (which is still noted $\{t_n\}_{n \in \mathbb{N}}$) such that $t_n = +\infty$ for all $n \in \mathbb{N}$, then for a.e. $s \in [0, +\infty)$

$$\begin{cases} f(y_{O, \alpha_n}(s), \alpha_n(s)) = f(O, \alpha_n(s)) = 0, \\ \ell(y_{O, \alpha_n}(s), \alpha_n(s)) = \ell(O, \alpha_n(s)). \end{cases}$$

In this case, $\alpha_n(s) \in \cup_{i=1}^N A_i^O$ for a.e. $s \in [0, +\infty)$. Therefore, for a.e. $s \in [0, +\infty)$

$$\ell(y_{O, \alpha_n}(s), \alpha_n(s)) = \ell(O, \alpha_n(s)) \geq -H_O^T,$$

and

$$v(O) + \varepsilon_n > J(O, \alpha_n) = \int_0^{+\infty} \ell(O, \alpha_n(s)) e^{-\lambda s} ds \geq \int_0^{+\infty} (-H_O^T) e^{-\lambda s} ds = -\frac{H_O^T}{\lambda}.$$

By letting n tend to ∞ , we get $v(O) \geq -\frac{H_O^T}{\lambda}$. On the other hand, since $v(O) \leq -\frac{H_O^T}{\lambda}$ by Lemma 2.12, this implies that $v(O) = -\frac{H_O^T}{\lambda}$.

Let us now assume that $0 \leq t_n < +\infty$ for all n large enough. Then, for a fixed n and for any positive $\delta \leq \delta_n$ where δ_n small enough, $y_{O, \alpha_n}(s)$ still belongs to some $\Gamma_{i(n)} \setminus \{O\}$ for all $s \in (t_n, t_n + \delta]$. We have

$$\begin{aligned}
v(O) + \varepsilon_n &> J(O, \alpha_n) \\
&= \int_0^{t_n} \ell(y_{O, \alpha_n}(s), \alpha_n(s)) e^{-\lambda s} ds + c_{i(n)} e^{-\lambda t_n} + \int_{t_n}^{t_n + \delta} \ell_{i(n)}(y_{O, \alpha_n}(s), \alpha_n(s)) e^{-\lambda s} ds \\
&\quad + e^{-\lambda(t_n + \delta)} J(y_{O, \alpha_n}(t_n + \delta), \alpha_n(\cdot + t_n + \delta)) \\
&\geq \int_0^{t_n} \ell(y_{O, \alpha_n}(s), \alpha_n(s)) e^{-\lambda s} ds + c_{i(n)} e^{-\lambda t_n} + \int_{t_n}^{t_n + \delta} \ell_{i(n)}(y_{O, \alpha_n}(s), \alpha_n(s)) e^{-\lambda s} ds \\
&\quad + e^{-\lambda(t_n + \delta)} v(y_{O, \alpha_n}(t_n + \delta)) \\
&= \int_0^{t_n} \ell(y_{O, \alpha_n}(s), \alpha_n(s)) e^{-\lambda s} ds + c_{i(n)} e^{-\lambda t_n} + \int_{t_n}^{t_n + \delta} \ell_{i(n)}(y_{O, \alpha_n}(s), \alpha_n(s)) e^{-\lambda s} ds \\
&\quad + e^{-\lambda(t_n + \delta)} v_{i(n)}(y_{O, \alpha_n}(t_n + \delta)).
\end{aligned}$$

By letting δ tend to 0,

$$v(O) + \varepsilon_n \geq \int_0^{t_n} \ell(y_{O, \alpha_n}(s), \alpha_n(s)) e^{-\lambda s} ds + c_{i(n)} e^{-\lambda t_n} + e^{-\lambda t_n} v_{i(n)}(O).$$

Note that $y_{O, \alpha_n}(s) = O$ for all $s \in [0, t_n]$, i.e., $f(O, \alpha_n(s)) = 0$ a.e. $s \in [0, t_n]$. Hence

$$\begin{aligned}
v(O) + \varepsilon_n &\geq \int_0^{t_n} \ell(O, \alpha_n(s)) e^{-\lambda s} ds + c_{i(n)} e^{-\lambda t_n} + e^{-\lambda t_n} v_{i(n)}(O) \\
&\geq \int_0^{t_n} (-H_O^T) e^{-\lambda s} ds + c_{i(n)} e^{-\lambda t_n} + e^{-\lambda t_n} v_{i(n)}(O) \\
&= \frac{1 - e^{-\lambda t_n}}{\lambda} (-H_O^T) + c_{i(n)} e^{-\lambda t_n} + e^{-\lambda t_n} v_{i(n)}(O).
\end{aligned}$$

Choose a subsequence $\{\varepsilon_{n_k}\}_{k \in \mathbb{N}}$ of $\{\varepsilon_n\}_{n \in \mathbb{N}}$ such that for some $i_0 \in \{1, \dots, N\}$, $c_{i(n_k)} = c_{i_0}$ for all k . By letting k tend to ∞ , recall that $\lim_{k \rightarrow \infty} t_{n_k} = \bar{t}$, we have three possible cases

1. If $\bar{t} = +\infty$, then $v(O) \geq -\frac{H_O^T}{\lambda}$. By Lemma 2.12, we obtain $v(O) = -\frac{H_O^T}{\lambda}$.
2. If $\bar{t} = 0$, then $v(O) \geq c_{i_0} + v_{i_0}(O)$. By (2.3), we obtain a contradiction.
3. If $\bar{t} \in (0, +\infty)$, then $v(O) \geq \frac{1 - e^{-\lambda \bar{t}}}{\lambda} (-H_O^T) + [c_{i_0} + v_{i_0}(O)] e^{-\lambda \bar{t}}$. By (2.3), $c_{i_0} + v_{i_0}(O) > v(O)$, so

$$v(O) > \frac{1 - e^{-\lambda \bar{t}}}{\lambda} (-H_O^T) + v(O) e^{-\lambda \bar{t}}.$$

This yields $v(O) > -\frac{H_O^T}{\lambda}$, and finally obtain a contradiction by Lemma 2.12.

□

3. THE HAMILTON-JACOBI SYSTEMS. VISCOSITY SOLUTIONS

3.1. Test-functions

Definition 3.1. A function $\varphi : \Gamma_1 \times \cdots \times \Gamma_N \rightarrow \mathbb{R}^N$ is an admissible test-function if there exists $(\varphi_i)_{i=\overline{1,N}}$, $\varphi_i \in C^1(\Gamma_i)$, such that $\varphi(x_1, \dots, x_N) = (\varphi_1(x_1), \dots, \varphi_N(x_N))$. The set of admissible test-function is denoted by $\mathcal{R}(\mathcal{G})$.

3.2. Definition of viscosity solution

Definition 3.2 (Hamiltonian). We define the Hamiltonian $H_i : \Gamma_i \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$H_i(x, p) = \max_{a \in A_i} \{-pf_i(x, a) - \ell_i(x, a)\}$$

and the Hamiltonian $H_i^+(O, \cdot) : \mathbb{R} \rightarrow \mathbb{R}$ by

$$H_i^+(O, p) = \max_{a \in A_i^+} \{-pf_i(O, a) - \ell_i(O, a)\},$$

where $A_i^+ = \{a_i \in A_i : f_i(O, a_i) \geq 0\}$. Recall that *the tangential Hamiltonian at O* , H_O^T , has been defined in (2.1).

We now introduce the Hamilton-Jacobi system for the case with entry costs

$$\begin{aligned} \lambda u_i(x) + H_i\left(x, \frac{du_i}{dx_i}(x)\right) &= 0 \quad \text{if } x \in \Gamma_i \setminus \{O\}, \\ \lambda u_i(O) + \max\left\{-\lambda \min_{j \neq i} \{u_j(O) + c_j\}, H_i^+\left(O, \frac{du_i}{dx_i}(O)\right), H_O^T\right\} &= 0 \quad \text{if } x = O, \end{aligned} \quad (3.1)$$

for all $i = \overline{1, N}$ and the Hamilton-Jacobi system with exit costs

$$\begin{aligned} \lambda \widehat{u}_i(x) + H_i\left(x, \frac{d\widehat{u}_i}{dx_i}(x)\right) &= 0 \quad \text{if } x \in \Gamma_i \setminus \{O\}, \\ \lambda \widehat{u}_i(O) + \max\left\{-\lambda \min_{j \neq i} \{\widehat{u}_j(O) + d_j\}, H_i^+\left(O, \frac{d\widehat{u}_i}{dx_i}(O)\right), H_O^T - \lambda d_i\right\} &= 0 \quad \text{if } x = O, \end{aligned} \quad (3.2)$$

for all $i = \overline{1, N}$ and their viscosity solutions.

Definition 3.3 (Viscosity solution with entry costs).

• A function $u := (u_1, \dots, u_N)$ where $u_i \in USC(\Gamma_i; \mathbb{R})$ for all $i = \overline{1, N}$, is called a *viscosity sub-solution* of (3.1) if for any $(\varphi_1, \dots, \varphi_N) \in \mathcal{R}(\mathcal{G})$, any $i = \overline{1, N}$ and any $x_i \in \Gamma_i$ such that $u_i - \varphi_i$ has a *local maximum point* on Γ_i at x_i , then

$$\begin{aligned} \lambda u_i(x_i) + H_i\left(x_i, \frac{d\varphi_i}{dx_i}(x_i)\right) &\leq 0 \quad \text{if } x_i \in \Gamma_i \setminus \{O\}, \\ \lambda u_i(O) + \max\left\{-\lambda \min_{j \neq i} \{u_j(O) + c_j\}, H_i^+\left(O, \frac{d\varphi_i}{dx_i}(O)\right), H_O^T\right\} &\leq 0 \quad \text{if } x_i = O. \end{aligned}$$

• A function $u := (u_1, \dots, u_N)$ where $u_i \in LSC(\Gamma_i; \mathbb{R})$ for all $i = \overline{1, N}$, is called a *viscosity super-solution* of (3.1) if for any $(\varphi_1, \dots, \varphi_N) \in \mathcal{R}(\mathcal{G})$, any $i = \overline{1, N}$ and any $x_i \in \Gamma_i$ such that $u_i - \varphi_i$ has a *local minimum*

point on Γ_i at x_i , then

$$\begin{aligned} \lambda u_i(x_i) + H_i\left(x_i, \frac{d\varphi_i}{dx_i}(x_i)\right) &\geq 0 \quad \text{if } x_i \in \Gamma_i \setminus \{O\}, \\ \lambda u_i(O) + \max\left\{-\lambda \min_{j \neq i} \{u_j(O) + c_j\}, H_i^+\left(O, \frac{d\varphi_i}{dx_i}(O)\right), H_O^T\right\} &\geq 0 \quad \text{if } x_i = O. \end{aligned}$$

• A functions $u := (u_1, \dots, u_N)$ where $u_i \in C(\Gamma_i; \mathbb{R})$ for all $i = \overline{1, N}$, is called a *viscosity solution* of (3.1) if it is both a viscosity sub-solution and a viscosity super-solution of (3.1).

Definition 3.4 (Viscosity solution with exit costs).

• A function $\hat{u} := (\hat{u}_1, \dots, \hat{u}_N)$ where $\hat{u}_i \in USC(\Gamma_i; \mathbb{R})$ for all $i = \overline{1, N}$, is called a *viscosity sub-solution* of (3.2) if for any $(\psi_1, \dots, \psi_N) \in \mathcal{R}(\mathcal{G})$, any $i = \overline{1, N}$ and any $y_i \in \Gamma_i$ such that $\hat{u}_i - \psi_i$ has a *local maximum point* on Γ_i at y_i , then

$$\begin{aligned} \lambda \hat{u}_i(y_i) + H_i\left(y_i, \frac{d\psi_i}{dx_i}(y_i)\right) &\leq 0 \quad \text{if } y_i \in \Gamma_i \setminus \{O\}, \\ \lambda \hat{u}_i(O) + \max\left\{-\lambda \min_{j \neq i} \{\hat{u}_j(O)\} - \lambda d_i, H_i^+\left(O, \frac{d\psi_i}{dx_i}(O)\right), H_O^T - \lambda d_i\right\} &\leq 0 \quad \text{if } y_i = O. \end{aligned}$$

• A function $\hat{u} := (\hat{u}_1, \dots, \hat{u}_N)$ where $\hat{u}_i \in LSC(\Gamma_i; \mathbb{R})$ for all $i = \overline{1, N}$, is called a *viscosity super-solution* of (3.2) if for any $(\psi_1, \dots, \psi_N) \in \mathcal{R}(\mathcal{G})$, any $i = \overline{1, N}$ and any $y_i \in \Gamma_i$ such that $\hat{u}_i - \psi_i$ has a *local minimum point* on Γ_i at y_i , then

$$\begin{aligned} \lambda \hat{u}_i(y_i) + H_i\left(y_i, \frac{d\psi_i}{dx_i}(y_i)\right) &\geq 0 \quad \text{if } y_i \in \Gamma_i \setminus \{O\}, \\ \lambda \hat{u}_i(O) + \max\left\{-\lambda \min_{j \neq i} \{\hat{u}_j(O)\} - \lambda d_i, H_i^+\left(O, \frac{d\psi_i}{dx_i}(O)\right), H_O^T - \lambda d_i\right\} &\geq 0 \quad \text{if } y_i = O. \end{aligned}$$

• A functions $\hat{u} := (\hat{u}_1, \dots, \hat{u}_N)$ where $\hat{u}_i \in C(\Gamma_i; \mathbb{R})$ for all $i = \overline{1, N}$, is called a *viscosity solution* of (3.2) if it is both a viscosity sub-solution and a viscosity super-solution of (3.2).

Remark 3.5. This notion of viscosity solution is consistent with the one of [3]. It can be seen in Section 6 when all the switching costs are zero, our definition and the one of [3] coincide.

4. CONNECTIONS BETWEEN THE VALUE FUNCTIONS AND THE HAMILTON-JACOBI SYSTEMS

Let v be the value function of the optimal control problem with entry costs and \hat{v} be a value function of the optimal control problem with exit costs. Recall that $v_i, \hat{v}_i : \Gamma_i \rightarrow \mathbb{R}$ are defined in Lemma 2.8 by

$$\begin{cases} v_i(x) = v(x) & \text{if } x \in \Gamma_i \setminus \{O\}, \\ v_i(O) = \lim_{\Gamma_i \setminus \{O\} \ni x \rightarrow O} v(x), \end{cases} \quad \text{and} \quad \begin{cases} \hat{v}_i(x) = \hat{v}(x) & \text{if } x \in \Gamma_i \setminus \{O\}, \\ \hat{v}_i(O) = \lim_{\Gamma_i \setminus \{O\} \ni x \rightarrow O} \hat{v}(x). \end{cases}$$

We wish to prove that $v := (v_1, v_2, \dots, v_N)$ and $\hat{v} := (\hat{v}_1, \dots, \hat{v}_N)$ are respectively viscosity solutions of (3.1) and (3.2). In fact, since $\mathcal{G} \setminus \{O\}$ is a finite union of open intervals in which the classical theory can be applied, we obtain that v_i and \hat{v}_i are viscosity solutions of

$$\lambda u(x) + H_i(x, Du(x)) = 0 \quad \text{in } \Gamma_i \setminus \{O\}.$$

Therefore, we can restrict ourselves to prove the following theorem.

Theorem 4.1. *For $i = \overline{1, N}$, the function v_i satisfies*

$$\lambda v_i(O) + \max \left\{ -\lambda \min_{j \neq i} \{v_j(O) + c_j\}, H_i^+ \left(O, \frac{dv_i}{dx_i}(O) \right), H_O^T \right\} = 0$$

in the viscosity sense. The function \widehat{v}_i satisfies

$$\lambda \widehat{v}_i(O) + \max \left\{ -\lambda \min_{j \neq i} \{\widehat{v}_j(O) + d_j\}, H_i^+ \left(O, \frac{d\widehat{v}_i}{dx_i}(O) \right), H_O^T - \lambda d_i \right\} = 0$$

in the viscosity sense.

The proof of Theorem 4.1 follows from Lemmas 4.2 and 4.5. We focus on v_i since the proof for \widehat{v}_i is similar.

Lemma 4.2. *For $i = \overline{1, N}$, the function v_i is a viscosity sub-solution of (3.1) at O .*

Proof of Lemma 4.2. From Theorem 2.9,

$$\lambda v_i(O) + \max \left\{ -\lambda \min_{j \neq i} \{v_j(O) + c_j\}, H_O^T \right\} \leq 0.$$

It is thus sufficient to prove that

$$\lambda v_i(O) + H_i^+ \left(O, \frac{dv_i}{dx_i}(O) \right) \leq 0$$

in the viscosity sense. Let $a_i \in A_i$ be such that $f_i(O, a_i) > 0$. Setting $\alpha(t) \equiv a_i$ then $(y_{x,\alpha}, \alpha) \in \mathcal{T}_x$ for all $x \in \Gamma_i$. Moreover, for all $x \in \Gamma_i \setminus \{O\}$, $y_{x,\alpha}(t) \in \Gamma_i \setminus \{O\}$ (the trajectory cannot approach O since the speed pushes it away from O for $y_{x,\alpha} \in \Gamma_i \cap B(O, r)$). Note that it is not sufficient to choose $a_i \in A_i$ such that $f(O, a_i) = 0$ since it can lead to $f(x, a_i) < 0$ for all $x \in \Gamma_i \setminus \{O\}$. Next, for $\tau > 0$ fixed and any $x \in \Gamma_i$, if we choose

$$\alpha_x(t) = \begin{cases} \alpha(t) = a_i & 0 \leq t \leq \tau, \\ \widehat{a}(t - \tau) & t \geq \tau, \end{cases} \quad (4.1)$$

then $y_{x,\alpha_x}(t) \in \Gamma_i \setminus \{O\}$ for all $t \in [0, \tau]$. It yields

$$v_i(x) \leq J(x, \alpha_x) = \int_0^\tau \ell_i(y_{x,\alpha}(s), a_i) e^{-\lambda s} ds + e^{-\lambda \tau} J(y_{x,\alpha}(\tau), \widehat{a}).$$

Since this holds for any \widehat{a} (α_x is arbitrary for $t > \tau$), we deduce that

$$v_i(x) \leq \int_0^\tau \ell_i(y_{x,\alpha_x}(s), a_i) e^{-\lambda s} ds + e^{-\lambda \tau} v_i(y_{x,\alpha_x}(\tau)). \quad (4.2)$$

Since $f_i(\cdot, a)$ is Lipschitz continuous by [H1], we also have for all $t \in [0, \tau]$,

$$\begin{aligned} |y_{x,\alpha_x}(t) - y_{O,\alpha_O}(t)| &= \left| x + \int_0^t f_i(y_{x,\alpha}(s), a_i) e_i ds - \int_0^t f_i(y_{O,\alpha}(s), a_i) e_i ds \right| \\ &\leq |x| + L \int_0^t |y_{x,\alpha}(s) - y_{O,\alpha}(s)| ds, \end{aligned}$$

where α_0 satisfies (4.1) with $x = O$. According to Grönwall's inequality,

$$|y_{x,\alpha_x}(t) - y_{O,\alpha_O}(t)| \leq |x| e^{Lt},$$

for $t \in [0, \tau]$, yielding that $y_{x,\alpha_x}(t)$ tends to $y_{O,\alpha_O}(t)$ when x tends to O . Hence, from (4.2), by letting $x \rightarrow O$, we obtain

$$v_i(O) \leq \int_0^\tau \ell_i(y_{O,\alpha_O}(s), a_i) e^{-\lambda s} ds + e^{-\lambda \tau} v_i(y_{O,\alpha_O}(\tau)).$$

Let φ be a function in $C^1(\Gamma_i)$ such that $0 = v_i(O) - \varphi(O) = \max_{\Gamma_i}(v_i - \varphi)$. This yields

$$\frac{\varphi(O) - \varphi(y_{O,\alpha_O}(\tau))}{\tau} \leq \frac{1}{\tau} \int_0^\tau \ell_i(y_{O,\alpha_O}(s), a_i) e^{-\lambda s} ds + \frac{(e^{-\lambda \tau} - 1) v_i(y_{O,\alpha_O}(\tau))}{\tau}.$$

By letting τ tend to 0, we obtain that

$$-f_i(O, a_i) \frac{d\varphi}{dx_i}(O) \leq \ell_i(O, a_i) - \lambda v_i(O).$$

Hence,

$$\lambda v_i(O) + \sup_{a \in A_i: f_i(O, a) > 0} \left\{ -f_i(O, a) \frac{dv_i}{dx_i}(O) - \ell_i(O, a) \right\} \leq 0$$

in the viscosity sense. Finally, from Corollary A.2 in Appendix A, we have

$$\sup_{a \in A_i: f_i(O, a) > 0} \left\{ -f_i(O, a) \frac{d\varphi_i}{dx_i}(O) - \ell_i(O, a) \right\} = \max_{a \in A_i: f_i(O, a) \geq 0} \left\{ -f_i(O, a) \frac{d\varphi_i}{dx_i}(O) - \ell_i(O, a) \right\}.$$

The proof is complete. \square

Lemma 4.3. *If*

$$v_i(O) < \min \left\{ \min_{j \neq i} \{v_j(O) + c_j\}, -\frac{H_O^T}{\lambda} \right\}, \quad (4.3)$$

then there exist $\bar{\tau} > 0, r > 0$ and $\varepsilon_0 > 0$ such that for any $x \in (\Gamma_i \setminus \{O\}) \cap B(O, r)$, any $\varepsilon < \varepsilon_0$ and any ε -optimal control law $\alpha_{\varepsilon, x}$ for x ,

$$y_{x,\alpha_{\varepsilon,x}}(s) \in \Gamma_i \setminus \{O\}, \quad \text{for all } s \in [0, \bar{\tau}].$$

Remark 4.4. Roughly speaking, this lemma takes care of the case $\lambda v_i + H_i^+ \left(x, \frac{dv_i}{dx_i}(O) \right) \leq 0$, i.e., the situation when the trajectory does not leave Γ_i , see introduction.

Proof of Lemma 4.3. Suppose by contradiction that there exist sequences $\{\varepsilon_n\}, \{\tau_n\} \subset \mathbb{R}^+$ and $\{x_n\} \subset \Gamma_i \setminus \{O\}$ such that $\varepsilon_n \searrow 0, x_n \rightarrow O, \tau_n \searrow 0$ and a control law α_n such that α_n is ε_n -optimal control law and $y_{x_n, \alpha_n}(\tau_n) = O$. This implies that

$$v_i(x_n) + \varepsilon_n > J(x_n, \alpha_n) = \int_0^{\tau_n} \ell(y_{x_n, \alpha_n}(s), \alpha_n(s)) e^{-\lambda s} ds + e^{-\lambda \tau_n} J(O, \alpha_n(\cdot + \tau_n)). \quad (4.4)$$

Since ℓ is bounded by M by [H1], then $v_i(x_n) + \varepsilon_n \geq -\tau_n M + e^{-\lambda\tau_n} v(O)$. By letting n tend to ∞ , we obtain

$$v_i(O) \geq v(O). \quad (4.5)$$

From (4.3), it follows that

$$\min \left\{ \min_{j \neq i} \{v_j(O) + c_j\}, -\frac{H_O^T}{\lambda} \right\} > v(O).$$

However, $v(O) = \min \left\{ \min_j \{v_j(O) + c_j\}, -\frac{H_O^T}{\lambda} \right\}$ by Theorem 2.9. Therefore, $v(O) = v_i(O) + c_i > v_i(O)$, which is a contradiction with (4.5). \square

Lemma 4.5. *The function v_i is a viscosity super-solution of (3.1) at O .*

Proof of Lemma 4.5. We adapt the proof of Oudet [21] and start by assuming that

$$v_i(O) < \min \left\{ \min_{j \neq i} \{v_j(O) + c_j\}, -\frac{H_O^T}{\lambda} \right\}.$$

We need to prove that

$$\lambda v_i(O) + H_i^+ \left(O, \frac{dv_i}{dx_i}(O) \right) \geq 0$$

in the viscosity sense. Let $\varphi \in C^1(\Gamma_i)$ be such that

$$0 = v_i(O) - \varphi(O) \leq v_i(x) - \varphi(x) \quad \text{for all } x \in \Gamma_i, \quad (4.6)$$

and $\{x_\varepsilon\} \subset \Gamma_i \setminus \{O\}$ be any sequence such that x_ε tends to O when ε tends to 0. From the dynamic programming principle and Lemma 4.3, there exists $\bar{\tau}$ such that for any $\varepsilon > 0$, there exists $(y_\varepsilon, \alpha_\varepsilon) := (y_{x_\varepsilon, \alpha_\varepsilon}, \alpha_\varepsilon) \in \mathcal{T}_{x_\varepsilon}$ such that $y_\varepsilon(\tau) \in \Gamma_i \setminus \{O\}$ for any $\tau \in [0, \bar{\tau}]$ and

$$v_i(x_\varepsilon) + \varepsilon \geq \int_0^\tau \ell_i(y_\varepsilon(s), \alpha_\varepsilon(s)) e^{-\lambda s} ds + e^{-\lambda\tau} v_i(y_\varepsilon(\tau)).$$

Then, according to (4.6)

$$\begin{aligned} v_i(x_\varepsilon) - v_i(O) + \varepsilon &\geq \int_0^\tau \ell_i(y_\varepsilon(s), \alpha_\varepsilon(s)) e^{-\lambda s} ds + e^{-\lambda\tau} [\varphi(y_\varepsilon(\tau)) - \varphi(O)] \\ &\quad - v_i(O) (1 - e^{-\lambda\tau}). \end{aligned} \quad (4.7)$$

Next,

$$\begin{cases} \int_0^\tau \ell_i(y_\varepsilon(s), \alpha_\varepsilon(s)) e^{-\lambda s} ds = \int_0^\tau \ell_i(y_\varepsilon(s), \alpha_\varepsilon(s)) ds + o(\tau), \\ [\varphi(y_\varepsilon(\tau)) - \varphi(O)] e^{-\lambda\tau} = \varphi(y_\varepsilon(\tau)) - \varphi(O) + \tau o_\varepsilon(1) + o(\tau), \end{cases}$$

and

$$\begin{cases} v_i(x_\varepsilon) - v_i(O) &= o_\varepsilon(1), \\ v_i(O)(1 - e^{-\lambda\tau}) &= o(\tau) + \tau\lambda v_i(O), \end{cases}$$

where the notation $o_\varepsilon(1)$ is used for a quantity which is independent on τ and tends to 0 as ε tends to 0. For $k \in \mathbb{N}^*$ the notation $o(\tau^k)$ is used for a quantity that is independent on ε and such that $\frac{o(\tau^k)}{\tau^k} \rightarrow 0$ as $\tau \rightarrow 0$.

Finally, $\mathcal{O}(\tau^k)$ stands for a quantity independent on ε such that $\frac{\mathcal{O}(\tau^k)}{\tau^k}$ remains bounded as $\tau \rightarrow 0$. From (4.7), we obtain that

$$\tau\lambda v_i(O) \geq \int_0^\tau \ell_i(y_\varepsilon(s), \alpha_\varepsilon(s)) ds + \varphi(y_\varepsilon(\tau)) - \varphi(O) + \tau o_\varepsilon(1) + o(\tau) + o_\varepsilon(1). \quad (4.8)$$

Since $y_\varepsilon(\tau) \in \Gamma_i$ for all ε , one has

$$\varphi(y_\varepsilon(\tau)) - \varphi(x_\varepsilon) = \int_0^\tau \frac{d\varphi}{dx_i}(y_\varepsilon(s)) \dot{y}_\varepsilon(s) ds = \int_0^\tau \frac{d\varphi}{dx_i}(y_\varepsilon(s)) f_i(y_\varepsilon(s), \alpha_\varepsilon(s)) ds.$$

Hence, from (4.8)

$$\tau\lambda v_i(O) - \int_0^\tau \left[\ell_i(y_\varepsilon(s), \alpha_\varepsilon(s)) + \frac{d\varphi}{dx_i}(y_\varepsilon(s)) f_i(y_\varepsilon(s), \alpha_\varepsilon(s)) \right] ds \geq \tau o_\varepsilon(1) + o(\tau) + o_\varepsilon(1). \quad (4.9)$$

Moreover, $\varphi(x_\varepsilon) - \varphi(O) = o_\varepsilon(1)$ and that $\frac{d\varphi}{dx_i}(y_\varepsilon(s)) = \frac{d\varphi}{dx_i}(O) + o_\varepsilon(1) + \mathcal{O}(s)$. Thus

$$\lambda v_i(O) - \frac{1}{\tau} \int_0^\tau \left[\ell_i(y_\varepsilon(s), \alpha_\varepsilon(s)) + \frac{d\varphi}{dx_i}(O) f_i(y_\varepsilon(s), \alpha_\varepsilon(s)) \right] ds \geq o_\varepsilon(1) + \frac{o(\tau)}{\tau} + \frac{o_\varepsilon(1)}{\tau}. \quad (4.10)$$

Let $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$ and $\tau_m \rightarrow 0$ as $m \rightarrow \infty$ such that

$$(a_{mn}, b_{mn}) := \left(\frac{1}{\tau_m} \int_0^{\tau_m} f_i(y_{\varepsilon_n}(s), \alpha_{\varepsilon_n}(s)) e_i ds, \frac{1}{\tau_m} \int_0^{\tau_m} \ell_i(y_{\varepsilon_n}(s), \alpha_{\varepsilon_n}(s)) ds \right) \longrightarrow (a, b) \in \mathbb{R}e_i \times \mathbb{R}$$

as $n, m \rightarrow \infty$. By [H1] and [H2]

$$\begin{cases} f_i(y_{\varepsilon_n}(s), \alpha_{\varepsilon_n}(s)) e_i &= f_i(O, \alpha_{\varepsilon_n}(s)) + L|y_{\varepsilon_n}(s)| = f_i(O, \alpha_{\varepsilon_n}(s)) e_i + o_n(1) + o_m(1), \\ \ell_i(y_{\varepsilon_n}(s), \alpha_{\varepsilon_n}(s)) e_i &= \ell_i(O, \alpha_{\varepsilon_n}(s)) + \omega(|y_{\varepsilon_n}(s)|) = \ell_i(O, \alpha_{\varepsilon_n}(s)) e_i + o_n(1) + o_m(1). \end{cases}$$

It follows that

$$\begin{aligned} (a_{mn}, b_{mn}) &= \left(\frac{1}{\tau_m} \int_0^{\tau_m} f_i(O, \alpha_{\varepsilon_n}(s)) e_i ds, \frac{1}{\tau_m} \int_0^{\tau_m} \ell_i(O, \alpha_{\varepsilon_n}(s)) ds \right) + o_n(1) + o_m(1) \\ &\in \text{FL}_i(O) + o_n(1) + o_m(1), \end{aligned}$$

since $\text{FL}_i(O)$ is closed and convex. Sending $n, m \rightarrow \infty$, we obtain $(a, b) \in \text{FL}_i(O)$ so there exists $\bar{a} \in A_i$ such that

$$\lim_{m, n \rightarrow \infty} \left(\frac{1}{\tau_m} \int_0^{\tau_m} f_i(y_{\varepsilon_n}(s), \alpha_{\varepsilon_n}(s)) e_i ds, \frac{1}{\tau_m} \int_0^{\tau_m} \ell_i(y_{\varepsilon_n}(s), \alpha_{\varepsilon_n}(s)) ds \right) = (f_i(O, \bar{a}) e_i, \ell_i(O, \bar{a})). \quad (4.11)$$

On the other hand, from Lemma 4.3, $y_{\varepsilon_n}(s) \in \Gamma_i \setminus \{O\}$ for all $s \in [0, \tau_m]$. This yields

$$y_{\varepsilon_n}(\tau_m) = \left[\int_0^{\tau_m} f_i(y_{\varepsilon_n}(s), \alpha_{\varepsilon_n}(s)) ds \right] e_i + x_{\varepsilon_n}.$$

Since $|y_{\varepsilon_n}(\tau_m)| > 0$, then

$$\frac{1}{\tau_m} \int_0^{\tau_m} f_i(y_{\varepsilon_n}(s), \alpha_{\varepsilon_n}(s)) ds \geq -\frac{|x_{\varepsilon_n}|}{\tau_m}.$$

Let ε_n tend to 0, then let τ_m tend to 0, one gets $f_i(O, \bar{a}) \geq 0$, so $\bar{a} \in A_i^+$. Hence, from (4.10) and (4.11), replacing ε by ε_n and τ by τ_m , let ε_n tend to 0, then let τ_m tend to 0, we finally obtain

$$\lambda v_i(O) + \max_{a \in A_i^+} \left\{ -f_i(O, a) \frac{d\varphi}{dx_i}(O) - \ell_i(O, a) \right\} \geq \lambda v_i(O) + \left[-f_i(O, \bar{a}) \frac{d\varphi}{dx_i}(O) - \ell_i(O, \bar{a}) \right] \geq 0.$$

□

5. COMPARISON PRINCIPLE AND UNIQUENESS

Inspired by [6, 7], we begin by proving some properties of sub and super viscosity solutions of (3.1). The following three lemmas are reminiscent of Lemma 3.4, Theorem 3.1 and Lemma 3.5 in [3].

Lemma 5.1. *Let $w = (w_1, \dots, w_N)$ be a viscosity super-solution of (3.1). Let $x \in \Gamma_i \setminus \{O\}$ and assume that*

$$w_i(O) < \min \left\{ \min_{j \neq i} \{w_j(O) + c_j\}, -\frac{H_O^T}{\lambda} \right\}. \quad (5.1)$$

Then for all $t > 0$,

$$w_i(x) \geq \inf_{\alpha_i(\cdot), \theta_i} \left(\int_0^{t \wedge \theta_i} \ell_i(y_x^i(s), \alpha_i(s)) e^{-\lambda s} ds + w_i(y_x^i(t \wedge \theta_i)) e^{-\lambda(t \wedge \theta_i)} \right),$$

where $\alpha_i \in L^\infty(0, \infty; A_i)$, y_x^i is the solution of $y_x^i(t) = x + \left[\int_0^t f_i(y_x^i(s), \alpha_i(s)) ds \right] e_i$ and θ_i satisfies $y_x^i(\theta_i) = 0$ and θ_i lies in $[\tau_i, \bar{\tau}_i]$, where τ_i is the exit time of y_x^i from $\Gamma_i \setminus \{O\}$ and $\bar{\tau}_i$ is the exit time of y_x^i from Γ_i .

Proof of Lemma 5.1. According to (5.1), the function w_i is a viscosity super-solution of the following problem in Γ_i

$$\begin{cases} \lambda w_i(x) + H_i \left(x, \frac{dw_i}{dx_i}(x) \right) = 0 & \text{if } x \in \Gamma_i \setminus \{O\}, \\ \lambda w_i(O) + H_i^+ \left(O, \frac{dw_i}{dx_i}(O) \right) = 0 & \text{if } x = O. \end{cases} \quad (5.2)$$

Hence, we can apply the result in [3, Lemma 3.4]. We refer to [6] for a detailed proof. The main point of that proof uses the results of Blanc [8, 9] on minimal super-solutions of exit time control problems. \square

Lemma 5.2 (Super-optimality). *Under assumption [H], let $w = (w_1, \dots, w_N)$ be a viscosity super-solution of (3.1) that satisfies (5.1); then there exists a sequence $\{\eta_k\}_{k \in \mathbb{N}}$ of strictly positive real numbers such that $\lim_{k \rightarrow \infty} \eta_k = \eta > 0$ and a sequence $x_k \in \Gamma_i \setminus \{O\}$ such that $\lim_{k \rightarrow \infty} x_k = O$, $\lim_{k \rightarrow \infty} w_i(x_k) = w_i(O)$ and for each k , there exists a control law α_i^k such that the corresponding trajectory $y_{x_k}(s) \in \Gamma_i$ for all $s \in [0, \eta_k]$ and*

$$w_i(x_k) \geq \int_0^{\eta_k} \ell_i(y_{x_k}(s), \alpha_i^k(s)) e^{-\lambda s} ds + w_i(y_{x_k}(\eta_k)) e^{-\lambda \eta_k}.$$

Proof of Lemma 5.2. According to (5.1) $\widehat{w}_i(O) < -\frac{H_O^T}{\lambda}$. Hence, this proof is complete by applying the proof of Theorem 3.1 in [3]. \square

Lemma 5.3. *Under assumption [H], let $u = (u_1, \dots, u_N)$ be a viscosity sub-solution of (3.1). Then u_i is Lipschitz continuous in $B(O, r) \cap \Gamma_i$. Therefore, there exists a test function $\varphi_i \in C^1(\Gamma_i)$ which touches u_i from above at O .*

Proof of Lemma 5.3. Since u is a viscosity sub-solution of (3.1), u_i is a viscosity sub-solution of (5.2). Recall that $H_i(x, \cdot)$ is coercive for any $x \in \Gamma_i \cap B(O, r)$, we can apply the proof in Lemma 3.2 of [3], which is based on arguments due to Ishii and contained in [18]. \square

Lemma 5.4 (Sub-optimality). *Under assumption [H], let $u = (u_1, \dots, u_N)$ be a viscosity sub-solution of (3.1). Consider $i = \overline{1, N}$, $x \in \Gamma_i \setminus \{O\}$ and $\alpha_i \in L^\infty(0, \infty; A_i)$. Let $T > 0$ be such that $y_x(t) = x + \left[\int_0^t f_i(y_x(s), \alpha_i(s)) ds \right] e_i$ belongs to Γ_i for any $t \in [0, T]$, then*

$$u_i(x) \leq \int_0^T \ell_i(y_x(s), \alpha_i(s)) e^{-\lambda s} ds + u_i(y_x(T)) e^{-\lambda T}.$$

Proof of Lemma 5.4. Since u is a viscosity sub-solution of (3.1), u_i is a viscosity sub-solution of (5.2) and satisfies $u_i(O) \leq -\frac{H_O^T}{\lambda}$. Hence, we can apply the proof in Lemma 3.5 of [3]. \square

Remark 5.5. Under assumption [H], Lemmas 5.1–5.4 hold for viscosity sub- and super-solution \widehat{u} and \widehat{w} respectively, of the exit cost control problem if (5.1) replaced by

$$\widehat{w}_i(O) < \min \left\{ \min_{j \neq i} \{\widehat{w}_j(O)\} + d_i, -\frac{H_O^T}{\lambda} + d_i \right\}.$$

Theorem 5.6 (Comparison Principle). *Under assumption [H], let u be a bounded viscosity sub-solution of (3.1) and w be a bounded viscosity super-solution of (3.1); then $u \leq w$ in \mathcal{G} , componentwise. This theorem also holds for viscosity sub- and super-solution \widehat{u} and \widehat{w} , respectively, of the exit cost control problem (3.2).*

We give two proofs of Theorem 5.6. The first one is inspired by [3] and uses the previously stated lemmas. The second one uses the elegant arguments proposed in [19].

Proof of Theorem 5.6 inspired by [3]. We focus on u and w , the arguments used for the comparison of \widehat{u} and \widehat{w} are totally similar. Suppose by contradiction that there exists $x \in \Gamma_i$ such that $u_i(x) - w_i(x) > 0$. By classical comparison arguments for the boundary value problem, see [5], $\sup_{\partial \Gamma_i} \{u_i - w_i\}^+ \geq \sup_{\Gamma_i} \{u_i - w_i\}^+$, so we have

$$u_i(O) - w_i(O) = \max_{x \in \Gamma_i} \{u_i(x) - w_i(x)\} > 0.$$

By definition of viscosity sub-solution

$$\lambda u_i(O) + H_O^T \leq 0. \quad (5.3)$$

This implies $\lambda w_i(O) + H_O^T < 0$. We now consider the two following cases.

Case 1: If $w_i(O) < \min_{j \neq i} \{w_j(O) + c_j\}$, from Lemma 5.2 (using the same notations),

$$w_i(x_k) \geq \int_0^{\eta_k} \ell_i(y_{x_k}(s), \alpha_i^k(s)) e^{-\lambda s} ds + w_i(y_{x_k}(\eta_k)) e^{-\lambda \eta_k}.$$

Moreover, according to Lemma 5.4, we also have

$$u_i(x_k) \leq \int_0^{\eta_k} \ell_i(y_{x_k}(s), \alpha_i^k(s)) e^{-\lambda s} ds + u_i(y_{x_k}(\eta_k)) e^{-\lambda \eta_k}.$$

This yields

$$u_i(x_k) - w_i(x_k) \leq [u_i(y_{x_k}(\eta_k)) - w_i(y_{x_k}(\eta_k))] e^{-\lambda \eta_k} \leq [u_i(O) - w_i(O)] e^{-\lambda \eta_k}.$$

By letting k tend to ∞ , one gets

$$u_i(O) - w_i(O) \leq [u_i(O) - w_i(O)] e^{-\lambda \eta}.$$

This implies that $u_i(O) - w_i(O) \leq 0$ and leads to a contradiction.

Case 2: If $w_i(O) \geq \min_{j \neq i} \{w_j(O) + c_j\}$, then there exists $j_0 \neq i$ such that

$$w_{j_0}(O) + c_{j_0} = \min_{j=1, N} \{w_j(O) + c_j\} = \min_{j \neq i} \{w_j(O) + c_j\} \leq w_i(O),$$

because $c_i > 0$. Since c_{j_0} is positive

$$w_{j_0}(O) < \min_{j \neq j_0} \{w_j(O) + c_j\}. \quad (5.4)$$

Next, by Lemma 5.3, there exists a test function φ_i in $C^1(J_i)$ that touches u_i from above at O , it yields

$$\lambda u_i(O) - \lambda \min_{j \neq i} \{u_j(O) + c_j\} \leq \lambda u_i(O) + \max \left\{ -\lambda \min_{j \neq i} \{u_j(O) + c_j\}, H_i^+ \left(O, \frac{d\varphi_i}{dx_i}(O) \right), H_0^T \right\} \leq 0.$$

Therefore

$$w_{j_0}(O) + c_{j_0} \leq w_i(O) < u_i(O) \leq \min_{j \neq i} \{u_j(O) + c_j\} \leq u_{j_0}(O) + c_{j_0}.$$

Thus

$$w_{j_0}(O) < u_{j_0}(O). \quad (5.5)$$

Replacing index i by j_0 in (5.3), we get

$$\lambda w_{j_0}(O) + H_O^T < 0. \quad (5.6)$$

By (5.4) and (5.6), (5.1) holds true. Repeating the proof of Case 1 with j_0 , we reach a contradiction with (5.5). It ends the proof. \square

The comparison principle can also be obtained alternatively, using the arguments which were very recently proposed by Lions and Souganidis in [19]. This new proof is self-combined and the arguments do not rely at all on optimal control theory, but are deeply connected to the ideas used by Soner [23, 24] and Capuzzo-Dolcetta and Lions [11] for proving comparison principles for state-constrained Hamilton-Jacobi equations

Proof of Theorem 5.6 inspired by [19]. We start as in first proof. We argue by contradiction without loss of generality, assuming that there exists i such that

$$u_i(O) - w_i(O) = \max_{\Gamma_i} \{u_i(x) - w_i(x)\} > 0.$$

Therefore $w_i(O) < -\frac{H_O^T}{\lambda}$. We now consider the two following cases.

Case 1: If $w_i(O) < \min_{j \neq i} \{w_j(O) + c_j\}$, then w_i is a viscosity super-solution of (5.2). Recall that by Lemma 5.3, there exists a positive number L such that for $i = \overline{1, N}$, u_i is Lipschitz continuous with Lipschitz constant L in $\Gamma_i \cap B(0, r)$. We consider the function

$$\begin{aligned} \Psi_{i,\varepsilon} : \Gamma_i \times \Gamma_i &\longrightarrow \mathbb{R} \\ (x, y) &\longrightarrow u_i(x) - w_i(y) - \frac{1}{2\varepsilon} [-|x| + |y| + \delta(\varepsilon)]^2 - \gamma(|x| + |y|), \end{aligned}$$

where $\delta(\varepsilon) = (L+1)\varepsilon$ and $\gamma \in \left(0, \frac{1}{2}\right)$. It is clear that $\Psi_{i,\varepsilon}$ attains its maximum $M_{\varepsilon,\gamma}$ at $(x_{\varepsilon,\gamma}, y_{\varepsilon,\gamma}) \in \Gamma_i \times \Gamma_i$. By classical techniques, we check that $x_{\varepsilon,\gamma}, y_{\varepsilon,\gamma} \rightarrow O$ and that $\frac{(x_{\varepsilon,\gamma} - y_{\varepsilon,\gamma})^2}{\varepsilon} \rightarrow 0$ as $\varepsilon \rightarrow 0$. Indeed, one has

$$\begin{aligned} &u_i(x_{\varepsilon,\gamma}) - w_i(y_{\varepsilon,\gamma}) - \frac{[-|x_{\varepsilon,\gamma}| + |y_{\varepsilon,\gamma}| + \delta(\varepsilon)]^2}{2\varepsilon} - \gamma(|x_{\varepsilon,\gamma}| + |y_{\varepsilon,\gamma}|) \\ &\geq \max_{\Gamma_i} \{u_i(x) - w_i(x) - 2\gamma|x|\} - \frac{\delta^2(\varepsilon)}{2\varepsilon} \end{aligned} \quad (5.7)$$

$$\geq u_i(O) - w_i(O) - \frac{(L+1)^2}{2}\varepsilon. \quad (5.8)$$

Since $u_i(O) - w_i(O) > 0$, the term in (5.8) is positive when ε is small enough. We also deduce from the above inequality and from the boundedness of u_i and w_i that, maybe after the extraction of a subsequence, $x_{\varepsilon,\gamma}, y_{\varepsilon,\gamma} \rightarrow x_\gamma$ as $\varepsilon \rightarrow 0$, for some $x_\gamma \in \Gamma_i$. From (5.7),

$$u_i(x_{\varepsilon,\gamma}) - w_i(y_{\varepsilon,\gamma}) - \frac{(|x_{\varepsilon,\gamma}| - |y_{\varepsilon,\gamma}|)^2}{2\varepsilon} - \frac{(-|x_{\varepsilon,\gamma}| + |y_{\varepsilon,\gamma}|)\delta(\varepsilon)}{\varepsilon} \geq \max_{\Gamma_i} \{u_i(x) - w_i(x) - 2\gamma|x|\}.$$

Taking the lim sup on both sides of this inequality when $\varepsilon \rightarrow 0$,

$$\begin{aligned}
u_i(x_\gamma) - w_i(x_\gamma) - 2\gamma|x_\gamma| &\geq \max_{\Gamma_i} \{u_i(x) - w_i(x) - 2\gamma|x|\} + \limsup_{\varepsilon \rightarrow 0} \frac{(|x_{\varepsilon,\gamma}| - |y_{\varepsilon,\gamma}|)^2}{2\varepsilon} \\
&\geq u_i(O) - w_i(O) + \limsup_{\varepsilon \rightarrow 0} \frac{(|x_{\varepsilon,\gamma}| - |y_{\varepsilon,\gamma}|)^2}{2\varepsilon} \\
&\geq u_i(O) - w_i(O) + \liminf_{\varepsilon \rightarrow 0} \frac{(|x_{\varepsilon,\gamma}| - |y_{\varepsilon,\gamma}|)^2}{2\varepsilon} \\
&\geq u_i(O) - w_i(O).
\end{aligned}$$

Recalling that $u_i(O) - w_i(O) = \max_{\Gamma_i} (u_i - w_i)$, we obtain from the inequalities above that $x_\gamma = O$ and that

$$\lim_{\varepsilon \rightarrow 0} \frac{(|x_{\varepsilon,\gamma}| - |y_{\varepsilon,\gamma}|)^2}{2\varepsilon} = 0. \quad (5.9)$$

We claim that if $\varepsilon > 0$, then $x_{\varepsilon,\gamma} \neq O$. Indeed, assume by contradiction that $x_{\varepsilon,\gamma} = O$:

1. if $y_{\varepsilon,\gamma} > 0$, then

$$M_{\varepsilon,\gamma} = u_i(O) - w_i(y_{\varepsilon,\gamma}) - \frac{1}{2\varepsilon} [|y_{\varepsilon,\gamma}| + \delta(\varepsilon)]^2 - \gamma|y_{\varepsilon,\gamma}| \geq u_i(y_{\varepsilon,\gamma}) - w_i(y_{\varepsilon,\gamma}) - \frac{\delta^2(\varepsilon)}{2\varepsilon} - 2\gamma|y_{\varepsilon,\gamma}|.$$

Since u_i is Lipschitz continuous in $B(O, r) \cap \Gamma_i$, we see that for ε small enough

$$L|y_{\varepsilon,\gamma}| \geq u_i(O) - u_i(y_{\varepsilon,\gamma}) \geq \frac{|y_{\varepsilon,\gamma}|^2}{2\varepsilon} + \frac{|y_{\varepsilon,\gamma}|\delta(\varepsilon)}{\varepsilon} - \gamma|y_{\varepsilon,\gamma}| \geq \frac{|y_{\varepsilon,\gamma}|\delta(\varepsilon)}{\varepsilon} - \gamma|y_{\varepsilon,\gamma}|.$$

Therefore, if $y_{\varepsilon,\gamma} \neq O$, then $L \geq L + 1 - \gamma$ which gives a contradiction since $\gamma \in \left(0, \frac{1}{2}\right)$.

2. Otherwise, if $y_{\varepsilon,\gamma} = O$, then

$$M_{\varepsilon,\gamma} = u_i(O) - w_i(O) - \frac{\delta^2(\varepsilon)}{2\varepsilon} \geq u_i(\varepsilon e_i) - w_i(O) - \frac{1}{2\varepsilon} [-\varepsilon + \delta(\varepsilon)]^2 - \gamma\varepsilon.$$

Since u_i is Lipschitz continuous in $B(O, r) \cap \Gamma_i$, we see that for ε small enough,

$$L\varepsilon \geq u_i(O) - u_i(\varepsilon e_i) \geq \frac{|y_{\varepsilon,\gamma}|^2}{2\varepsilon} + \frac{|y_{\varepsilon,\gamma}|\delta(\varepsilon)}{\varepsilon} - 2\gamma|y_{\varepsilon,\gamma}| \geq \frac{|y_{\varepsilon,\gamma}|\delta(\varepsilon)}{\varepsilon} - 2\gamma|y_{\varepsilon,\gamma}|.$$

This implies that $L \geq -\frac{1}{2} + L + 1 - \gamma$, which gives a contradiction since $\gamma \in \left(0, \frac{1}{2}\right)$.

Therefore the claim is proved. It follows that we can apply the viscosity inequality for u_i at $x_{\varepsilon,\gamma}$. Moreover, notice that the viscosity super-solution inequality (5.2) holds also for $y_{\varepsilon,\gamma} = 0$ since $H_i(O, p) \leq H_i^+(O, p)$ for any p . Therefore

$$\begin{aligned}
u_i(x_{\varepsilon,\gamma}) + H_i\left(x_{\varepsilon,\gamma}, \frac{-x_{\varepsilon,\gamma} + y_{\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} + \gamma\right) &\leq 0, \\
w_i(y_{\varepsilon,\gamma}) + H_i\left(y_{\varepsilon,\gamma}, \frac{-x_{\varepsilon,\gamma} + y_{\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} - \gamma\right) &\geq 0.
\end{aligned}$$

Subtracting the two inequalities,

$$u_i(x_{\varepsilon,\gamma}) - w_i(y_{\varepsilon,\gamma}) \leq H_i\left(y_{\varepsilon,\gamma}, \frac{-x_{\varepsilon,\gamma} + y_{\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} + \gamma\right) - H_i\left(x_{\varepsilon,\gamma}, \frac{-x_{\varepsilon,\gamma} + y_{\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} - \gamma\right). \quad (5.10)$$

Using [H1] and [H2], it is easy to see that there exists $\bar{M}_i > 0$ such that for any $x, y \in \Gamma_i, p, q \in \mathbb{R}$

$$\begin{aligned} |H_i(x, p) - H_i(y, q)| &\leq |H_i(x, p) - H_i(y, p)| + |H_i(y, p) - H_i(y, q)| \\ &\leq \bar{M}_i |x - y| (1 + |p|) + \bar{M}_i |p - q|. \end{aligned}$$

It yields

$$\begin{aligned} u_i(x_{\varepsilon,\gamma}) - w_i(y_{\varepsilon,\gamma}) &\leq \bar{M}_i \left[|x_{\varepsilon,\gamma} - y_{\varepsilon,\gamma}| \left(1 + \left| \frac{-x_{\varepsilon,\gamma} + y_{\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} - \gamma \right| \right) + 2|\gamma| \right] \\ &\leq \bar{M}_i \left[|x_{\varepsilon,\gamma} - y_{\varepsilon,\gamma}| \left(\gamma + 1 + \frac{\delta(\varepsilon)}{\varepsilon} \right) + \frac{|x_{\varepsilon,\gamma} - y_{\varepsilon,\gamma}|^2}{\varepsilon} + 2|\gamma| \right]. \end{aligned}$$

Applying (5.9), let ε tend to 0 and γ tend to 0, we obtain that $u_i(O) - w_i(O) \leq 0$, the desired contradiction.

Case 2: $w_i(O) \geq \min_{j \neq i} \{w_j(O) + c_j\} = w_{j_0}(O) + c_{j_0}$. Using the same arguments as in Case 2 of the first proof, we get

$$w_{j_0} < \min \left\{ \min_{j \neq j_0} \{w_j(O) + c_j\}, -\frac{H_O^T}{\lambda} \right\}$$

and $w_{j_0}(O) < u_{j_0}(O)$. Repeating Case 1, replacing the index i by j_0 , implies that $w_{j_0}(O) \geq u_{j_0}(O)$, the desired contradiction. □

Corollary 5.7 (Uniqueness). *If \mathbf{v} is the value function (with entry costs) and (v_1, \dots, v_N) is defined by*

$$v_i(x) = \begin{cases} \mathbf{v}(x) & \text{if } x \in \Gamma_i \setminus \{O\}, \\ \lim_{\delta \rightarrow 0^+} \mathbf{v}(\delta e_i) & \text{if } x = O, \end{cases}$$

then (v_1, \dots, v_N) is the unique bounded viscosity solution of (3.1).

Similarly, if $\hat{\mathbf{v}}$ is the value function (with exit costs) and $(\hat{v}_1, \dots, \hat{v}_N)$ is defined by

$$\hat{v}_i(x) = \begin{cases} \hat{\mathbf{v}}(x) & \text{if } x \in \Gamma_i \setminus \{O\}, \\ \lim_{\delta \rightarrow 0^+} \hat{\mathbf{v}}(\delta e_i) & \text{if } x = O, \end{cases}$$

then $(\hat{v}_1, \dots, \hat{v}_N)$ is the unique bounded viscosity solution of (3.2).

Remark 5.8. From Corollary 5.7, we see that in order to characterize the original value function with entry costs, we need to solve first the Hamilton-Jacobi system (3.1) and find the unique viscosity solution (v_1, \dots, v_N) . The original value function \mathbf{v} with entry costs satisfies

$$\mathbf{v}(x) = \begin{cases} v_i(x) & \text{if } x \in \Gamma_i \setminus \{O\}, \\ \min \left\{ \min_{i=1, \dots, N} \{v_i(O) + c_i\}, -\frac{H_O^T}{\lambda} \right\}, & \text{if } x = O. \end{cases}$$

The characterization of $v(O)$ follows from Theorem 2.9. The characterization of the original value function with exit costs \widehat{v} is similar.

6. A MORE GENERAL OPTIMAL CONTROL PROBLEM

In what follows, we generalize the control problem studied in the previous sections by allowing some of the entry (or exit) costs to be zero. The situation can be viewed as intermediary between the one studied in [3] when all the entry (or exit) costs were zero, and that studied above when all the entry or exit costs were positive. Accordingly, every result presented below will mainly be obtained by combining the arguments proposed above with those used in [3]. Hence, we will present the results and omit the proofs.

To be more specific, we consider the optimal control problems with non-negative entry cost $\overline{C} = \{\overline{c}_1, \dots, \overline{c}_m, \overline{c}_{m+1}, \dots, \overline{c}_N\}$ where $\overline{c}_i = 0$ if $i \leq m$ and $\overline{c}_i > 0$ if $i > m$, keeping all the assumptions and definitions of Section 2 unchanged. The value function associated to \overline{C} will be denoted by V . Similarly to Lemma 2.8, $V|_{\Gamma_i \setminus \{O\}}$ is continuous and Lipschitz continuous near O : therefore, it is possible to extend $V|_{\Gamma_i \setminus \{O\}}$ at O . This extension will be noted \mathcal{V}_i . Moreover, one can check that $\mathcal{V}_i(O) = \mathcal{V}_j(O)$ for all $i, j \leq m$, which means that $V|_{\cup_{i=1}^m \Gamma_i}$ is a continuous function which will be noted \mathcal{V}_c hereafter.

Combining the arguments in [3] and in Section 2 leads us to the following theorem.

Theorem 6.1. *The value function V satisfies*

$$\max_{i=m+1, N} \{\mathcal{V}_i(O)\} \leq V(O) = \mathcal{V}_c(O) \leq \min \left\{ \min_{i=m+1, N} \{\mathcal{V}_i(O) + \overline{c}_i\}, -\frac{H_O^T}{\lambda} \right\}.$$

Remark 6.2. In the case when $\overline{c}_i = 0$ for $i = \overline{1, N}$, V is continuous on \mathcal{G} and it is exactly the value function of the problem studied in [3].

We now define a set of admissible test-function and the Hamilton-Jacobi equation that will characterize V .

Definition 6.3. A function $\varphi : (\cup_{i=1}^m \Gamma_i) \times \Gamma_{m+1} \times \dots \times \Gamma_N \rightarrow \mathbb{R}^{N-m+1}$ of the form $\varphi(x_c, x_{m+1}, \dots, x_N) = (\varphi_c(x_c), \varphi_{m+1}(x_{m+1}), \dots, \varphi_N(x_N))$ is an admissible test-function if

- φ_c is continuous and for $i \leq m$, $\varphi_c|_{\Gamma_i}$ belongs to $C^1(\Gamma_i)$,
- for $i > m$, φ_i belongs to $C^1(\Gamma_i)$,
- the space of admissible test-function is noted $R(\mathcal{G})$.

Definition 6.4. A function $U = (U_c, U_{m+1}, \dots, U_N)$ where $U_c \in USC(\cup_{j=1}^m \Gamma_j; \mathbb{R})$, $U_i \in USC(\Gamma_i; \mathbb{R})$ is called a *viscosity sub-solution* of the Hamilton-Jacobi system if for any $(\varphi_c, \varphi_{m+1}, \dots, \varphi_N) \in R(\mathcal{G})$:

1. if $U_c - \varphi_c$ has a local maximum at $x_c \in \cup_{j=1}^m \Gamma_j$ and if
 - $x_c \in \Gamma_j \setminus \{O\}$ for some $j \leq m$, then

$$\lambda U_c(x_c) + H_j \left(x, \frac{d\varphi_c}{dx_j}(x_c) \right) \leq 0,$$

- $x_c = O$, then

$$\lambda U_c(O) + \max \left\{ -\lambda \min_{j>m} \{U_j(O) + \overline{c}_j\}, \max_{j \leq m} \left\{ H_j^+ \left(O, \frac{d\varphi_c}{dx_j^+}(O) \right) \right\}, H_O^T \right\} \leq 0;$$

2. if $U_i - \varphi_i$ has a local maximum point at $x_i \in \Gamma_i$ for $i > m$, and if

- $x_i \in \Gamma_i \setminus \{O\}$, then

$$\lambda U_i(x_i) + H_i\left(x, \frac{d\varphi_i}{dx_i}(x_i)\right) \leq 0,$$

- $x_i = O$, then

$$\lambda U_i(O) + \max\left\{-\lambda \min_{j>m, j \neq i} \{U_j(O) + \bar{c}_j\}, -\lambda U_c(O), H_i^+\left(O, \frac{d\varphi_i}{dx_i}(O)\right), H_O^T\right\} \leq 0.$$

A function $U = (U_c, U_{m+1}, \dots, U_N)$ where $U_c \in LSC(\cup_{j=1}^m \Gamma_j; \mathbb{R})$, $U_i \in LSC(\Gamma_i; \mathbb{R})$ is called a *viscosity super-solution* of the Hamilton-Jacobi system if

$$U_c(O) \geq U_i(O), \quad \text{for } i = \overline{m+1, N}, \quad (6.1)$$

and for any $(\varphi_c, \varphi_{m+1}, \dots, \varphi_N) \in R(\mathcal{G})$:

1. if $U_c - \varphi_c$ has a local maximum at $x_c \in \cup_{j=1}^m \Gamma_j$ and if
 - $x_c \in \Gamma_j \setminus \{O\}$ for some $j \leq m$, then

$$\lambda U_c(x_c) + H_j\left(x, \frac{d\varphi_c}{dx_j}(x_c)\right) \geq 0,$$

- $x_c = O$, then

$$\lambda U_c(O) + \max\left\{-\lambda \min_{j>m} \{U_j(O) + \bar{c}_j\}, \max_{j \leq m} \left\{H_j^+\left(O, \frac{d\varphi_c}{dx_j^+}(O)\right)\right\}, H_O^T\right\} \geq 0;$$

2. if $U_i - \varphi_i$ has a *local minimum point* at $x_i \in \Gamma_i$ for $i > m$, and if
 - $x_i \in \Gamma_i \setminus \{O\}$, then

$$\lambda U_i(x_i) + H_i\left(x, \frac{d\varphi_i}{dx_i}(x_i)\right) \geq 0,$$

- $x_i = O$ for $i > m$ then

$$\lambda U_i(O) + \max\left\{-\lambda \min_{j>m, j \neq i} \{U_j(O) + \bar{c}_j\}, -\lambda U_c(O), H_i^+\left(O, \frac{d\varphi_i}{dx_i}(O)\right), H_O^T\right\} \geq 0.$$

A function $U = (U_c, U_1, \dots, U_m)$ where $U_c \in C(\cup_{j \leq m} \Gamma_j; \mathbb{R})$ and $U_i \in C(\Gamma_i; \mathbb{R})$ for all $i > m$ is called a *viscosity solution* of the Hamilton-Jacobi system if it is both a viscosity sub-solution and a viscosity super-solution of the Hamilton-Jacobi system.

Remark 6.5. The term $-\lambda H_C(O)$ in the above definition accounts for the situation in which the trajectory enters $\cup_{j=1}^m \Gamma_j$. The term $\max_{j \leq m} \left\{H_j^+\left(O, \frac{d\varphi_c}{dx_j^+}(O)\right)\right\}$ accounts for the situation in which the trajectory enters Γ_{i_0} where $H_{i_0}^+\left(O, \frac{d\varphi_c}{dx_j^+}(O)\right) = \max_{j \leq m} \left\{H_j^+\left(O, \frac{d\varphi_c}{dx_j^+}(O)\right)\right\}$.

Remark 6.6. In the case when $\bar{c}_i = 0$ for $i = \overline{1, N}$, i.e., $m = N$, the term $-\lambda \min_{j>m} U_j(O) + \bar{c}_j$ vanishes. This implies that

$$\begin{aligned} \max \left\{ -\lambda \min_{j>m} \{U_j(O) + \bar{c}_j\}, \max_{j \leq m} \left\{ H_j^+ \left(O, \frac{\partial \varphi_c}{\partial e_j^+}(O) \right) \right\}, H_O^T \right\} &= \max_{j=\overline{1, N}} \left\{ H_j^+ \left(O, \frac{\partial \varphi_c}{\partial e_j^+}(O) \right) \right\} \\ &= H_O \left(\frac{\partial \varphi_c}{\partial e_1^+}(O), \dots, \frac{\partial \varphi_c}{\partial e_N^+}(O) \right). \end{aligned}$$

where $H_O(p_1, \dots, p_N)$ is defined in page 6 of [3]. This means that, in the case when all the entry costs \bar{c}_j vanish, we recover the notion of viscosity solution proposed in [3].

We now study the relationship between the value function V and the Hamilton-Jacobi system.

Theorem 6.7. *Let V be the value function corresponding to the entry costs \bar{C} , then $(\mathcal{V}_c, \mathcal{V}_{m+1}, \dots, \mathcal{V}_N)$ is a viscosity solution of the Hamilton-Jacobi system.*

Let us state the comparison principle for the Hamilton-Jacobi system.

Theorem 6.8. *Let $U = (U_c, U_{m+1}, \dots, U_N)$ and $W = (W_c, W_{m+1}, \dots, W_N)$ be a bounded viscosity sub-solution and a viscosity super-solution, respectively, of the Hamilton-Jacobi system. The following holds: $U \leq W$ in \mathcal{G} , i.e., $U_c \leq W_c$ on $\cup_{j=1}^m \Gamma_j$, and $U_i \leq W_i$ in Γ_i for all $i > m$.*

Proof of Theorem 6.8. Suppose by contradiction that there exists $i \in \{1, \dots, N\}$ and $x \in \Gamma_i$ such that

$$\begin{cases} U_c(x) - W_c(x) > 0 & \text{if } i \leq m, \\ U_i(x) - W_i(x) > 0 & \text{if } i > m, \end{cases}$$

then

$$\begin{cases} U_c(O) - W_c(O) = \max_{\cup_{j=1}^m \Gamma_j} \{U_c - W_c\} > 0 & \text{if } i \leq m, \\ U_i(O) - W_i(O) = \max_{\Gamma_i} \{U_i - W_i\} > 0 & \text{if } i > m, \end{cases}$$

since the case where the positive maximum is achieved outside the junction leads to a contradiction by classical comparison results.

Case 1: $U_c(O) - W_c(O) = \max_{\cup_{i=1}^m \Gamma_i} (U_c - W_c) > 0$

Sub-case 1-a: $W_c(O) < \min_{j>m} \{W_j(O) + \bar{c}_j\}$. Since $W_c(O) < U_c(O) \leq -\frac{H_O^T}{\lambda}$, the function W_c is a viscosity super-solution of

$$\begin{cases} \lambda W_c(x) + H_i \left(x, \frac{dW_c}{dx_i}(x) \right) = 0 & \text{if } i \leq m, x \in \Gamma_i \setminus \{O\}, \\ \lambda W_c(O) + H_c \left(\frac{dW_c}{dx_1^+}(O), \dots, \frac{dW_c}{dx_m^+}(O) \right) = 0 & \text{if } x = O. \end{cases}$$

where $H_c(p_1, \dots, p_m) = \max_{i \leq m} H_i^+(O, p_i)$. Applying Lemma A.3 in Appendix A, we obtain that $U_c(O) \leq W_c(O)$ in contradiction with the assumption.

Sub-case 1-b: $W_c(O) \geq \min_{j>m} \{W_j(O) + \bar{c}_j\} = W_{i_0}(O) + \bar{c}_{i_0}$. Since $\bar{c}_{i_0} > 0$, we first see that $W_{i_0}(O) < \min \left\{ \min_{j>m} \{W_j(O) + \bar{c}_j\}, W_c(O), -\frac{H_O^T}{\lambda} \right\}$. Hence, W_{i_0} is a viscosity super-solution of (5.2) replacing

i by i_0 . Moreover, since

$$U_{i_0}(O) + \bar{c}_{i_0} \geq \min_{j>m} (U_j(O) + \bar{c}_j) \geq U_c(O) > W_c(O) > W_{i_0}(O) + \bar{c}_{i_0},$$

then $U_{i_0}(O) > W_{i_0}(O)$. Applying the same argument as Case 1 in the second proof of Theorem 5.6 replacing i by i_0 , we obtain that $U_{i_0}(O) \leq W_{i_0}(O)$, which is contradictory.

Case 2: $U_i(O) - W_i(O) = \max_{\Gamma_i} (U_i - W_i) > 0$ for some $i > m$. Using the definition of viscosity sub-solutions and Case 1, we see that $W_i(O) < U_i(O) \leq U_c(O) \leq W_c(O)$.

Sub-case 2-a: $W_i(O) < \min_{j>m} \{W_j(O) + \bar{c}_j\}$. Since $U_i(O) < -\frac{H_O^T}{\lambda}$, we first see that $W_i(O) < \min \left\{ \min_{j>m} \{W_j(O) + \bar{c}_j\}, W_c(O), -\frac{H_O^T}{\lambda} \right\}$. Hence, W_i is a viscosity super-solution of (5.2). Applying the same argument as in Case 1 in the second proof of Theorem 5.6, we see that $U_i(O) \leq W_i(O)$, which is contradictory.

Sub-case 2-b: $W_i(O) \geq \min_{j>m} \{W_j(O) + \bar{c}_j\} = W_{i_0}(O) + \bar{c}_{i_0}$. Since $\bar{c}_{i_0} > 0$, we can check that $W_{i_0}(O) < \min \left\{ \min_{j>m} \{W_j(O) + \bar{c}_j\}, W_c(O), -\frac{H_O^T}{\lambda} \right\}$. Hence, W_{i_0} is a viscosity super-solution of (5.2) replacing i by i_0 . Moreover, since

$$U_{i_0}(O) + \bar{c}_{i_0} \geq \min_{j>m} (U_j(O) + \bar{c}_j) \geq U_c(O) > W_i(O) > W_{i_0}(O) + \bar{c}_{i_0},$$

then $U_{i_0}(O) > W_{i_0}(O)$. Applying the same argument as Case 1 in the second proof of Theorem 5.6 replacing i by i_0 , we obtain that $U_{i_0}(O) \leq W_{i_0}(O)$ which is contradictory. □

APPENDIX A

Lemma A.1. *For any $a \in A_i^+$, there exists a sequence $\{a_n\}$ such that $a_n \in A_i$ and*

$$\begin{aligned} f_i(O, a_n) &\geq \frac{\delta}{n} > 0, \\ |f_i(O, a_n) - f_i(O, a)| &\leq \frac{2M}{n}, \\ |\ell_i(O, a_n) - \ell_i(O, a)| &\leq \frac{2M}{n}. \end{aligned}$$

Proof of Lemma A.1. From assumption [H4], there exists $a_\delta \in A_i$ such that $f_i(O, a_\delta) = \delta$. Since $\text{FL}_i(O)$ is convex (by assumption [H3]), for any $n \in \mathbb{N}, a \in A_i^+$

$$\frac{1}{n} (f_i(O, a_\delta) e_i, \ell_i(O, a_\delta)) + \left(1 - \frac{1}{n}\right) (f_i(O, a), \ell_i(O, a) e_i) \in \text{FL}_i(O).$$

Then, there exists a sequence $\{a_n\}$ such that $a_n \in A_i$ and

$$\frac{1}{n} (f_i(O, a_\delta), \ell_i(O, a_\delta)) + \left(1 - \frac{1}{n}\right) (f_i(O, a), \ell_i(O, a)) = (f_i(O, a_n), \ell_i(O, a_n)) \in \text{FL}_i(O). \quad (\text{A.1})$$

Notice that $f_i(O, a) \geq 0$ since $a \in A_i^+$, this yields

$$f_i(O, a_n) \geq \frac{f_i(O, a_\delta)}{n} = \frac{\delta}{n} > 0.$$

From (A.1), we also have

$$|f_i(O, a_n) - f_i(O, a)| = \frac{1}{n} |f_i(O, a_\delta) - f_i(O, a)| \leq \frac{2M}{n},$$

and

$$|\ell_i(O, a_n) - \ell_i(O, a)| = \frac{1}{n} |\ell_i(O, a_\delta) - \ell_i(O, a)| \leq \frac{2M}{n}.$$

□

We can state the following corollary of Lemma A.1:

Corollary A.2. For $i = \overline{1, N}$ and $p_i \in \mathbb{R}$,

$$\max_{a \in A_i \text{ s.t. } f_i(O, a) \geq 0} \{-f_i(O, a) p_i - \ell_i(O, a)\} = \sup_{a \in A_i \text{ s.t. } f_i(O, a) > 0} \{-f_i(O, a) p_i - \ell_i(O, a)\}.$$

Lemma A.3. If U_c and W_c are respectively viscosity sub and super-solution of

$$\begin{aligned} \lambda U_c(x) + H_i\left(x, \frac{dU_c}{dx_i}(x)\right) &\leq 0 \text{ if } x \in \Gamma_i \setminus \{O\}, \\ \lambda U_c(O) + H_c\left(\frac{dU_c}{dx_1}(O), \dots, \frac{dU_c}{dx_m}(O)\right) &\leq 0 \text{ if } x = O, \end{aligned}$$

and

$$\begin{aligned} \lambda W_c(x) + H_i\left(x, \frac{dW_c}{dx_i}(x)\right) &\geq 0 \text{ if } x \in \Gamma_i \setminus \{O\}, \\ \lambda W_c(O) + H_c\left(\frac{dW_c}{dx_1}(O), \dots, \frac{dW_c}{dx_m}(O)\right) &\geq 0 \text{ if } x = O, \end{aligned}$$

then $U_c(x) \leq W_c(x)$ for all $x \in \bigcup_{i=1}^m \Gamma_i$.

Proof of Lemma A.3. Assume that there exists $\hat{x} \in \Gamma_i$ where $1 \leq i \leq m$ and $U_c(\hat{x}) - W_c(\hat{x}) > 0$. By classical comparison principle for the boundary problem on Γ_i , one gets

$$U_c(O) - W_c(O) = \max_{\Gamma_i} \{U_c(x) - W_c(x)\} > 0.$$

Applying again classical comparison principle for the boundary problem for each edge Γ_j

$$U_c(O) - W_c(O) = \max_{\bigcup_{i=1}^m \Gamma_i} \{U_c(x) - W_c(x)\} > 0.$$

For $j = \overline{1, N}$, we consider the function

$$\Psi_{j, \varepsilon, \gamma} : \Gamma_j \times \Gamma_j \longrightarrow \mathbb{R}$$

$$(x, y) \longrightarrow U_c(x) - W_c(y) - \frac{1}{2\varepsilon} [-|x| + |y| + \delta(\varepsilon)]^2 - \gamma(|x| + |y|),$$

where $\delta(\varepsilon) = (L+1)\varepsilon$, $\gamma \in \left(0, \frac{1}{2}\right)$.

The function $\Psi_{j,\varepsilon}$ attains its maximum at $(x_{j,\varepsilon,\gamma}, y_{j,\varepsilon,\gamma}) \in \Gamma_j \times \Gamma_j$. Applying the same argument as in the second proof of Theorem 5.6, we have $x_{j,\varepsilon,\gamma}, y_{j,\varepsilon,\gamma} \rightarrow O$ and $\frac{(x_{j,\varepsilon,\gamma} - y_{j,\varepsilon,\gamma})^2}{\varepsilon_n} \rightarrow 0$ as $\varepsilon \rightarrow 0$. Moreover, for any $j = \overline{1, m}$, $x_{j,\varepsilon,\gamma} \neq O$. We claim that $y_{j,\varepsilon,\gamma}$ must be O for ε small enough. Indeed, if there exists a sequence ε_n such that $y_{j,\varepsilon_n,\gamma} \in \Gamma_j \setminus \{O\}$, then applying viscosity inequalities, we have

$$\begin{aligned} U_c(x_{j,\varepsilon_n,\gamma}) + H_j \left(x_{j,\varepsilon_n,\gamma}, \frac{-x_{j,\varepsilon_n,\gamma} + y_{j,\varepsilon_n,\gamma} + \delta(\varepsilon_n)}{\varepsilon_n} + \gamma \right) &\leq 0, \\ W_c(y_{j,\varepsilon_n,\gamma}) + H_j \left(y_{j,\varepsilon_n,\gamma}, \frac{-x_{j,\varepsilon_n,\gamma} + y_{j,\varepsilon_n,\gamma} + \delta(\varepsilon_n)}{\varepsilon_n} - \gamma \right) &\geq 0. \end{aligned}$$

Subtracting the two inequalities and using (5.10) with H_j , we obtain

$$U_c(x_{j,\varepsilon_n,\gamma}) - W_c(y_{j,\varepsilon_n,\gamma}) \leq \overline{M}_j |x_{j,\varepsilon_n,\gamma} - y_{j,\varepsilon_n,\gamma}| \left(1 + \left| \frac{-x_{j,\varepsilon_n,\gamma} + y_{j,\varepsilon_n,\gamma} + \delta(\varepsilon_n)}{\varepsilon_n} - \gamma \right| \right) + \overline{M}_j 2\gamma.$$

Recall that we already have $\frac{(x_{j,\varepsilon_n,\gamma} - y_{j,\varepsilon_n,\gamma})^2}{\varepsilon_n} \rightarrow 0$ as $n \rightarrow \infty$. Let n tend to ∞ and γ tend to 0 then we obtain $U_c(O) - W_c(O) \leq 0$. It leads us to a contradiction. So this claim is proved.

Define the function $\Psi : \bigcup_{j=1}^m \Gamma_j \rightarrow \mathbb{R}$ by

$$\Psi|_{\Gamma_i}(y) = \frac{1}{2\varepsilon} \sum_{j \neq i} \left\{ [-|x_{i,\varepsilon,\gamma}| + \delta(\varepsilon)]^2 - \gamma |x_{i,\varepsilon,\gamma}| \right\} + \frac{1}{2\varepsilon} [-|x_{i,\varepsilon,\gamma}| + |y| + \delta(\varepsilon)]^2 + \gamma(-|x_{i,\varepsilon,\gamma}| + |y|).$$

We can see that Ψ is continuous on $\bigcup_{j=1}^m \Gamma_j$ and belongs to $C^1(\Gamma_j)$ for $j = \overline{1, m}$. Moreover, for $j = \overline{1, m}$ and for ε small enough, $y_{j,\varepsilon,\gamma} = O$ then the function $\Psi + W_c$ has a minimum point at O . It yields

$$\lambda W_c(O) + H_c \left(\frac{-\overline{x}_{1,\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon}, \dots, \frac{-\overline{x}_{m,\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} \right) \geq 0.$$

By definition of H_c , there exists $j_0 \in \{1, \dots, m\}$ such that

$$\lambda W_c(O) + H_{j_0}^+ \left(O, \frac{-\overline{x}_{j_0,\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} \right) \geq 0.$$

This implies

$$\lambda W_c(O) + H_{j_0} \left(O, \frac{-\overline{x}_{j_0,\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} \right) \geq 0$$

On the other hand, since $x_{j_0,\varepsilon,\gamma} \in \Gamma_{j_0} \setminus \{O\}$, we have

$$\lambda U_c(\overline{x}_{j_0,\varepsilon,\gamma}) + H_{j_0} \left(\overline{x}_{j_0,\varepsilon,\gamma}, \frac{-\overline{x}_{j_0,\varepsilon,\gamma} + \delta(\varepsilon)}{\varepsilon} \right) \leq 0.$$

Subtracting the two inequalities and using properties of Hamiltonian H_{j_0} , let ε tend to 0 then γ tend to 0, we obtain that $U_c(O) - W_c(O) \leq 0$, which is contradictory. \square

Acknowledgements. I would like to express my thanks to my advisors Y. Achdou, O. Ley and N. Tchou for suggesting me this work and for their help. I also thank the hospitality of Centre Henri Lebesgue and INSA de Rennes during the preparation of this work. Moreover, this work was partially supported by the ANR (Agence Nationale de la Recherche) through HJnet project ANR-12-BS01-0008-01 and MFG project ANR-16-CE40-0015-01.

REFERENCES

- [1] Y. Achdou, F. Camilli, A. Cutri and N. Tchou, Hamilton-Jacobi equations on networks. *IFAC Proc.* **44** (2011) 2577–2582.
- [2] Y. Achdou, F. Camilli, A. Cutri and N. Tchou, Hamilton-Jacobi equations constrained on networks. *NoDEA Nonlinear Differ. Equ. Appl.* **20** (2013) 413–445.
- [3] Y. Achdou, S. Oudet and N. Tchou, Hamilton-Jacobi equations for optimal control on junctions and networks. *ESAIM: COCV* **21** (2015) 876–899.
- [4] G. Barles, Discontinuous viscosity solutions of first-order Hamilton-Jacobi equations: a guided visit. *Nonlinear Anal.* **20** (1993) 1123–1134.
- [5] G. Barles, An introduction to the theory of viscosity solutions for first-order Hamilton-Jacobi equations and applications, in *Hamilton-Jacobi Equations: Approximations, Numerical Analysis and Applications*. Vol. 2074 of *Lecture Notes in Mathematics*. Springer, Heidelberg (2013) 49–109.
- [6] G. Barles, A. Briani and E. Chasseigne, A Bellman approach for two-domains optimal control problems in \mathbb{R}^N . *ESAIM: COCV* **19** (2013) 710–739.
- [7] G. Barles, A. Briani and E. Chasseigne, A Bellman approach for regional optimal control problems in \mathbb{R}^N . *SIAM J. Control Optim.* **52** (2014) 1712–1744.
- [8] A.-P. Blanc, Deterministic exit time control problems with discontinuous exit costs. *SIAM J. Control Optim.* **35** (1997) 399–434.
- [9] A.-P. Blanc, Comparison principle for the Cauchy problem for Hamilton-Jacobi equations with discontinuous data. *Nonlinear Anal.* **45** (2001) 1015–1037.
- [10] F. Camilli and C. Marchi, A comparison among various notions of viscosity solution for Hamilton-Jacobi equations on networks. *J. Math. Anal. Appl.* **407** (2013) 112–118.
- [11] I. Capuzzo-Dolcetta and P.-L. Lions, Hamilton-Jacobi equations with state constraints. *Trans. Am. Math. Soc.* **318** (1990) 643–683.
- [12] K.-J. Engel, M. Kramar Fijavz, R. Nagel and E. Sikolya, Vertex control of flows in networks. *Netw. Heterog. Media* **3** (2008) 709–722.
- [13] H. Frankowska and M. Mazzola, Discontinuous solutions of Hamilton-Jacobi-Bellman equation under state constraints. *Calc. Var. Partial Differ. Equ.* **46** (2013) 725–747.
- [14] M. Garavello and B. Piccoli, Conservation laws models, in *Traffic Flow on Networks*. Vol. 1 of *AIMS Series on Applied Mathematics*. American Institute of Mathematical Sciences (AIMS), Springfield, MO (2006).
- [15] P. J. Graber, C. Hermosilla and H. Zidani, Discontinuous solutions of Hamilton-Jacobi equations on networks. *J. Differ. Equ.* **263** (2017) 8418–8466.
- [16] C. Imbert and R. Monneau, Flux-limited solutions for quasi-convex Hamilton-Jacobi equations on networks. *Ann. Sci. Éc. Norm. Supér.* **50** (2017) 357–448.
- [17] C. Imbert, R. Monneau and H. Zidani, A Hamilton-Jacobi approach to junction problems and application to traffic flows. *ESAIM: COCV* **19** (2013) 129–166.
- [18] H. Ishii, A short introduction to viscosity solutions and the large time behavior of solutions of Hamilton-Jacobi equations, in *Hamilton-Jacobi Equations: Approximations, Numerical Analysis and Applications*. Vol. 2074 of *Lecture Notes in Mathematics*. Springer, Heidelberg (2013) 111–249.
- [19] P.-L. Lions and P. Souganidis, Viscosity solutions for junctions: well posedness and stability. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **27** (2016) 535–545.
- [20] P.-L. Lions and P. Souganidis, Well Posedness for Multi-dimensional Junction Problems With Kirchoff-Type Conditions. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **28** (2017) 807–816.
- [21] S. Oudet, Hamilton-Jacobi Equations for Optimal Control on Multidimensional Junctions. Preprint [arXiv:1412.2679](https://arxiv.org/abs/1412.2679) (2014).
- [22] D. Schieborn and F. Camilli, Viscosity solutions of Eikonal equations on topological networks. *Calc. Var. Partial Differ. Equ.* **46** (2013) 671–686.
- [23] H.M. Soner, Optimal control with state-space constraint. I. *SIAM J. Control Optim.* **24** (1986) 552–561.
- [24] H.M. Soner, Optimal control with state-space constraint. II. *SIAM J. Control Optim.* **24** (1986) 1110–1122.