

ON THE SHAPE OF SMALL LIQUID DROPS MINIMIZING NONLOCAL ENERGIES

KONSTANTINOS BESSAS¹, MATTEO NOVAGA² AND FUMIHIKO ONOUE^{3,*} 

Abstract. We study the equilibrium shape of liquid drops minimizing the fractional perimeter under the action of a potential energy. We prove, with a quantitative estimate, that the small volume minimizers are convex and uniformly close to a ball.

Mathematics Subject Classification. 49Q20, 53A10, 35R09, 35R11.

Received April 13, 2023. Accepted November 8, 2023.

1. INTRODUCTION

Given $s \in (0, 1)$ and $m > 0$, we study the following minimization problem:

$$\min \left\{ \mathcal{E}_{s,g}(E) := P_s(E) + \int_E g(x) \, dx : |E| = m \right\}, \quad (1.1)$$

where $P_s(E)$ is the s -fractional perimeter of a measurable set $E \subset \mathbb{R}^N$ defined as

$$P_s(E) := \int_E \int_{E^c} \frac{1}{|x - y|^{N+s}} \, dx \, dy, \quad (1.2)$$

and $\int_E g(x) \, dx$ is the potential energy of E associated with a locally Lipschitz and coercive function $g : \mathbb{R}^N \mapsto \mathbb{R}$. The notion of s -fractional perimeter was introduced by Caffarelli, Roquejoffre, and Savin in [5], who were motivated by the classical phase field model when long-range correlations exist. After their celebrated work, variational problems involving the s -fractional perimeter have been studied by many authors as an analogy of the classical ones and our paper is in this line of research.

Our research is motivated by the classical minimization problem for the equilibrium shape of liquid drops and crystals. The classical problem can be formulated as follows:

$$\min \left\{ \mathcal{E}(E) := \int_{\partial^* E} f(\nu_E) \, d\mathcal{H}^{N-1} + \int_E g(x) \, dx : |E| = m \right\}, \quad (1.3)$$

Keywords and phrases: Fractional perimeter, isoperimetric problem, regularity of minimizers, liquid drop model.

¹ Dipartimento di Matematica, Università di Pavia, Via Adolfo Ferrata 5, 27100 Pavia, Italy.

² Dipartimento di Matematica, Università di Pisa, Largo Bruno Pontecorvo 5, 56127 Pisa, Italy.

³ Zentrum Mathematik, Technische Universität München, Boltzmannstrasse 3, 85748 Garching, Germany.

* Corresponding author: fumihiko.onoue@tum.de

where $f : \mathbb{R}^N \rightarrow [0, \infty)$ is a convex, positively 1-homogeneous function, ν_E is the measure-theoretic unit normal to E , and ∂^*E is the reduced boundary (see the definition of ν_E and ∂^*E in [20]). Notice that, if f is the Euclidean norm, then the first term in \mathcal{E} is exactly the classical perimeter of a set $E \subset \mathbb{R}^N$ in the sense of De Giorgi.

Almgren originally proposed the following question on the shape of minimizers for Problem (1.3). This question is mentioned in [21].

Question If the potential g is convex, then is it true that the minimizer for Problem (1.3) is convex (at least connected) and unique?

The geometric properties of minimizers for liquid drop and crystal models associated with Problem (1.3) have been studied for decades by many authors, for instance, Almgren, Avron, Baer, De Philippis, Figalli, Finn, Goldman, Gonzalez, Maggi, Massari, McCann, Tamanini, Taylor, Zia, and so on [2, 3, 9, 12–15]. Partial answers to Question are given by some of these authors.

In particular, Figalli and Maggi in [12] extensively studied the shape of minimizers for Problem (1.3) in the situation where the contribution coming from the potential term $E \mapsto \int_E g(x) dx$ can be negligible compared with the perimeter term associated with f . They proved that the minimizers of \mathcal{E} are uniformly close to a Wulff shape when the volume is sufficiently small. In addition, if $N = 2$, they showed that the minimizers are convex and, if $N \geq 3$, assuming a stronger regularity on the 1-homogeneous function f and the potential g of the energy \mathcal{E} , they also proved the convexity of minimizers with a quantitative estimate for the second fundamental form on the boundaries of minimizers. We remark that De Philippis and Goldman in [9] proved that, if the function f is smooth and uniformly elliptic, and the potential g is convex and coercive, then any minimizer for Problem (1.3) without volume constraint is convex in any dimension. Moreover, under the same assumptions on f and g as above, they showed that, in dimension 2, any minimizer for Problem (1.3) (with volume constraint in this case) is convex and unique. On the other hand, Indrei in [16] proved the non-existence of minimizers for Problem (1.3) in dimension 2 if the potential g is merely convex and satisfies $g(0) = 0$. Moreover, Indrei and Karakhanyan in [17] recently showed the sharp quantitative stability for Problem (1.3) if the function $f = 1$ on \mathbb{S}^{N-1} and the potential g is radial and non-decreasing and $g(0) = 0$, which implies that the unique minimizer for Problem (1.3) is a ball.

To the best of our knowledge, the shape of minimizers for the nonlocal version of Problem (1.3), that is, Problem (1.1) is not well-understood and there are few references on the problem. For instance, Cesaroni and Novaga in [8] proved the existence and regularity of minimizers for Problem (1.1) in the case that the potential function is either coercive or \mathbb{Z}^N -periodic. Our goal in this paper is to study the shape of minimizers for Problem (1.1) and to establish a nonlocal version of some of the results shown in [12]. Precisely, we prove

Theorem 1.1. *Let $s \in (0, 1)$. Assume that the potential g satisfies (H1) and (H2) (see Sect. 2 for the assumptions) and let $c = c(g) := \min_{\mathbb{R}^N} g$. Then, there exist constants $m_0 = m_0(N, s, g) > 0$ and $R_0 = R_0(N, g) > 0$ such that, for any $m \in (0, m_0)$, every minimizer E_m of $\mathcal{E}_{s,g}$ with $|E_m| = m$ satisfies the following: there exists a point $x_m \in \mathbb{R}^N$ such that $|x_m| \leq R_0$, $\lim_{m \rightarrow 0} \text{dist}(x_m, \{g = c\}) = 0$ and*

$$x_m + B_{\sigma(m)(1-r_0)} \subset E_m \subset x_m + B_{\sigma(m)(1+r_0)},$$

for some $r_0 \in (0, Cm^{s^2/(2N^2)})$ and some constant $C > 0$ depending only on N , s , and g , where we set

$$\sigma(m) := \left(\frac{m}{|B_1|} \right)^{\frac{1}{N}}. \quad (1.4)$$

Moreover, we have that ∂E_m is of class $C^{2,\alpha}$ with $\alpha < s$ and ∂E_m converges to ∂B_1 in C^2 as $m \downarrow 0$, by proper scaling and translation. In particular, E_m is convex for small $m \in (0, m_0)$.

From a geometric point of view, the existence of minimizers for Problem (1.3) is related to the existence of a smooth hypersurface \mathcal{M} on which the following prescribed mean curvature equation holds:

$$H_{\mathcal{M}} + g = 0 \quad \text{on } \mathcal{M}, \quad (1.5)$$

where $H_{\mathcal{M}}$ is the mean curvature on \mathcal{M} . Indeed, if $E_m \subset \mathbb{R}^N$ is a smooth minimizer for Problem (1.3) with $|E_m| = m$, then we can obtain the following Euler–Lagrange equation:

$$H_{\partial E_m} + g = \lambda_m \quad \text{on } \partial E_m.$$

As a nonlocal analogy of the classical problem, the existence of minimizers of our energy $\mathcal{E}_{s,g}$ is related to the problem of finding a set E with a smooth boundary such that the following prescribed “fractional” mean curvature equation holds on ∂E :

$$H_{s,\partial E}(x) + g(x) = 0 \quad \text{for } x \in \partial E. \quad (1.6)$$

Here $H_{s,\partial E}(x)$ is the s -fractional mean curvature on ∂E at x , which is defined as

$$H_{s,\partial E}(x) := \text{P.V.} \int_{\mathbb{R}^N} \frac{\chi_{E^c}(y) - \chi_E(y)}{|y - x|^{N+s}} dy \quad (1.7)$$

where “P.V.” means the Cauchy principle value. Indeed, assuming that E_m is a minimizer of $\mathcal{E}_{s,g}$ among sets with volume $m > 0$ and ∂E_m is smooth, we can show that the following Euler–Lagrange equation holds true:

$$H_{s,\partial E_m} + g = \lambda_m \quad \text{on } \partial E_m \quad (1.8)$$

where λ_m is a Lagrange multiplier. As we mentioned above, the authors in [8] proved the existence of bounded minimizers with smooth boundary for Problem (1.1) and this implies the existence of compact hypersurfaces $\mathcal{M} \subset \mathbb{R}^N$ on which the equation (1.8) holds true for any $m > 0$. In this context, our main theorem may imply the existence of a compact, smooth hypersurface $\mathcal{M}_m \subset \mathbb{R}^N$ with non-negative mean curvature such that the equation (1.8) holds on \mathcal{M}_m for small $m > 0$ and \mathcal{M}_m , by properly rescaling, converges to the sphere in \mathbb{R}^N as $m \downarrow 0$.

The proof of Theorem 1.1 is divided in several steps. The former part of Theorem 1.1 (the uniform closeness of minimizers to the Euclidean ball) will be proved in Section 3 and the latter part of Theorem 1.1 (the regularity and convexity of minimizers) will be proved in Section 5.

The organization of our paper is as follows: in Section 2 we fix some notation and we introduce the mathematical setting of our problem.

In Section 3 we first establish properties of minimizers of (2.1) and consider a suitable rescaling for them. Then, we prove the uniform closeness of the rescaled minimizers to the unit Euclidean ball for small volumes in Theorem 3.7. The strategy follows some arguments in [8, 10, 12]. Precisely, we derive suitable integro-differential inequalities for estimating the volume of the difference between rescaled minimizers and balls centred in the origin. A similar argument can be found in [8], Proposition 3.2, where the authors, applying a nonlocal version of the so-called Almgren’s Lemma (see [12], Lem. 3), prove the boundedness of minimizers of (2.1) for a fixed value of the volume. The first main difference with the approach in [8] is that our estimates not only aim at obtaining boundedness of minimizer, but also at closeness in L^∞ to the unit ball (for small volumes). The second one is that our estimates are uniform with respect to the volume of minimizers. To obtain the uniformity with respect to the volume, we adapt ideas in [12] coming from the classical case. In particular, we do not employ a nonlocal Almgren’s Lemma such as [8], Lemma 3.1, but we made use of a suitable dilation. Indeed, with our approach, the constants involved in our estimates depend only on the volume of the minimizers rather than on the minimizers themselves.

In Section 4, with the uniform proximity result in our hands, we can reformulate Problem (2.1) into a variational unconstrained problem, which becomes crucial for establishing regularity.

In Section 5, we obtain the regularity and convexity of minimizers for Problem (1.1), employing several results on the regularity of the so-called “almost minimizers” or “ Λ -minimizers”, where Λ is independent of minimizers, of the s -fractional perimeter and the bootstrap argument for integro-differential equations shown in [4]. As a consequence, by applying the regularity result for the “ Λ -minimizers” shown in [11], Corollary 3.6 and from the uniform closeness of the minimizers to balls that we prove in Section 3, we obtain the convergence of the minimizers to the Euclidean ball in C^2 -topology, which implies in particular the convexity of minimizers.

2. NOTATION AND SETTING OF THE PROBLEM

We denote the Euclidean norm of $x \in \mathbb{R}^N$ by $|x|$, the Euclidean ball of radius $r > 0$ and centre $x \in \mathbb{R}^N$ by $B_r(x)$ and the Euclidean ball of volume $m > 0$ and centre $x \in \mathbb{R}^N$ by $B^m(x)$. We also set $B_r := B_r(0)$ and $B^m := B^m(0)$.

If E is a non-empty subsets of \mathbb{R}^N , then $\text{dist}(x, E) := \inf\{|x - y| : y \in E\}$ for all $x \in \mathbb{R}^N$.

If E, F are two subsets of \mathbb{R}^N we indicate by $E \triangle F$ their *symmetric difference*, i.e. $E \triangle F := (E \setminus F) \cup (F \setminus E)$.

We denote the Lebesgue measure of \mathbb{R}^N by \mathcal{L}^N and the class of all Lebesgue measurable sets of \mathbb{R}^N by \mathcal{M}_N . For $E \in \mathcal{M}_N$ we also set $|E| := \mathcal{L}^N(E)$ and we denote the topological boundary of $E \in \mathcal{M}_N$ by ∂E . \mathcal{H}^{N-1} denotes the $(N - 1)$ -dimensional Hausdorff measure of \mathbb{R}^N .

Let $g : \mathbb{R}^N \rightarrow \mathbb{R}$ be

(H1) coercive, i.e. s.t. $\lim_{|x| \rightarrow +\infty} g(x) = +\infty$;

(H2) locally Lipschitz.

Let E_m be a minimizer of the constrained problem (1.1) with volume m , that is

$$E_m \in \arg \min \left\{ \mathcal{E}_{s,g}(E) := P_s(E) + \int_E g(x) dx : |E| = m \right\}. \quad (2.1)$$

Moreover, we recall the definition of (1.4) for every $m > 0$, so that $B^m = B_{\sigma(m)}$ (the ball of volume m is equal to the ball of radius $\sigma(m)$).

Without loss of generality (thanks to the structure of the volume constrained problem in (2.1)) in the whole work we assume

$$\inf_{\mathbb{R}^N} g = g(0) = 0. \quad (2.2)$$

Indeed, fixed any $\hat{x} \in \mathbb{R}^N$ such that $g(\hat{x}) = \inf_{\mathbb{R}^N} g$, by replacing the potential g with \hat{g} , defined by $\hat{g}(x) := g(x + \hat{x}) - \inf_{\mathbb{R}^N} g$, on the one hand we observe that \hat{g} satisfies (H1), (H2) and (2.1) on the other one we note that the minimizers of $\mathcal{E}_{s,\hat{g}}$ with volume m are the same minimizers of $\mathcal{E}_{s,g}$ with volume m (upon a translation via \hat{x}).

Remark 2.1. We observe that E_m is well-defined, since the problem (1.1) admits solution if the potential g is measurable, bounded from below and satisfies assumption (H1). This can be proved via the Direct Method as done in [8], Proposition 5.3.

As far as uniqueness is concerned, to the best of our knowledge, there are no complete results available in the literature. Even in the classical case (see Problem (1.3)) a general answer has not been provided yet, even though there are some partial results. Precisely, from [9], Remark 1.6 it follows that for large volumes, minimizers of (1.3) are unique when the potential g is convex and coercive.

We also mention that in [21], Theorem 1.1 uniqueness (up to translations) is established in $2D$, for any volume but among convex competitors, when the potential g is convex and the set where it vanishes is bounded but non-empty.

For every $m > 0$, we let

$$\mathcal{S}_m := \{E_m^i\}_{i \in \mathcal{I}_m} \quad (2.3)$$

be the family of all the minimizers of $\mathcal{E}_{s,g}$ with volume constraint $|E_m^i| = m$. We observe that $\mathcal{S}_m \neq \emptyset$ because there always exists a minimizer of $\mathcal{E}_{s,g}$ for every $m > 0$. In the sequel, by a little abuse of notation, we will drop the dependence on $i \in \mathcal{I}_m$ in E_m^i when it is not essential to highlight it (and also in other quantities possibly depending on $i \in \mathcal{I}_m$).

3. UNIFORM PROXIMITY TO THE UNIT BALL

Definition 3.1. Let $E \in \mathcal{M}_N$, i.e a measurable set of \mathbb{R}^N , such that $0 < |E| < +\infty$. We define the *Fraenkel asymmetry* of E as

$$A(E) := \inf \left\{ \frac{|E \triangle B^{|E|}(x)|}{|E|} : x \in \mathbb{R}^N \right\},$$

where $B^{|E|}(x)$ is the ball centred at x with volume $|E|$. Moreover, if $P_s(E) < +\infty$ we define the *Wulff s -deficit* of E as

$$\delta_s(E) := \frac{P_s(E)}{P_s(B^{|E|})} - 1,$$

where $B^{|E|}$ is the ball centred at the origin with volume $|E|$.

Remark 3.2. We observe that A and δ_s are scaling and translation invariant, that is:

$$A(E) = A(\lambda E + x),$$

$$\delta_s(E) = \delta_s(\lambda E + x),$$

for every $\lambda \in \mathbb{R} \setminus \{0\}$, $x \in \mathbb{R}^N$ and $E \in \mathcal{M}_N$ with $0 < |E| < +\infty$. Furthermore, \bar{x} attains the infimum in the definition of $A(E)$ if and only if $x + \bar{x}$ attains the infimum in the definition of $A(\lambda E + x)$.

We state the following quantitative isoperimetric inequality for the fractional perimeter P_s (see also [11]).

Theorem 3.3. Let $N \geq 2$ and $s \in (0, 1)$. Let $E \in \mathcal{M}_N$, i.e a measurable set of \mathbb{R}^N , such that $0 < |E| < +\infty$. Then, there exists $C(N, s) > 0$ s.t.

$$\delta_s(E) \geq C(N, s) A(E)^2.$$

We are now ready to establish properties of minimizers of (1.1).

Proposition 3.4 (Properties of Minimizers). Let $s \in (0, 1)$ and assume that g satisfies (H1), (H2) and (2.2). Let $\mathcal{S}_m := \{E_m^i\}_{i \in \mathcal{I}_m}$ be as in (2.3) and $\sigma(m)$ be as in (1.4) for every $m > 0$. Then, there exists $C(N, s) > 0$ s.t. for every $m > 0$ and $i \in \mathcal{I}_m$

$$\delta_s(E_m^i) \leq C(N, s) \sigma(m)^s \sup_{B_{\sigma(m)}} g, \quad (3.1)$$

$$A(E_m^i) = \frac{|E_m^i \triangle B^m(x_m^i)|}{m} \leq C(N, s) \sqrt{\sigma(m)^s \sup_{B_{\sigma(m)}} g}, \quad (3.2)$$

for some $x_m = x_m^i \in \mathbb{R}^N$.

Moreover, there exist $m_0 = m_0(N, s, g) > 0$ and $R_0 = R_0(N, g) > 0$ such that for every $m \leq m_0$ and $i \in \mathcal{I}_m$

$$|x_m^i| \leq R_0, \quad (3.3)$$

and

$$\lim_{m \rightarrow 0} \sup_{i \in \mathcal{I}_m} \text{dist}(x_m^i, \{g = 0\}) = 0. \quad (3.4)$$

Furthermore, if we let

$$\tilde{E}_m := \frac{(E_m - x_m)}{\sigma(m)} \quad (3.5)$$

and

$$g_m(x) := \sigma(m)^s g(\sigma(m)x + x_m), \quad (3.6)$$

then

$$\tilde{E}_m \in \arg \min \left\{ P_s(F) + \int_F g_m(x) \, dx : |F| = |B_1| \right\} \quad (3.7)$$

for every $m > 0$.

Proof. This proof follows the strategy adopted in part of [12], Theorem 9 combined with [8], Proposition 3.2.

Let $m > 0$ and $i \in \mathcal{I}_m$. By the non-negativity of g (recall (2.2)) and the minimality of E_m^i (see (2.3)),

$$P_s(E_m^i) \leq P_s(E_m^i) + \int_{E_m^i} g \, dx \leq P_s(B^m) + \int_{B^m} g \, dx. \quad (3.8)$$

Then,

$$\delta_s(E_m^i) \leq \frac{\int_{B^m} g(x) \, dx}{P_s(B^m)} = \frac{\sigma(m)^N \int_{B_1} g(\sigma(m)\tilde{x}) \, d\tilde{x}}{\sigma(m)^{N-s} P_s(B_1)} \leq \frac{|B_1|}{P_s(B_1)} \sigma(m)^s \sup_{B_{\sigma(m)}} g,$$

so that (3.1) is proved. Since by [8], Proposition 3.2 E_m^i is bounded, the infimum in the definition of $A(E_m^i)$ (see Def. 3.1) is actually a minimum. This observation joint with Theorem 3.3 and (3.1) implies (3.2) for some $x_m^i \in \mathbb{R}^N$.

Thanks to the local boundedness of g (see (H2)), from (3.2) we can find a constant $m_0(N, s, g) > 0$ such that

$$\frac{|E_m^i \setminus B_{\sigma(m)}(x_m^i)|}{m} \leq \frac{1}{2},$$

for all $m \leq m_0$ and $i \in \mathcal{I}_m$; which (since $|E_m^i| = m$) is equivalent to

$$|E_m^i \cap B_{\sigma(m)}(x_m^i)| \geq \frac{m}{2}, \quad (3.9)$$

for all $m \leq m_0$ and $i \in \mathcal{I}_m$.

Since (H1) holds, we observe that $\{g = 0\}$ is contained in a compact subset of \mathbb{R}^N . Then, (3.4) implies (3.3) for some $R_0 = R_0(N, g) > 0$ (possibly choosing $m_0 = m_0(N, s, g) > 0$ smaller).

We now claim that (3.4) holds.

We define $d_m := \sup_{i \in \mathcal{I}_m} \text{dist}(x_m^i, \{g = 0\})$ for every $m \in (0, m_0]$ and we consider the set $I_{m_0} := \{m \in (0, m_0] : d_m > 2\sigma(m)\}$.

We first observe that the fact that $\sigma(m)$ converges to zero as $m \rightarrow 0$ implies,

$$\lim_{\substack{m \rightarrow 0 \\ m \in (0, m_0] \setminus I_{m_0}}} d_m = 0, \quad (3.10)$$

whenever 0 is an accumulation point for $(0, m_0] \setminus I_{m_0}$.

If $m \in I_{m_0}$, then $B_{\sigma(m)}(x_m^i) \subset \mathbb{R}^N \setminus (\{g = 0\} + B_{d_m - 2\sigma(m)})$ for some $i \in \mathcal{I}_m$ (by definition of d_m and the triangular inequality).

We now observe that by the fractional isoperimetric inequality and (3.8) it immediately follows

$$\int_{E_m^i} g \, dx \leq \int_{B^m} g \, dx. \quad (3.11)$$

Then, by (3.9), (3.11) and the non-negativity of g (see (2.2)):

$$\frac{m}{2} \inf_{\mathbb{R}^N \setminus (\{g=0\} + B_{d_m - 2\sigma(m)})} g \leq \int_{E_m^i \cap B_{\sigma(m)}(x_m^i)} g \, dx \leq \int_{B^m} g \, dx \leq m \sup_{B_{\sigma(m)}} g,$$

for some $i \in \mathcal{I}_m$.

Therefore, since g is continuous and $g(0) = 0$ (see (2.2) and (H2)):

$$\limsup_{\substack{m \rightarrow 0 \\ m \in I_{m_0}}} \inf_{\mathbb{R}^N \setminus (\{g=0\} + B_{d_m - 2\sigma(m)})} g \leq 2 \limsup_{m \rightarrow 0} \sup_{B_{\sigma(m)}} g = 0, \quad (3.12)$$

whenever 0 is an accumulation point for I_{m_0} .

From (3.12) and the coercivity of g (see (H1)) and the non-negativity of g (see (2.2)) we then deduce that

$$\lim_{\substack{m \rightarrow 0 \\ m \in I_{m_0}}} d_m = 0, \quad (3.13)$$

whenever 0 is an accumulation point for I_{m_0} .

Therefore, (3.4) is proved by combining (3.10) and (3.13).

Since

$$\mathcal{E}_{s,g}(\sigma(m)F + x_m) = \sigma(m)^{N-s} \mathcal{E}_{s,g_m}(F)$$

for every $F \in \mathcal{M}_N$ with $|F| = |B_1|$, (3.7) follows. \square

In order to prove the uniform closeness of the rescaled minimizers \tilde{E}_m to the Euclidean ball for small volumes (see Thm. 3.7), we need two technical lemmas.

Lemma 3.5 (Estimate for the Fractional Derivative). *Let $a, b \in \mathbb{R}$ such that $a < b$ and $s \in (0, 1)$.*

(i) Let $f : [a, +\infty) \rightarrow [0, +\infty)$ be absolutely continuous and monotone non-increasing such that $\lim_{r \rightarrow +\infty} f(r) = 0$. Then,

$$-(1-s) \int_a^b dr \int_r^{+\infty} dt \frac{f'(t)}{(t-r)^s} \leq (b-a)^{1-s} f(a) \quad (3.14)$$

if the integral in (3.14) is finite.

(ii) Let $f : [0, b] \rightarrow [0, +\infty)$ be absolutely continuous and monotone non-decreasing such that $f(0) = 0$. Then,

$$(1-s) \int_a^b dr \int_0^r dt \frac{f'(t)}{(r-t)^s} \leq (b-a)^{1-s} f(b) \quad (3.15)$$

if the integral in (3.15) is finite.

Proof. We prove only (i) since the proof of (ii) is analogous.

By Fubini's Theorem:

$$\begin{aligned} & -(1-s) \int_a^b dr \int_r^{+\infty} dt \frac{f'(t)}{(t-r)^s} = \int_a^{+\infty} f'(t) dt \int_a^{t \wedge b} dr \frac{d}{dr} [(t-r)^{1-s}] \\ & = \int_a^{+\infty} f'(t) [(t-t \wedge b)^{1-s} - (t-a)^{1-s}] dt \\ & = \int_a^b (-f'(t))(t-a)^{1-s} dt + \int_b^{+\infty} (-f'(t))[(t-a)^{1-s} - (t-b)^{1-s}] dt \\ & \leq (b-a)^{1-s} \int_a^{+\infty} (-f'(t)) = (b-a)^{1-s} f(a) \end{aligned}$$

We also used that the function $(b, +\infty) \ni t \mapsto (t-a)^{1-s} - (t-b)^{1-s}$ is non-negative and monotone non-increasing, so it can be bounded by $(b-a)^{1-s}$. \square

Lemma 3.6. Let $a, b \in \mathbb{R}$ such that $a < b$, $s \in (0, 1)$ and $c > 0$.

(i) Let $u : [a, b] \rightarrow [0, +\infty)$ be monotone non-increasing such that

$$a + \frac{(2u(a))^{\frac{s}{N}} N}{sc} \leq b,$$

and

$$2c \int_{\varrho}^b u(r)^{\frac{N-s}{N}} dr \leq u(\varrho), \quad (3.16)$$

for every $a < \varrho < b$.

Then,

$$u\left(a + \frac{(2u(a))^{\frac{s}{N}} N}{sc}\right) = 0 \quad (3.17)$$

(ii) Let $w : [a, b] \rightarrow [0, +\infty)$ be monotone non-decreasing such that

$$b - \frac{(2w(b))^{\frac{s}{N}} N}{sc} \geq a,$$

and

$$2c \int_a^\varrho w(r)^{\frac{N-s}{N}} dr \leq w(\varrho),$$

for every $a < \varrho < b$.

Then,

$$w\left(b - \frac{(2w(b))^{\frac{s}{N}} N}{sc}\right) = 0 \tag{3.18}$$

Proof. The proof follows the argument in [10], Lemma 4.1 with slight adjustments.

We prove (i).

We highlight that in this proof the constant c in (3.16) cannot vary from line to line: it has to be considered fixed. Now we argue in a similar fashion as in [10], Lemma 4.1. We define the auxiliary function $h : [a, +\infty) \rightarrow \mathbb{R}$ as

$$h(\varrho) := \begin{cases} \left[(2u(a))^{\frac{s}{N}} - c \frac{s}{N} (\varrho - a) \right]^{\frac{N}{s}} & \text{if } a \leq \varrho \leq a + \frac{(2u(a))^{\frac{s}{N}} N}{sc} \\ 0 & \text{if } \varrho > a + \frac{(2u(a))^{\frac{s}{N}} N}{sc} \end{cases}$$

for every $\varrho \geq a$. We observe that h is continuous, $h(a) = 2u(a)$ and it satisfies

$$c \int_\varrho^b h(r)^{\frac{N-s}{N}} dr = h(\varrho), \tag{3.19}$$

for every $\varrho \geq a$. Our goal is now to prove that

$$u(\varrho) \leq h(\varrho) \quad \forall \varrho : a \leq \varrho \leq a + \frac{(2u(a))^{\frac{s}{N}} N}{sc}. \tag{3.20}$$

We define $I := \{\varrho \in (a, b) : u \geq h \text{ in } [\varrho, b]\}$ and we observe that it is an interval and that

$$a + \frac{(2u(a))^{\frac{s}{N}} N}{sc} \in I.$$

Now, we let

$$R_* := \inf I \in \left[a, a + \frac{(2u(a))^{\frac{s}{N}} N}{sc} \right]. \tag{3.21}$$

Therefore, we can find a sequence $\{R_n\}_{n \in \mathbb{N}} \subseteq [a, R_*]$ converging to R_* , such that $h(R_n) \geq u(R_n)$ for every $n \in \mathbb{N}$. So, combining (3.16) and (3.19), we get

$$\begin{aligned} h(R_n) &\geq u(R_n) \geq 2c \int_{R_n}^b u(r)^{\frac{N-s}{N}} dr \geq 2c \int_{R_n}^{R_*} u(r)^{\frac{N-s}{N}} dr + 2c \int_{R_*}^b h(r)^{\frac{N-s}{N}} dr \\ &= 2c \int_{R_n}^{R_*} u(r)^{\frac{N-s}{N}} dr + 2h(R_*) \geq 2h(R_*). \end{aligned}$$

Letting n go to $+\infty$, we obtain $h(R_*) \geq \lim_{r \rightarrow R_*^-} u(r) \geq 2h(R_*)$, which implies $u(R_*) = h(R_*) = 0$. By definition

of h and thanks to (3.21), we finally observe that $R_* = a + \frac{(2u(a))^{\frac{s}{N}} N}{sc}$. The proof of (i) is complete.

The proof of (ii) is similar to the proof of (i) and thus is left to the reader. \square

Theorem 3.7 (Uniform Closeness of Rescaled Minimizers to B_1). *Let $s \in (0, 1)$ and g s.t. (H1), (H2) and (2.2) hold. For every $m > 0$ let E_m be a minimizer of $\mathcal{E}_{s,g}$ with $|E_m| = m$ and let \tilde{E}_m be as in (3.5). There exist $m_0 = m_0(N, s, g) > 0$ and $C = C(N, s, g) > 0$ such that if $0 < m \leq m_0(N, s, g)$, then \tilde{E}_m satisfies*

$$B_{1-r_0} \subset \tilde{E}_m \subset B_{1+r_0},$$

for some $0 < r_0 \leq C(N, s, g)m^{s^2/(2N^2)}$.

Proof. The proof is in the spirit of the one of [12], Theorem 5.

We choose $m_0 = m_0(N, s, g)$ smaller than the constant m_0 in the statement of Theorem 3.4 so that

$$\sigma(m_0) < 1, \tag{3.22}$$

and we choose $R_0 = R_0(N, g)$ and x_m as in the statement of Theorem 3.4 (so that $|x_m| \leq R_0$ whenever $m \in (0, m_0)$, as in Prop. 3.4).

Then, we fix $0 < m \leq m_0(N, s, g)$.

For convenience, we put $E := \tilde{E}_m$ and we change our focus from the variable m to $\varepsilon := \sqrt{\sigma(m)^s}$. Therefore, applying Proposition 3.4, we deduce that there exists $\varepsilon_0(N, s, g) > 0$ such that $0 < \varepsilon \leq \varepsilon_0(N, s, g)$ and

$$\delta_s(E) \leq C(N, s, g)\varepsilon^2; \tag{3.23}$$

$$|E \triangle B_1| \leq C(N, s, g)\varepsilon.$$

We claim (possibly choosing a smaller $\varepsilon_0(N, s, g)$ or, equivalently, a smaller $m_0(N, s, g)$) that there exists a constant $C(N, s, g)$ such that

$$B_{1-r_0} \subset E \subset B_{1+r_0},$$

for some $r_0 = r_0(\varepsilon, N, s, g) \leq C(N, s, g)\varepsilon^{s/N}$. From (3.23) (in alternative we can simply use the minimality, testing E against B_1) we deduce

$$P_s(E) \leq C(N, s, g). \tag{3.24}$$

STEP 1: proof of the inclusion $E \subset B_{1+r_0}$.

For every $r > 0$ we define the non-negative and monotone non-increasing absolutely continuous function

$$u(r) := |E \setminus B_r|.$$

Thanks to the monotonicity and the hypotheses we observe that for every $r \geq 1$

$$u(r) \leq u(1) \leq C(N, s, g)\varepsilon. \quad (3.25)$$

If the set $\{r > 1 : u(r) > 0\}$ is empty, there is nothing to prove, so we assume it is not. We let

$$r_1 := \sup\{r > 1 : u(r) > 0\} \in (1, +\infty]. \quad (3.26)$$

Our goal is to prove that

$$r_1 - 1 \leq C(N, s, g)\varepsilon^{s/N}. \quad (3.27)$$

We fix $r > 0$ such that $1 < r < \min\{2, r_1\}$. By definition of r_1 we have that $|E \cap B_r| < |E|$.

We consider the following competitor (with volume equal to $|E| = |B_1|$) for the volume constrained problem,

$$F := \lambda(E \cap B_r),$$

with

$$\lambda = \lambda(r) := \left(\frac{|E|}{|E \cap B_r|} \right)^{\frac{1}{N}} > 1.$$

After a computation, thanks to (3.25), we observe that λ is arbitrarily close to 1 (upon choosing $\varepsilon_0(N, s, g)$ small enough). In particular (upon choosing $\varepsilon_0(N, s, g)$ small enough) we get

$$0 < \lambda - 1 \leq C(N)u(r) \quad (3.28)$$

and, since $r < 2$,

$$F \subseteq B_{\lambda r} \subseteq B_3.$$

At this point we make use of the minimality of E (see (3.7)) to write

$$\begin{aligned} & P_s(E) + \int_E g_m(x) \, dx \\ & \leq P_s(F) + \int_F g_m(x) \, dx = \lambda^{N-s} P_s(E \cap B_r) + \lambda^N \int_{E \cap B_r} g_m(\lambda x) \, dx \end{aligned}$$

Now, we recall [10], Lemma 2.1 and making use of the isoperimetric inequality for P_s and the fact that $E = (E \setminus B_r) \sqcup (E \cap B_r)$ we obtain

$$\begin{aligned} C(N, s)(u(r))^{\frac{N-s}{N}} & \leq P_s(E \setminus B_r) = P_s(E) - P_s(E \cap B_r) + 2 \int_{E \setminus B_r} \int_{E \cap B_r} \frac{1}{|x-y|^{N+s}} \, dx dy \\ & \leq (\lambda^{N-s} - 1)P_s(E \cap B_r) \end{aligned} \quad (3.29)$$

$$+ \lambda^N \int_{E \cap B_r} g_m(\lambda x) \, dx - \int_E g_m(x) \, dx \quad (3.30)$$

$$+ 2 \int_{E \setminus B_r} \int_{E \cap B_r} \frac{1}{|x-y|^{N+s}} \, dx dy. \quad (3.31)$$

We first estimate (3.29) using (3.24), (3.25), (3.28) (possibly choosing a smaller $\varepsilon_0(N, s, g)$) and the convexity of balls (see [11], Lem. B.1):

$$(\lambda^{N-s} - 1)P_s(E \cap B_r) \leq C(N, s)(\lambda - 1)P_s(E) \leq C(N, s, g)u(r).$$

We then estimate (3.30) using the fact that g is locally Lipschitz (see (H2)) and non-negative (see (2.2)) together with (3.25), (3.28), (3.22) and the fact that $r < 2$ (possibly choosing a smaller $\varepsilon_0(N, s, g)$):

$$\begin{aligned} & \lambda^N \int_{E \cap B_r} g_m(\lambda x) \, dx - \int_E g_m(x) \, dx \\ &= (\lambda^N - 1) \int_{E \cap B_r} g_m(\lambda x) \, dx + \int_{E \cap B_r} (g_m(\lambda x) - g_m(x)) \, dx - \int_{E \setminus B_r} g_m(x) \, dx \\ &\leq C(N)(\lambda - 1)|E \cap B_r| \sup_{B_{3+R_0}} g + (\lambda - 1)r \operatorname{Lip}(g; B_{3+R_0})|E \cap B_r| \\ &\leq C(N, s, g)u(r). \end{aligned}$$

Now we deal with (3.31). Arguing exactly as in [8], equation (15), we find

$$\int_{E \setminus B_r} \int_{E \cap B_r} \frac{1}{|x-y|^{N+s}} \, dx dy \leq -\frac{N\omega_N}{s} \int_r^{+\infty} \frac{u'(t)}{(t-r)^s} \, dt.$$

Gathering all the previous inequalities together, we find that for every $1 < r < \min\{2, r_1\}$,

$$C(N, s)(u(r))^{\frac{N-s}{N}} \leq C(N, s, g)u(r) - \frac{N\omega_N}{s} \int_r^{+\infty} \frac{u'(t)}{(t-r)^s} \, dt. \quad (3.32)$$

Now, recalling (3.25), upon choosing $\varepsilon_0(N, s, g)$ smaller, we can rewrite (3.32) as

$$C(N, s, g)(u(r))^{\frac{N-s}{N}} \leq -\frac{N\omega_N}{s} \int_r^{+\infty} \frac{u'(t)}{(t-r)^s} \, dt. \quad (3.33)$$

Now we integrate both sides of (3.33) with respect to the variable r and apply Lemma 3.5 part (i) to obtain that for some $c = c(N, s, g) > 0$

$$2c \int_{\varrho}^{\min\{2, r_1\}} u(r)^{\frac{N-s}{N}} \, dr \leq u(\varrho), \quad (3.34)$$

for every $1 < \varrho < \min\{2, r_1\}$.

Recalling (3.25), upon choosing $C(N, s, g)$ larger and $\varepsilon_0(N, s, g)$ smaller, we can assume

$$1 + \frac{(2u(1))^{\frac{s}{N}} N}{sc} \leq 1 + C(N, s, g)\varepsilon^{\frac{s}{N}} < 2. \quad (3.35)$$

Without loss of generality we can also assume that

$$1 + \frac{(2u(1))^{\frac{s}{N}} N}{sc} < r_1, \quad (3.36)$$

otherwise (3.35) immediately implies (3.27).

(3.34), (3.35), (3.36) allow us to invoke Lemma 3.6 part (i) to deduce that

$$u\left(1 + \frac{(2u(1))^{\frac{s}{N}} N}{sc}\right) = 0. \quad (3.37)$$

From (3.37), recalling (3.35) and (3.26), we obtain that $r_1 \leq 1 + \frac{(2u(1))^{\frac{s}{N}} N}{sc} \leq 1 + C(N, s, g)\varepsilon^{\frac{s}{N}}$.

STEP 2: proof of the inclusion $B_{1-r_0} \subset E$.

Thanks to the previous step, upon choosing $m_0 = m_0(N, s, g)$, we can assume

$$E \subset B_2. \quad (3.38)$$

For every $r > 0$ we define the non-negative and monotone non-decreasing absolutely continuous function

$$w(r) := |B_r \setminus E|.$$

Thanks to the monotonicity and the hypotheses we observe that for every $0 \leq r \leq 1$

$$w(r) \leq w(1) \leq C(N, s, g)\varepsilon. \quad (3.39)$$

We observe that $0 \in \{r \in [0, 1] : w(r) = 0\}$ and we let

$$r_2 := \sup\{r \in [0, 1] : w(r) = 0\} \in [0, 1]. \quad (3.40)$$

Our goal is to prove that

$$1 - r_2 \leq C(N, s, g)\varepsilon^{s/N}. \quad (3.41)$$

Without loss of generality we suppose that $r_2 < 1$, otherwise (3.41) trivialises itself, and we fix $r > 0$ such that $r_2 < r < 1$. By definition of r_2 we have that $|E \cup B_r| > |E|$.

We consider the following competitor (with volume equal to $|E| = |B_1|$) for the volume constrained problem,

$$G := \mu(E \cup B_r),$$

with

$$\mu = \mu(r) := \left(\frac{|E|}{|E \cup B_r|}\right)^{\frac{1}{N}} < 1.$$

After a computation, thanks to (3.39), we observe that μ is arbitrarily close to 1 (upon choosing $\varepsilon_0(N, s, g)$ small enough). In particular (upon choosing $\varepsilon_0(N, s, g)$ small enough) we get

$$0 < 1 - \mu \leq C(N)w(r) \quad (3.42)$$

and, since $r < 1$,

$$F \subseteq B_{\mu r} \subseteq B_1.$$

At this point we make use of the minimality of E (see (3.7)) to write

$$\begin{aligned} & P_s(E) + \int_E g_m(x) \, dx \\ & \leq P_s(G) + \int_G g_m(x) \, dx = \mu^{N-s} P_s(E \cup B_r) + \mu^N \int_{E \cup B_r} g_m(\mu x) \, dx \end{aligned}$$

Now, we recall [10], Lemma 2.1 and making use of the isoperimetric inequality for P_s , the invariance under complementation of P_s and the fact that $E^c = (B_r \setminus E) \sqcup (B_r \cup E)^c$ we obtain

$$\begin{aligned} & C(N, s)(w(r))^{\frac{N-s}{N}} \leq P_s(B_r \setminus E) \\ & = P_s(E^c) - P_s((E \cup B_r)^c) + 2 \int_{B_r \setminus E} \int_{(E \cup B_r)^c} \frac{1}{|x-y|^{N+s}} \, dx dy \\ & \leq (\mu^{N-s} - 1)P_s(E \cup B_r) \end{aligned} \tag{3.43}$$

$$+ \mu^N \int_{E \cup B_r} g_m(\mu x) \, dx - \int_E g_m(x) \, dx \tag{3.44}$$

$$+ 2 \int_{B_r \setminus E} \int_{(E \cup B_r)^c} \frac{1}{|x-y|^{N+s}} \, dx dy. \tag{3.45}$$

We first observe that (3.43) is negative.

We then estimate (3.44) using the fact that g is locally Lipschitz (see (H2)) and non-negative (see (2.2)) together with (3.38), (3.42), (3.22) and the fact that $r < 2$:

$$\begin{aligned} & \mu^N \int_{E \cup B_r} g_m(\mu x) \, dx - \int_E g_m(x) \, dx \\ & = (\mu^N - 1) \int_{E \cup B_r} g_m(\mu x) \, dx + \int_E (g_m(\mu x) - g_m(x)) \, dx + \int_{B_r \setminus E} g_m(\mu x) \, dx \\ & \leq (1 - \mu)2 \operatorname{Lip}(g; B_{2+R_0}) + w(r) \sup_{B_{1+R_0}} g \\ & \leq C(N, s, g)w(r). \end{aligned}$$

Now we deal with (3.45). Following the scheme of the proof of [8], equation (15), we write

$$\begin{aligned} & \int_{B_r \setminus E} \int_{(E \cup B_r)^c} \frac{1}{|x-y|^{N+s}} \, dx dy \leq \int_{B_r \setminus E} \int_{B_r^c} \frac{1}{|x-y|^{N+s}} \, dx dy \\ & \leq \int_{B_r \setminus E} \int_{\mathbb{R}^N \setminus B_{r-|y|}(y)} \frac{1}{|x-y|^{N+s}} \, dx dy \leq \int_{B_r \setminus E} \int_{\mathbb{R}^N \setminus B_{r-|y|}} \frac{1}{|z|^{N+s}} \, dz dy \\ & \leq N\omega_N \int_{B_r \setminus E} \int_{r-|y|}^{+\infty} \frac{1}{\varrho^{1+s}} \, d\varrho dy = \frac{N\omega_N}{s} \int_{B_r \setminus E} \frac{1}{(r-|y|)^s} \, dy \\ & = \frac{N\omega_N}{s} \int_0^r \frac{1}{(r-t)^s} \mathcal{H}^{N-1}(\partial B_t \setminus E) \, dt \\ & = \frac{N\omega_N}{s} \int_0^r \frac{w'(t)}{(r-t)^s} \, dt. \end{aligned}$$

Gathering all the previous inequalities together, we find that for every r s.t. $r_2 < r < 1$,

$$C(N, s)(w(r))^{\frac{N-s}{N}} \leq C(N, s, g)w(r) + \frac{N\omega_N}{s} \int_0^r \frac{w'(t)}{(r-t)^s} dt. \quad (3.46)$$

Now, recalling (3.39), upon choosing $\varepsilon_0(N, s, g)$ smaller, we can rewrite (3.46) as

$$C(N, s, g)(w(r))^{\frac{N-s}{N}} \leq \frac{N\omega_N}{s} \int_0^r \frac{w'(t)}{(r-t)^s} dt. \quad (3.47)$$

Now we integrate both sides of (3.47) with respect to the variable r and apply Theorem 3.5 part (ii) to obtain that for some $c = c(N, s, g) > 0$

$$2c \int_{r_2}^\varrho w(r)^{\frac{N-s}{N}} dr \leq w(\varrho), \quad (3.48)$$

for every $r_2 < \varrho < 1$.

Recalling (3.39), upon choosing $C(N, s, g)$ larger and $\varepsilon_0(N, s, g)$ smaller, we can assume

$$1 - \frac{(2w(1))^{\frac{s}{N}} N}{sc} \geq 1 - C(N, s, g)\varepsilon^{\frac{s}{N}} > 0. \quad (3.49)$$

Without loss of generality we can also assume that

$$1 - \frac{(2w(1))^{\frac{s}{N}} N}{sc} > r_2, \quad (3.50)$$

otherwise (3.49) immediately implies (3.41).

(3.48) and (3.50) allow us to invoke Theorem 3.6 part (ii) to deduce that

$$w\left(1 - \frac{(2w(1))^{\frac{s}{N}} N}{sc}\right) = 0. \quad (3.51)$$

From (3.51), recalling (3.49) and (3.40), we obtain that $r_2 \geq 1 - \frac{(2w(1))^{\frac{s}{N}} N}{sc} \geq 1 - C(N, s, g)\varepsilon^{\frac{s}{N}}$. \square

4. UNCONSTRAINED MINIMIZATION PROBLEM

Now we show that our minimization problem (1.1) is equivalent to the rescaled and unconstrained problem

$$\min \left\{ \mathcal{E}_{s, g_m}^\mu(E) := P_s(E) + \int_E g_m(x) dx + \mu||E| - |B_1|| : E \subset \mathbb{R}^N \right\}$$

for $\mu > 0$ sufficiently large. Here we recall that g_m is defined by (3.6), that is,

$$g_m(x) := \sigma(m)^s g(\sigma(m)x + x_m)$$

for $x \in \mathbb{R}^N$ where $\sigma(m)$ is defined by (1.4) and x_m is given in Theorem 3.4.

Proposition 4.1 (Uniform version of Lemma 3.4 in [8]). *Let $s \in (0, 1)$ and assume that g satisfies (H1), (H2) and (2.2). For every $m > 0$ let E_m be a minimizer of $\mathcal{E}_{s,g}$ with $|E_m| = m$ and let \tilde{E}_m be as in (3.5). There exist $m_0 = m_0(N, s, g) > 0$ and $\mu_0 = \mu_0(N, s, g) > 0$ such that if $0 < m \leq m_0(N, s, g)$, then,*

$$\tilde{E}_m \subset B_{3/2},$$

and

$$\tilde{E}_m \in \arg \min \left\{ P_s(F) + \int_F g_m(x) \, dx + \mu \|F\| - |B_1| : F \subset B_3 \right\}, \quad (4.1)$$

for every $\mu \geq \mu_0$, where g_m is defined by (3.6).

Proof. We revisit the proof of [8], Lemma 3.4 making use of the extra information given by Theorem 3.7 and tracking carefully all the constants.

In this proof $c = c(N, s, g) > 0$, $c' = c'(N, s, g) > 0$ are fixed constants defined by,

$$\begin{aligned} c &= P_s(B_1) + (\|g\|_{L^\infty(B_{3+R_0})})2|B_3|, \\ c' &= \left[2c \frac{(N-s)}{N\omega_N} + \text{Lip}(g; B_{6+R_0}) \frac{2 \text{diam}(B_3)}{N} \right] (1 + \varepsilon_1)^N + \|g\|_{L^\infty(B_{3+R_0})}, \end{aligned}$$

where $R_0 = R_0(N, s, g)$ is as in the statement of Proposition 3.4.

Let $\varepsilon_1 = \varepsilon_1(N, s) \in (0, 1/2)$ be a small constant such that

$$\left| \frac{\lambda - 1}{\lambda^N - 1} \right| \leq \frac{2}{N} \quad \left| \frac{\lambda^{N-s} - 1}{\lambda^N - 1} \right| \leq \frac{2(N-s)}{N} \quad (4.2)$$

for all $\lambda > 0$ with $|\lambda - 1| \leq \varepsilon_1$.

Let $\varepsilon_2 = \varepsilon_2(N, s) > 0$ be a constant such that $\varepsilon_2 < \min \left\{ 1 - \frac{1}{(1+\varepsilon_1)^N}, \frac{1}{(1-\varepsilon_1)^N} - 1 \right\}$.

Let $\mu_0 = \mu_0(N, s, g) > 0$ be a constant such that $\mu_0 > \max \left\{ \frac{c}{\varepsilon_2 |B_1|}, c' \right\}$.

We fix $m_0 = m_0(N, s, g) \in (0, 1)$ as in the statement of Theorem 3.7 (or smaller) and m such that $0 < m \leq m_0(N, s, g)$.

We fix $\mu \geq \mu_0$ and we assume by contradiction that $\exists G_\mu \subseteq B_3$ of finite s -perimeter such that

$$P_s(G_\mu) + \int_{G_\mu} g_m(x) \, dx + \mu \left| |G_\mu| - |B_1| \right| < P_s(\tilde{E}_m) + \int_{\tilde{E}_m} g_m(x) \, dx \quad (4.3)$$

Then, since \tilde{E}_m solves (3.7),

$$\begin{aligned} P_s(G_\mu) + \mu \left| |G_\mu| - |B_1| \right| &\leq P_s(\tilde{E}_m) + \int_{\tilde{E}_m} g_m(x) \, dx - \int_{G_\mu} g_m(x) \, dx \leq \\ &\leq P_s(B_1) + \int_{B_1} g_m(x) \, dx - \int_{G_\mu} g_m(x) \, dx \leq \\ &\leq P_s(B_1) + (\|g_m\|_{L^\infty(B_3)}) |G_\mu \triangle B_1| \leq P_s(B_1) + (\|g\|_{L^\infty(B_{3+R_0})}) 2|B_3| = c. \end{aligned}$$

Which, implies

$$P_s(G_\mu) \leq c \qquad \|G_\mu\| - |B_1| \leq \frac{c}{\mu} \leq \frac{c}{\mu_0} \leq \varepsilon_2 |B_1|.$$

In particular, $\frac{|B_1|}{(1+\varepsilon_1)^N} \leq |G_\mu| \leq \frac{|B_1|}{(1-\varepsilon_1)^N}$. We observe that $|G_\mu| \neq |B_1|$, since \tilde{E}_m solves (3.7) and (4.3) holds.

At this point let $\tilde{G}_\mu := \lambda_\mu G_\mu$, with $\lambda_\mu := (|B_1|/|G_\mu|)^{1/N}$. Being the volume of \tilde{G}_μ equal to $|B_1|$, by minimality of \tilde{E}_m for (3.7), we get:

$$P_s(\tilde{E}_m) + \int_{\tilde{E}_m} g_m(x) \, dx \leq P_s(\tilde{G}_\mu) + \int_{\tilde{G}_\mu} g_m(x) \, dx. \quad (4.4)$$

The combination between (4.3) and (4.4) leads to:

$$\mu \leq \frac{P_s(\tilde{G}_\mu) - P_s(G_\mu)}{\|G_\mu\| - |B_1|} + \frac{\int_{\tilde{G}_\mu} g_m(x) \, dx - \int_{G_\mu} g_m(x) \, dx}{\|G_\mu\| - |B_1|} \quad (4.5)$$

Applying exactly the same argument in the proof of [8], Lemma 3.4, we obtain:

$$\frac{P_s(\tilde{G}_\mu) - P_s(G_\mu)}{\|G_\mu\| - |B_1|} = \frac{(\lambda_\mu^{N-s} - 1)P_s(G_\mu)}{|\lambda_\mu^N - 1||G_\mu|}, \quad (4.6)$$

and

$$\begin{aligned} & \frac{\int_{\tilde{G}_\mu} g_m(x) \, dx - \int_{G_\mu} g_m(x) \, dx}{\|G_\mu\| - |B_1|} \\ &= \frac{\int_{\tilde{G}_\mu} g_m(x) \, dx - \lambda_\mu^N \int_{G_\mu} g_m(x) \, dx}{\|G_\mu\| - |B_1|} + \frac{(\lambda_\mu^N - 1) \int_{G_\mu} g_m(x) \, dx}{\|G_\mu\| - |B_1|} \\ &\leq \frac{|B_1| \operatorname{Lip}(g; B_{6+R_0}) |\lambda_\mu - 1| \operatorname{diam}(B_3)}{|\lambda_\mu^N - 1||G_\mu|} + \|g\|_{L^\infty(B_{3+R_0})}. \end{aligned} \quad (4.7)$$

By (4.2), (4.5), (4.6), (4.7) we deduce that

$$\mu \leq c' < \mu_0,$$

that is the desired contradiction. □

5. REGULARITY AND CONVEXITY OF MINIMIZERS

In this section, our goal is to show the following lemma:

Lemma 5.1. *Let $s \in (0, 1)$ and assume that g satisfies (H1), (H2) and (2.2). For every $m > 0$ let E_m be a minimizer of $\mathcal{E}_{s,g}$ with $|E_m| = m$ and let \tilde{E}_m be as in (3.5). Then, there exists $\tilde{m}_0 > 0$, which depends only on N , s and g , such that, for any $m \in (0, \tilde{m}_0)$ the boundary of \tilde{E}_m is of class $C^{2,\alpha}$ for any $\alpha \in (0, s)$. Moreover, $\partial\tilde{E}_m$ converges to ∂B_1 in C^2 -sense, in particular, \tilde{E}_m is convex for any $m \in (0, \tilde{m}_0)$.*

We remark that Lemma 5.1 actually implies that the latter part of Theorem 1.1 is valid. Therefore, combining Lemma 5.1 with Theorem 3.7, we conclude the proof of our main theorem, that is, Theorem 1.1.

In order to prove Lemma 5.1, we exploit the regularity results proved by Caputo and Guillen [6]; Figalli, Fusco, Maggi, Millot, and Morini [11]; Savin and Valdinoci [22]; and Barrios, Figalli, and Valdinoci [4]. Before recalling the regularity results, we first give the definition of the so called “almost” minimizers or Λ -minimizers for the s -fractional perimeter P_s based on the definition given by Figalli, Fusco, Maggi, Millot, and Morini [11] or Caputo and Guillen [6].

Definition 5.2 ((Λ, δ) -Minimizers or Almost Minimizers of P_s). Let $s \in (0, 1)$, $\Lambda > 0$ and $\delta > 0$. We say that a measurable set $E \subset \mathbb{R}^N$ is a (Λ, δ) -*minimizer* of P_s if

$$P_s(E; B_r(x)) \leq P_s(F; B_r(x)) + \Lambda |E \Delta F| \quad (5.1)$$

for any measurable set $F \subset \mathbb{R}^N$, $x \in \partial E$, and $r \in (0, \delta)$ with $E \Delta F \subset B_r(x)$.

Note that the similar concept of the almost minimality for P_s , which is more general, was also given by Caputo and Guillen [6]. See [6] for the details.

Next we recall the regularity result on the almost minimizers or (Λ, δ) -minimizers of P_s shown by Figalli, Fusco, Maggi, Millot, and Morini [11], Corollary 3.5 (see also [6]). This result is a nonlocal analogue of the theory of Tamanini on almost minimal surfaces.

Theorem 5.3 (cf. Corollary 3.5 in [11]). *If $N \geq 2$, $\Lambda > 0$, and $s_0 \in (0, 1)$, then there exist $0 < \varepsilon_0 < 1$, $C_0 > 0$, and $\alpha \in (0, 1)$, depending on n , Λ , and s_0 only, with the following property: if E is a (Λ, δ) -minimizer of P_s for $s \in [s_0, 1)$ in the sense of Definition 5.2 and*

$$0 \in \partial E, \quad \partial E \cap B_1(0) \subset \{y \in \mathbb{R}^N : |y \cdot e| < \varepsilon_0\} \quad (5.2)$$

for some $e \in \mathbb{S}^{N-1}$, then $\partial E \cap B_{1/2}(0)$ is of class $C^{1,\alpha}$ and the $C^{1,\alpha}$ -norm of its graph function is bounded by C_0 .

In other words, Theorem 5.3 with [6], Theorem 1.2 implies that the boundary of an almost minimizer of P_s is of class $C^{1,\alpha}$ with some $\alpha \in (0, 1)$ outside of a closed set of Hausdorff dimension at most $N - 2$.

Next we recall the regularity result of fractional minimal cones in \mathbb{R}^2 by Savin and Valdinoci [22].

Theorem 5.4 ([22]). *Assume that $E \subset \mathbb{R}^2$ is an s -fractional minimal cone, namely, E is a minimizer of P_s in any ball and satisfies that $E = tE$ for any $t > 0$. Then E is a half-plane.*

In particular, by combining the blow-up and blow-down arguments in [5], one may obtain that s -fractional minimal surfaces in \mathbb{R}^2 are fully $C^{1,\alpha}$ -regular for any $\alpha \in (0, s)$.

Corollary 5.5 ([22]). *If E is an s -fractional minimal set in an open set $\Omega \subset \mathbb{R}^2$, then $\partial E \cap \Omega'$ is a $C^{1,\alpha}$ -curve for any $\Omega' \Subset \Omega$.*

Originally, the regularity of nonlocal (fractional) minimal surfaces, which are defined by the boundaries of sets minimizing the fractional perimeter, was obtained by Caffarelli, Roquejoffre, and Savin [5]. Precisely, they proved that every fractional minimal surface is locally $C^{1,\alpha}$ outside of a closed singular set of Hausdorff dimension at most $N - 2$. Moreover, thanks to Theorem 5.5, it follows that the singular set of fractional minimal surfaces has Hausdorff dimension at most $N - 3$.

Now we prove that the rescaled minimizers for our problem are (Λ, δ) -minimizers of P_s where Λ is independent of the minimizers.

Lemma 5.6 ((Λ, δ) -Minimality or Almost Minimality). *Let $s \in (0, 1)$ and $\delta \in (0, 1)$. Assume that g satisfies (H1), (H2) and (2.2). For every $m > 0$, let E_m be a minimizer of $\mathcal{E}_{s,g}$ with $|E_m| = m$ and let \tilde{E}_m be as in*

(3.5). Then there exist $m_0 = m_0(N, s, g) > 0$ and $\Lambda = \Lambda(N, s, g)$ such that, for every $m \in (0, m_0)$, \tilde{E}_m is a (Λ, δ) -minimizer of P_s in the sense of Definition 5.2.

Proof. We fix $\delta \in (0, 1)$. From Proposition 4.1, we know that \tilde{E}_m is also a solution to the problem (4.1), for some large constant $\mu > 0$ independent of m and $\tilde{E}_m \subset B_{3/2}$. Now, letting $F \subset \mathbb{R}^N$, $x \in \partial\tilde{E}_m$, and $r \in (0, \delta)$ with $\tilde{E}_m \triangle F \subset B_r(x)$, from the minimality of \tilde{E}_m , we have that

$$\mathcal{E}_{s, g_m}^\mu(\tilde{E}_m) \leq \mathcal{E}_{s, g_m}^\mu(F), \quad (5.3)$$

since $F \subset B_3$. Hence, from (5.3), we can compute as follows:

$$\begin{aligned} & P_s(\tilde{E}_m) - P_s(F) \\ &= \mathcal{E}_{s, g_m}^\mu(\tilde{E}_m) - \int_{\tilde{E}_m} g_m(x) \, dx - \mu \left| |\tilde{E}_m| - |B_1| \right| - \mathcal{E}_{s, g_m}^\mu(F) + \int_F g_m(x) \, dx + \mu \left| |F| - |B_1| \right| \\ &\leq \int_{\mathbb{R}^N} |\chi_{\tilde{E}_m} - \chi_F| g_m(x) \, dx + \mu \left| |F| - |\tilde{E}_m| \right| \\ &\leq \int_{\tilde{E}_m \triangle F} g_m(x) \, dx + \mu |\tilde{E}_m \triangle F|. \end{aligned} \quad (5.4)$$

Moreover, from the fact that $\tilde{E}_m \subset B_{3/2}$ we deduce that $\tilde{E}_m \triangle F \subset B_r(x) \subset B_3$.

Therefore, by Proposition 3.4 and the definition of g_m , we have

$$\int_{\tilde{E}_m \triangle F} g_m(x) \, dx \leq \sigma(m)^s \|g\|_{L^\infty(B_R(0))} |\tilde{E}_m \triangle F|, \quad (5.5)$$

where $R > 0$ is a constant independent of m . Hence, from (5.4) and (5.5), we obtain

$$P_s(\tilde{E}_m) \leq P_s(F) + (\sigma(m_0)^s \|g\|_{L^\infty(B_R(0))} + \mu) |\tilde{E}_m \triangle F|$$

for any $F \subset \mathbb{R}^N$, $x \in \partial\tilde{E}_m$, and $r \in (0, \delta)$ with $\tilde{E}_m \triangle F \subset B_r(x)$ where $m_0 > 0$ is as in Proposition 4.1. This implies that \tilde{E}_m is an almost minimizer or (Λ, δ) -minimizer of P_s in the sense of (5.1) with $\Lambda := \sigma(m_0)^s \|g\|_{L^\infty(B_R(0))} + \mu < \infty$, which is independent of minimizers. \square

As a consequence of Lemma 5.6 and the aforementioned regularity results for the almost minimizers of the s -fractional perimeter, we obtain

Lemma 5.7 ($C^{1, \alpha}$ -regularity for Minimizers of \mathcal{E}_{s, g_m}). *Let $s \in (0, 1)$ and assume that g satisfies (H1), (H2) and (2.2). For every $m > 0$ let E_m be a minimizer of $\mathcal{E}_{s, g}$ with $|E_m| = m$ and let \tilde{E}_m be as in (3.5). There exists $m_0 = m_0(N, s, g) > 0$ s.t. if $m \in (0, m_0)$, then the boundary of \tilde{E}_m is of class $C^{1, \alpha}$ for some $\alpha \in (0, 1)$ outside of a closed set of Hausdorff dimension at most $N - 3$.*

Proof. The idea is to apply Theorem 5.3 to our problem. To this aim, it is sufficient to show that every minimizer \tilde{E}_m is indeed a (Λ, δ) -minimizer in the sense of (5.1) for some $\Lambda > 0$ depending on N, s and g only, which is guaranteed by Lemma 5.6.

Therefore, from Theorem 5.3 (see also [11], Cor. 3.5) and the regularity of minimal cones in [22], we obtain that the boundary of \tilde{E}_m is of class $C^{1, \alpha}$ for some $\alpha \in (0, 1)$ outside of a closed set of Hausdorff dimension at most $N - 3$. \square

From the uniform closeness of minimizers to the Euclidean ball stated in Theorem 3.7, we can immediately obtain the Hausdorff convergence of the minimizers to the unit ball.

Then, with the Hausdorff convergence of the minimizers at our disposal, we can exploit the regularity criterion (5.2) and the smoothness of the limit set B_1 via the argument in [20], Theorem 26.6 in order to obtain the following result (see also [11], Thm. 3.3 and Cor. 3.6).

Lemma 5.8. *Let $s \in (0, 1)$. Assume that g satisfies (H1), (H2) and (2.2). For any $m > 0$, let E_m be a minimizer of $\mathcal{E}_{s,g}$ with $|E_m| = m$ and let \tilde{E}_m be as in (3.5). Then, there exist a constant $m_0 > 0$ and a bounded sequence of functions $\{\varphi_m\}_{m \in (0, m_0)} \subset C^{1,\alpha}(\partial B_1)$ with some $\alpha \in (0, 1)$ (independent of m) such that, for any $m \in (0, m_0)$,*

$$\partial \tilde{E}_m = \{(1 + \varphi_m(x))x : x \in \partial B_1(0)\} \quad \text{and} \quad \lim_{m \downarrow 0} \|\varphi_m\|_{C^1(\partial B_1(0))} = 0. \quad (5.6)$$

Proof. First of all, we recall that, from Theorem 3.7, there exist positive constants $m_0 = m_0(N, s, g)$ and $C = C(N, s, g)$ such that, if $m \in (0, m_0)$, \tilde{E}_m satisfies the inclusion

$$B_{1-r_0}(0) \subset \tilde{E}_m \subset B_{1+r_0}(0) \quad (5.7)$$

for some $r_0 \in (0, Cm^{s^2/(2N^2)})$. Thus, (5.7) implies that \tilde{E}_m converges to $B_1(0)$ as $m \downarrow 0$ in L^1 -sense and moreover, by definition, $\partial \tilde{E}_m$ converges to $\partial B_1(0)$ in the Hausdorff distance.

Now, as we observe in Lemma 5.6 and Lemma 5.7, we obtain that \tilde{E}_m is also an almost minimizer of P_s and that $\partial \tilde{E}_m$ is of class $C^{1,\alpha}$ with some $\alpha \in (0, 1)$ for $m \in (0, m_0)$ outside of a closed set of Hausdorff dimension at most $N - 3$. Although the singular sets can appear on each $\partial \tilde{E}_m$, by combining the convergence of $\{\partial \tilde{E}_m\}_{m \in (0, m_0)}$ to ∂B_1 in the Hausdorff distance with the regularity criterion (5.2), we can show that there exists a constant $m'_0 > 0$ such that, for each $m \in (0, m'_0)$, there exists a function $\varphi_m \in C^{1,\alpha}(\partial B_1)$ such that

$$\partial \tilde{E}_m = \{(1 + \varphi_m(x))x : x \in \partial B_1\}.$$

Therefore, the representation of $\partial \tilde{E}_m$ in (5.6) holds. We refer to [11], Corollary 3.6 for the similar argument.

Finally, we show the convergence of the functions $\{\varphi_m\}_{m \in (0, m'_0)}$ in C^1 sense as in (5.6) by using the argument in [20], Theorem 26.6. Let $z_m \in \partial \tilde{E}_m$ be any point and let $R \in (0, 1)$. By the Hausdorff convergence of $\{\partial \tilde{E}_m\}_{m \in (0, m'_0)}$, we can choose a subsequence (denoted by the same indices) such that $z_m \rightarrow z_0 \in \partial B_1(0)$ as $m \downarrow 0$. We define the cylinder $C(x, R) \subset \mathbb{R}^N$ centred at $x = (x', x_N) \in \mathbb{R}^{N-1} \times \mathbb{R}$ by

$$\{(y', y_N) \in \mathbb{R}^{N-1} \times \mathbb{R} : |y' - x'| < R, \quad |y_N - x_N| < R\}.$$

Then, from the regularity of each $\partial \tilde{E}_m$ and $\partial B_1(0)$, for each $m \in (0, m'_0)$, we can choose a $C^{1,\alpha}$ -function $u_m : B'_R(z'_0) \rightarrow \mathbb{R}$ and a smooth function $u_0 : B'_R(z'_0) \rightarrow \mathbb{R}$ such that $\text{Lip}(u_m) \leq 1$ for any m and

$$\begin{aligned} C(z_0, R) \cap \tilde{E}_m &= \{(x', x_N) : -R < x_N < u_m(x')\} \\ C(z_0, R) \cap B_1(0) &= \{(x', x_N) : -R < x_N < u_0(x')\} \end{aligned}$$

where $z_0 := (z'_0, z_{0,N}) \in \mathbb{R}^{N-1} \times \mathbb{R}$ and $B'_R(x')$ is the $(N - 1)$ -dimensional ball of radius R centred at $x' \in \mathbb{R}^{N-1}$. From the regularity criterion (5.2) and the same argument as in, for instance, [20], Theorem 26.3, we further obtain the Hölder continuity of the function $\nabla' u_m$, that is,

$$|\nabla' u_m(y') - \nabla' u_m(z')| \leq C |y' - z'|^\alpha < +\infty \quad (5.8)$$

for any $y', z' \in B'_R(z'_0)$ and some constant $C > 0$ independent of m , where ∇' is the gradient in \mathbb{R}^{N-1} and $\alpha \in (0, 1)$ is as in Lemma 5.7. Moreover, thanks to the convergence $\tilde{E}_m \rightarrow B_1(0)$ in L^1 -sense, we have

$$\int_{B'_R(z'_0)} |u_m(z') - u_0(z')| dz' = |\tilde{E}_m \triangle E \cap C(z_0, R)| \xrightarrow{m \downarrow 0} 0.$$

Thus, we obtain

$$\int_{B'_R(z'_0)} \xi(z') \nabla' u_m(z') dz' \xrightarrow{m \downarrow 0} \int_{B'_R(z'_0)} \xi(z') \nabla' u_0(z') dz' \quad (5.9)$$

for any $\xi \in C_c^\infty(B'_R(z_0))$. From (5.8), we have that $\{\nabla' u_m\}_{m \in (0, m'_0)}$ is equi-continuous and bounded in $C^0(B'_R)$, and thus, by the Ascoli-Arzelà theorem, we can extract a subsequence (denoted by the same indices) such that $\nabla' u_m \rightarrow v$ as $m \downarrow 0$ uniformly in $B'_R(z'_0)$. From (5.9), the limit v coincides with $\nabla' u_0$. Therefore, we obtain that the graph functions $\{u_m\}_{m \in (0, m'_0)}$ of $\partial \tilde{E}_m$ converge to the graph function u_0 of ∂B_1 in C^1 sense. From the choice of φ_m , this implies that $\|\varphi_m\|_{C^1(\partial B_1)}$ converges to 0. Therefore, we conclude the proof of (5.6). \square

Now we improve the regularity of the boundary of minimizers of \mathcal{E}_{s, g_m} by employing the regularity result for solutions to integro-differential equations via the bootstrap argument. This result was obtained by Barrios, Figalli, and Valdinoci [4], Theorem 1.6. They proved the following regularity theorem on the solutions to integro-differential equations. For simplicity, we do not describe all of the theorem here. See [4], Theorem 1.6 for the full statement.

Theorem 5.9. *Let $s \in (0, 1)$, $\beta \in (0, 1]$, and $r > 0$. Let $v \in L^\infty(\mathbb{R}^{N-1})$ be a solution (in the viscosity sense) to the integro-differential equation*

$$\int_{\mathbb{R}^{N-1}} L_r(x', y') (v(x' + y') + v(x' - y') - 2v(x')) dy' = F(x', v(x'))$$

for any $x' \in B'_r(0) \subset \mathbb{R}^{N-1}$ where L_r satisfies the following assumptions:

(A1) *There exist constants $a_0, r_0 > 0$ and $\eta \in (0, \frac{a_0}{4})$ such that*

$$\frac{(1-s)(a_0 - \eta)}{|y'|^{N+s}} \leq L_r(x', y') \leq \frac{(1-s)(a_0 + \eta)}{|y'|^{N+s}}$$

for any $x' \in B'_r(0)$ and $y' \in B'_{r_0}(0) \setminus \{0\}$;

(A2) *$L_r \in C^{0, \beta}(B'_r(0) \times (\mathbb{R}^{N-1} \setminus \{0\}))$ and there exists a constant $C_0 > 0$ such that*

$$\|L_r(\cdot, y')\|_{C^{0, \beta}(B'_r)} \leq \frac{C_0}{|y'|^{N+s}}$$

for any $y' \in \mathbb{R}^{N-1} \setminus \{0\}$;

and $F \in C^{0, \beta}(B'_r(0) \times \mathbb{R})$. Then, if η is sufficiently small, then $v \in C^{1+s+\alpha}(B'_{r/2}(0))$ for any $\alpha < \beta$. Moreover, we have the estimate

$$\|v\|_{C^{1+s+\alpha}(B'_{r/2}(0))} \leq C \left(1 + \|v\|_{L^\infty(\mathbb{R}^{N-1})} + \|F\|_{L^\infty(B'_r(0) \times \mathbb{R})}\right),$$

where $C > 0$ is a constant depending on N, s, C_0 , and $\|F\|_{C^{0, \beta}(B'_r(0) \times \mathbb{R})}$.

Taking into account all of the above arguments, we can obtain that the boundary of minimizers of \mathcal{E}_{s,g_m} with the volume $|B_1|$ has $C^{2,\beta}$ -regularity for any $\beta \in (0, s)$. Precisely, we prove

Lemma 5.10 (Improved Regularity of Minimizers). *Let $s \in (0, 1)$ and $m_0 > 0$ be as in Lemma 5.8. Assume that g satisfies (H1), (H2) and (2.2). Then, if E_m is a minimizer of $\mathcal{E}_{s,g}$ with $|E_m| = m$ for any $m \in (0, m_0)$, $\partial\tilde{E}_m$ is of class $C^{2,\beta}$ for any $\beta \in (0, s)$, where \tilde{E}_m is defined as in (3.5).*

Proof. We may assume that $0 \in \partial\tilde{E}_m$ by translation. By Lemma 5.8, we can represent $\partial\tilde{E}_m \cap (B'_r(0) \times (-r, r)) \subset \mathbb{R}^{N-1} \times \mathbb{R}$ for $r > 0$ as a graph of a $C^{1,\alpha}$ -function u for some $\alpha \in (0, 1)$ by choosing the proper coordinate. From the fact that \tilde{E}_m is a minimizer of \mathcal{E}_{s,g_m} with volume constraint and by employing the computation shown in [4], we may obtain the following Euler–Lagrange equation in the viscosity sense:

$$\begin{aligned} & \int_{\mathbb{R}^{N-1}} L_r(x', y')(u(x' + y') + u(x' - y') - 2u(x')) dy' \\ & = G(x', u(x')) + \tilde{\lambda}_m - g_m(x', u(x')) \quad \text{for } x' \in B'_{r'/2}(0) \subset \mathbb{R}^{N-1} \end{aligned}$$

for $0 < r' < r$, where L_r satisfies (A1) and (A2) in Theorem 5.9, $\text{spt } L_r(x', \cdot) \subset B'_{r'/2}(0)$ for any $x' \in B'_{r'/2}(0)$, G is some smooth function (see [4], Sect. 3 for the precise expression), and $\tilde{\lambda}_m$ is a Lagrange multiplier. Then, since the potential g is locally Lipschitz, we now apply Theorem 5.9 several times, if necessary, to conclude that the regularity of u can be improved up to $C^{2,\beta}$ with $\beta \in (0, s)$. From the compactness of $\partial\tilde{E}_m$ and by a simple covering argument, we obtain the $C^{2,\beta}$ -regularity of $\partial\tilde{E}_m$ for any $\beta \in (0, s)$ with the following estimate

$$\|u\|_{C^{2,\beta}(B'_{r'/4})} \leq C \left(1 + \|u\|_{L^\infty(B'_r)} + |\tilde{\lambda}_m| + \|g_m\|_{L^\infty(B_{2R_0})} \right),$$

where $C > 0$ is a constant depending only on N, s , and $\|g\|_{C^{0,1}(B_{2R_0})}$ and R_0 is as in Proposition 3.4. \square

Finally, we are ready to prove Lemma 5.1, that is, the main lemma in this section.

Proof of Lemma 5.1. Taking into account all the arguments in Proposition 3.4, Theorem 3.7, Lemma 5.8, and Lemma 5.10, we obtain that there exist a constant $m_0 = m_0(N, s, g) > 0$ and a sequence of functions $\{\varphi_m\}_{m \in (0, m_0)} \subset C^{2,\alpha}(\partial B_1(0))$ for any $\alpha \in (0, s)$ such that

$$\partial\tilde{E}_m = \{(1 + \varphi_m(x))x : x \in \partial B_1\}, \|\varphi_m\|_{L^\infty(\partial B_1)} \leq C_1 m^{\frac{s^2}{2N^2}}, \lim_{m \downarrow 0} \|\varphi_m\|_{C^1(\partial B_1)} = 0 \quad (5.10)$$

where $C_1 > 0$ is a constant depending only on N, s , and g . Moreover, from the proof of Lemma 5.10 and (5.10), we can obtain

$$\|\varphi_m\|_{C^{2,\alpha}(\partial B_1)} \leq C_2(1 + |\tilde{\lambda}_m|) \quad (5.11)$$

for any $m \in (0, m_0)$, by choosing $m_0 > 0$ small if necessary, where $C_2 > 0$ is a constant depending only on N, s , and g and $\tilde{\lambda}_m$ is a Lagrange multiplier as in Lemma 5.10.

Now we derive the upper bound of $|\tilde{\lambda}_m|$ from the minimality of \tilde{E}_m . Since \tilde{E}_m is a minimizer of \mathcal{E}_{s,g_m} with $|\tilde{E}_m| = |B_1|$, by considering the Euler–Lagrange equation with volume constraint, we have

$$\frac{d}{dt} \Big|_{t=0} \left(P_s(\Phi_t(\tilde{E}_m)) + \int_{\Phi_t(\tilde{E}_m)} g_m \right) = \tilde{\lambda}_m \frac{d}{dt} \Big|_{t=0} |\Phi_t(\tilde{E}_m)| \quad (5.12)$$

where $\{\Phi_t\}_{|t|<1}$ is one-parameter diffeomorphism associated with $T \in C_c^\infty(\mathbb{R}^N; \mathbb{R}^N)$, that is, $\Phi_t(x) := x + tT(x)$ for any $x \in \mathbb{R}^N$ and $|t| < 1$. Then, substituting the identity function $\text{Id} : x \mapsto x$ with T (by using a suitable cut-off function or by approximation), we obtain, from (5.12),

$$N|B_1|\tilde{\lambda}_m = (N-s)P_s(\tilde{E}_m) + \int_{\tilde{E}_m} \nabla g_m(x) \cdot x \, dx + N \int_{\tilde{E}_m} g_m(x) \, dx. \quad (5.13)$$

From Theorem 3.7 and the assumption on g , we obtain

$$\left| \int_{\tilde{E}_m} \nabla g_m(x) \cdot x \, dx + N \int_{\tilde{E}_m} g_m(x) \, dx \right| \leq C(N, s, g) m^{\frac{s}{N}} \quad (5.14)$$

for any $m \in (0, m_0)$ where $C = C(N, s, g) > 0$ is a constant and R_0 is given in Proposition 3.4 depending only on N, s , and g . Then, from (5.13) and (5.14) and by the minimality of \tilde{E}_m , we obtain

$$|\tilde{\lambda}_m| \leq (N|B_1|)^{-1} ((N-s)P_s(B_1) + C(N, s, g)m^{\frac{s}{N}}) \quad (5.15)$$

for any $m \in (0, m_0)$. Therefore, from (5.11) and (5.15), we obtain

$$\|\varphi_m\|_{C^{2,\alpha}(\partial B_1)} \leq C_3(N, s, g) < \infty \quad (5.16)$$

for any $m \in (0, m_0)$ and some constant $C_3 > 0$ independent of m .

Next, because of the continuous embedding $C^{2,\alpha}(\bar{\Omega}) \subset C^{2,\beta}(\bar{\Omega}) \subset C^0(\bar{\Omega})$ for any smooth bounded open set Ω and $\beta < \alpha$, we may observe that the following interpolation inequality is valid (see, for instance, [19], Cor. 1.2.19 and Cor. 1.2.7): let $\alpha \in (0, 1)$. Then, for any $\beta \in (0, \alpha)$, there exist $\vartheta(\alpha, \beta) \in (0, 1)$ and a constant $C = C(N, \alpha, \beta) > 0$ such that

$$\|u\|_{C^{2,\beta}} \leq C \|u\|_{C^0}^{\vartheta(\alpha,\beta)} \|u\|_{C^{2,\alpha}}^{1-\vartheta(\alpha,\beta)}$$

for any $u \in C^{2,\alpha}$. From this interpolation inequality, we deduce that, for $\beta \in (0, \alpha)$, there exist constants $\vartheta(\alpha, \beta) \in (0, 1)$ and $\tilde{C}_3(N, \alpha, \beta) > 0$ such that

$$\|\varphi_m\|_{C^{2,\beta}(\partial B_1(0))} \leq C_3(N, \alpha, \beta) \|\varphi_m\|_{L^\infty(\partial B_1(0))}^{\vartheta(\alpha,\beta)} \|\varphi_m\|_{C^{2,\alpha}(\partial B_1(0))}^{1-\vartheta(\alpha,\beta)}$$

for $m \in (0, m_0)$. Therefore, from (5.10) and (5.16), we finally obtain that there exist constants $\tilde{m}_0 = \tilde{m}_0(N, s, g) > 0$ and $C_4(N, s, g, \alpha) > 0$ such that

$$\|\varphi_m\|_{C^2(\partial B_1(0))} \leq C_4(N, s, g, \alpha) m^{\frac{\vartheta(\alpha)s^2}{2N^2}} \quad (5.17)$$

for any $m \in (0, \tilde{m}_0)$. This implies that $\partial \tilde{E}_m$ is close to $\partial B_1(0)$ in C^2 -sense and, in particular, \tilde{E}_m is convex for sufficiently small $m > 0$ because the second fundamental form of $\partial \tilde{E}_m$ converges pointwise to that of $\partial B_1(0)$ which is positive definite. \square

Remark 5.11. In the proof of Lemma 5.1, we have observed the convexity of minimizers for small volumes, assuming that g is locally Lipschitz, coercive, and $\inf_{\mathbb{R}^N} g = g(0) = 0$.

In the classical problem (1.3), assuming that g is convex and coercive, convexity of minimizers for large volumes holds true by [7] (see also [9], Rem. 1.6). However, we do not know if such property still holds in our problem due to the lack of the analogue of Alvarez-Lasry-Lions theory [1, 18] in the nonlocal setting.

In addition, if we further assume that g is monotone non-decreasing and radially symmetric, that is, there exists a monotone non-decreasing function $G : [0, \infty) \rightarrow \mathbb{R}$ such that $g(x) = G(|x|)$ for any $x \in \mathbb{R}^N$, then we can show that the unit ball B_1 is a minimizer of \mathcal{E}_{s,g_m} with volume $|B_1|$ for all $m > 0$.

Indeed, from the isoperimetric inequality for P_s and a direct computation, we have

$$P_s(E^*) \leq P_s(E), \quad (5.18)$$

$$\int_{E^*} g_m(x) dx \leq \int_E g_m(x) dx, \quad (5.19)$$

for any measurable set $E \subset \mathbb{R}^N$ with finite measure, where we denote by E^* the open ball of radius equal to $|E|^{1/N}|B_1|^{-1/N}$ centred at the origin.

Furthermore, if G is strictly increasing, we observe that (5.19) becomes an equality if and only if $E = E^*$. Therefore, since the equality in (5.18) is attained if and only if $E = E^*$ (up to translations) in this case we conclude that the unit ball B_1 is the unique minimizer of \mathcal{E}_{s,g_m} with volume $|B_1|$ for all $m > 0$.

Corollary 5.12. *Under the assumptions of Theorem 5.1, for every $m > 0$ the following Euler–Lagrange equation holds:*

$$H_{s,\partial E_m} + g = \lambda_m \quad \text{on } \partial E_m, \quad (5.20)$$

where λ_m is a Lagrange multiplier associated to the volume constraint such that

$$\lim_{m \rightarrow 0} \lambda_m = +\infty. \quad (5.21)$$

Proof. By making the same computation as in (5.13), we can show that the identity

$$Nm\lambda_m = (N-s)P_s(E_m) + \int_{E_m} \nabla g(x) \cdot x dx + N \int_{E_m} g(x) dx \quad (5.22)$$

holds for $m > 0$ and that $\lambda_m \rightarrow +\infty$ as $m \downarrow 0$. □

Remark 5.13 (H_s -bubble problem). The classical H -bubble problem, which was originally raised by S.-T. Yau [23], Problem 59, is formulated as follows: given a function $H : \mathbb{R}^N \rightarrow \mathbb{R}$, the question is to find an immersed hypersurface $\mathcal{M} \subset \mathbb{R}^N$ such that its mean curvature at $p \in \mathcal{M}$ is equal to $H(p)$. (1.1) is related to a nonlocal version of this geometric problem. Using the same notation as in Lemma 5.1 and Corollary 5.12, we observe that if the function $(0, +\infty) \ni m \mapsto \lambda_m$ were continuous in a neighbourhood of zero, this would imply that it would be also surjective when restricted to a neighbourhood of zero (thanks to (5.21)). As a consequence, there would exist $\bar{\lambda} > 0$ such that we would be able to solve a fractional version of the H -bubble problem, that is: we might find a hypersurface ∂E satisfying the equation

$$H_{s,\partial E} + g = \lambda \quad \text{on } \partial E,$$

for every $\lambda > \bar{\lambda}$.

Regarding a sufficient condition for the continuity of the Lagrange multiplier λ_m , we remark that, if for $m_0 > 0$ sufficiently small a set E_{m_0} with $|E_{m_0}| = m_0 > 0$ is the unique minimizer of $\mathcal{E}_{s,g}$ among sets with volume m_0 , then the Lagrange multiplier λ_m is continuous in a neighbourhood of zero (with respect to $m > 0$).

Indeed, to prove this we first take any sequence $\{m_i\}_i \subset (0, \infty)$ converging to a fixed $m_0 \in (0, \infty)$. We observe, from the minimality of E_{m_i} , that $\sup_i P_s(E_{m_i}) < \infty$. Then, from the compactness of P_s , the uniqueness

of E_{m_0} , and the coercivity of g , we deduce that $E_{m_i} \rightarrow E_{m_0}$ in L^1 -sense. Thus, from the minimality of E_{m_0} , the uniform boundedness of E_{m_i} and by using the dominated convergence theorem, we have

$$\int_{E_{m_i}} g(x) \, dx \xrightarrow{i \rightarrow \infty} \int_{E_{m_0}} g(x) \, dx. \quad (5.23)$$

From the minimality of E_{m_i} , we can also obtain the continuity of P_s at m_0 , that is,

$$P_s(E_{m_i}) \xrightarrow{i \rightarrow \infty} P_s(E_{m_0}).$$

Indeed, from the minimality of E_{m_i} , setting $F_{m_i} := \left(\frac{m_i}{m_0}\right)^{1/N} E_{m_0}$, we have

$$\begin{aligned} E_{s,g}[m_i] &:= \inf\{\mathcal{E}_{s,g}(E) \mid |E| = m_i\} \\ &\leq \mathcal{E}_{s,g}(F_{m_i}) \\ &= \left(\frac{m_i}{m_0}\right)^{\frac{N-s}{N}} P_s(E_{m_0}) + \frac{m_i}{m_0} \int_{E_{m_0}} g\left(\left(\frac{m_i}{m_0}\right)^{\frac{1}{N}} x\right) \, dx \end{aligned}$$

for any $i \in \mathbb{N}$. Thus, from the minimality of E_{m_0} , we obtain that

$$\limsup_{i \rightarrow \infty} E_{s,g}[m_i] \leq E_{s,g}[m_0].$$

In the same way, we can also obtain that $\liminf_{i \rightarrow \infty} E_{s,g}[m_i] \geq E_{s,g}[m_0]$. From this, we derive the continuity of $E_{s,g}[m]$ at m_0 . Therefore, from the minimality of E_{m_i} and E_{m_0} and from (5.23), we obtain that P_s is continuous at m_0 .

Finally, from the assumption on g and the uniform boundedness of E_{m_i} , we can also obtain that the potential term in (5.22) is continuous at m_0 , that is,

$$\int_{E_{m_i}} \nabla g(x) \cdot x \, dx + N \int_{E_{m_i}} g(x) \, dx \xrightarrow{i \rightarrow \infty} \int_{E_{m_0}} \nabla g(x) \cdot x \, dx + N \int_{E_{m_0}} g(x) \, dx.$$

In conclusion, from (5.22), we obtain the continuity of λ_m .

Acknowledgements. K.B. and M.N. are members of INdAM-GNAMPA. The work of KB is partially supported by the INdAM-GNAMPA 2023 Project *Problemi variazionali per funzionali e operatori non-locali*, codice CUP_E53C22001930-001, and by the PRIN Project 2022J4FYNJ. The research of KB was also supported by the INdAM-GNAMPA 2022 Project *Fenomeni non locali in problemi locali*, codice CUP_E55F22000270001, and by the PRIN Project 2017BTM7SN. The work of F.O. was supported by the DFG Collaborative Research Center TRR 109, ‘‘Discretization in Geometry and Dynamics’’.

REFERENCES

- [1] O. Alvarez, J.-M. Lasry and P.-L. Lions, Convex viscosity solutions and state constraints. *J. Math. Pures Appl.* **76** (1997) 265–288.
- [2] J.E. Avron, J.E. Taylor and R.K.P. Zia, Equilibrium shapes of crystals in a gravitational field: crystals on a table. *J. Statist. Phys.* **33** (1983) 493–522.
- [3] E. Baer, Minimizers of anisotropic surface tensions under gravity: higher dimensions via symmetrization. *Arch. Ration. Mech. Anal.* **215** (2015) 531–578.
- [4] B. Barrios, A. Figalli and E. Valdinoci, Bootstrap regularity for integro-differential operators and its application to nonlocal minimal surfaces. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **13** (2014) 609–639.

- [5] L. Caffarelli, J.-M. Roquejoffre and O. Savin, Nonlocal minimal surfaces. *Commun. Pure Appl. Math.* **63** (2010) 1111–1144.
- [6] M.C. Caputo and N. Guillen, Regularity for non-local almost minimal boundaries and applications. arXiv (2010).
- [7] V. Caselles and A. Chambolle, Anisotropic curvature-driven flow of convex sets. *Nonlinear Anal.* **65** (2006) 1547–1577.
- [8] A. Cesaroni and M. Novaga, Volume constrained minimizers of the fractional perimeter with a potential energy. *Discrete Contin. Dyn. Syst. Ser. S* **10** (2017) 715–727.
- [9] G. De Philippis and M. Goldman, A two-point function approach to connectedness of drops in convex potentials. *Commun. Anal. Geom.* **30** (2022) 815–841.
- [10] A. Di Castro, M. Novaga, B. Ruffini and E. Valdinoci, Nonlocal quantitative isoperimetric inequalities. *Calc. Var. Partial Diff. Equ.* **54** (2464) 2421–2464.
- [11] A. Figalli, N. Fusco, F. Maggi, V. Millot and M. Morini, Isoperimetry and stability properties of balls with respect to nonlocal energies. *Commun. Math. Phys.* **336** (2015) 441–507.
- [12] A. Figalli and F. Maggi, On the shape of liquid drops and crystals in the small mass regime. *Arch. Ration. Mech. Anal.* **201** (2011) 143–207.
- [13] R. Finn, The sessile liquid drop. I. Symmetric case. *Pacific J. Math.* **88** (1980) 541–587.
- [14] R. Finn, Equilibrium capillary surfaces, Vol. 284 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York (1986).
- [15] E. Gonzalez, U. Massari and I. Tamanini, Existence and regularity for the problem of a pendent liquid drop. *Pacific J. Math.* **88** (1980) 399–420.
- [16] E. Indrei, On the equilibrium shape of a crystal. arXiv (2020).
- [17] E. Indrei and A. Karakhanyan, Minimizing the free energy. arXiv (2023).
- [18] N. Korevaar, Capillary surface convexity above convex domains. *Indiana Univ. Math. J.* **32** (1983) 73–81.
- [19] A. Lunardi, Analytic semigroups and optimal regularity in parabolic problems. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel (1995).
- [20] F. Maggi, Sets of finite perimeter and geometric variational problems, Vol. 135 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge (2012).
- [21] R.J. McCann, Equilibrium shapes for planar crystals in an external field. *Commun. Math. Phys.* **195** (1998) 699–723.
- [22] O. Savin and E. Valdinoci, Regularity of nonlocal minimal cones in dimension 2. *Calc. Var. Part. Diff. Equ.* **48** (2013) 33–39.
- [23] S.-T. Yau, Problem section, in *Seminar on Differential Geometry*, Vol. 102 of *Ann. of Math. Stud.* Princeton University Press, Princeton, N.J. (1982) 669–706.



Please help to maintain this journal in open access!

This journal is currently published in open access under the Subscribe to Open model (S2O). We are thankful to our subscribers and supporters for making it possible to publish this journal in open access in the current year, free of charge for authors and readers.

Check with your library that it subscribes to the journal, or consider making a personal donation to the S2O programme by contacting subscribers@edpsciences.org.

More information, including a list of supporters and financial transparency reports, is available at <https://edpsciences.org/en/subscribe-to-open-s2o>.